

Evidence for resonant structures in $e^+e^- \rightarrow \pi^+\pi^-h_c^*$

YUAN Chang-Zheng(苑长征)

Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China

Abstract: The cross sections of $e^+e^- \rightarrow \pi^+\pi^-h_c$ at center-of-mass energies from 3.90 to 4.42 GeV were measured by the BESIII and the CLEO-c experiments. Resonant structures are evident in the $e^+e^- \rightarrow \pi^+\pi^-h_c$ line shape. The fit to the line shape results in a narrow structure at a mass of (4216 ± 18) MeV/ c^2 and a width of (39 ± 32) MeV, and a possible wide structure of mass (4293 ± 9) MeV/ c^2 and width (222 ± 67) MeV. Here, the errors are combined statistical and systematic errors. This may indicate that the Y(4260) state observed in $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ has a fine structure in it.

Key words: charmonium, charmoniumlike states, e^+e^- annihilation, Exotics, Hybrids

PACS: 14.40.Rt, 14.40.Pq, 13.66.Bc **DOI:** 10.1088/1674-1137/38/4/043001

The observation of the Y-states in the exclusive production of $\pi^+\pi^-J/\psi$ [1–4] and $\pi^+\pi^-\psi(3686)$ [5–7] from the B-factories is a great puzzle in understanding vector charmonium states [8]. According to the potential models, there are five vector states above the well-known 1D state $\psi(3770)$ and below around 4.7 GeV/ c^2 ; namely, the 3S, 2D, 4S, 3D, and 5S states [8]. However, experimentally, besides the three well known structures observed in inclusive hadronic cross section (i.e., the $\psi(4040)$, $\psi(4160)$, and $\psi(4415)$ [9]) there are four Y-states (i.e., the Y(4008), Y(4260), Y(4360), and Y(4660)) [1–7]. This suggests that at least some of these structures are not charmonium states and, therefore, a number of scenarios have arisen for interpreting one or more of these structures [8].

The BESIII experiment [10] running near the open charm threshold supplies further information to understand the properties of these vector states. Amongst this information, the most relevant measurement is the study of $e^+e^- \rightarrow \pi^+\pi^-h_c$ [11]. Besides the observation of a charged charmoniumlike state $Z_c(4020)$, BESIII has reported the cross section measurement of $e^+e^- \rightarrow \pi^+\pi^-h_c$ at 13 center-of-mass (CM) energies, from 3.900 to 4.420 GeV [11]. The measurements are listed in Table 1. In the studies, the h_c is reconstructed via its electric-dipole (E1) transition $h_c \rightarrow \gamma\eta_c$ with η_c to 16 exclusive hadronic final states: $p\bar{p}$, $2(\pi^+\pi^-)$, $2(K^+K^-)$, $K^+K^-\pi^+\pi^-$, $p\bar{p}\pi^+\pi^-$, $3(\pi^+\pi^-)$, $K^+K^-2(\pi^+\pi^-)$, $K_S^0K^\pm\pi^\mp$, $K_S^0K^\pm\pi^\mp\pi^\pm\pi^\mp$, $K^+K^-\pi^0$,

$p\bar{p}\pi^0$, $\pi^+\pi^-\eta$, $K^+K^-\eta$, $2(\pi^+\pi^-)\eta$, $\pi^+\pi^-\pi^0\pi^0$, and $2(\pi^+\pi^-)\pi^0\pi^0$.

Table 1. $e^+e^- \rightarrow \pi^+\pi^-h_c$ cross sections measured from the BESIII experiment. For the first three energy points, besides the upper limits, the central values and the statistical errors that will be used in the fits below are also listed. The second errors are systematic errors and the third errors are from the uncertainty in $\mathcal{B}(h_c \rightarrow \gamma\eta_c)$ [12].

\sqrt{s}/GeV	$\sigma(e^+e^- \rightarrow \pi^+\pi^-h_c)/\text{pb}$
3.900	0.0 ± 6.0 or < 8.3
4.009	1.9 ± 1.9 or < 5.0
4.090	0.0 ± 7.4 or < 13
4.190	$17.7 \pm 9.8 \pm 1.6 \pm 2.8$
4.210	$34.8 \pm 9.5 \pm 3.2 \pm 5.5$
4.220	$41.9 \pm 10.7 \pm 3.8 \pm 6.6$
4.230	$50.2 \pm 2.7 \pm 4.6 \pm 7.9$
4.245	$32.7 \pm 10.3 \pm 3.0 \pm 5.1$
4.260	$41.0 \pm 2.8 \pm 3.7 \pm 6.4$
4.310	$61.9 \pm 12.9 \pm 5.6 \pm 9.7$
4.360	$52.3 \pm 3.7 \pm 4.8 \pm 8.2$
4.390	$41.8 \pm 10.8 \pm 3.8 \pm 6.6$
4.420	$49.4 \pm 12.4 \pm 4.5 \pm 7.6$

The CLEO-c experiment did a similar analysis, but with significant signal only at CM energy 4.17 GeV [13], the result is $\sigma = (15.6 \pm 2.3 \pm 1.9 \pm 3.0)$ pb, where the third error is from the uncertainty in $\mathcal{B}[\psi(3686) \rightarrow \pi^0 h_c]$.

Received 24 December 2013

* Supported by Ministry of Science and Technology of China (2009CB825203) and National Natural Science Foundation of China (NSFC) (10825524, 10935008, 11235011)



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Article funded by SCOAP³ and published under licence by the Chinese Physical Society, the Institute of High Energy Physics of the Chinese Academy of Sciences, the Institute of Modern Physics of the Chinese Academy of Sciences, and IOP Publishing Ltd

The cross sections are of the same order of magnitude as those of the $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ measured by BESIII [14] and other experiments [3, 4], but with a different line shape (see Fig. 1). There is a broad structure at high energy with a possible local maximum at around 4.23 GeV. We try to use the BESIII and the CLEO-c measurements to extract the resonant structures in $e^+e^- \rightarrow \pi^+\pi^-h_c$.




Fig. 1. A comparison between the cross sections of $e^+e^- \rightarrow \pi^+\pi^-h_c$ from BESIII (dots with error bars) [11] and those of $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ from Belle (open circles with error bars) [4]. The errors are statistical only.

Since the systematic error ($\pm 18.1\%$) of the BESIII experiment is common for all the data points, we only use the statistical errors in the fits below. The CLEO-c measurement is completely independent from the BESIII experiment and all of the errors added in quadrature (± 4.2 pb) are taken as the total error, which is used in the fits. We use a least χ^2 method with [15]

$$\chi^2 = \sum_{i=1}^{14} \frac{(\sigma_i^{\text{meas}} - \sigma^{\text{fit}}(m_i))^2}{(\Delta\sigma_i^{\text{meas}})^2},$$

where $\sigma_i^{\text{meas}} \pm \Delta\sigma_i^{\text{meas}}$ is the experimental measurement, and $\sigma^{\text{fit}}(m_i)$ is the cross section value calculated from the model below with the parameters from the fit. Here, m_i is the energy corresponds to the i th energy point.

Since the line shape above 4.42 GeV is unknown, it is not clear whether or not the large cross section at high energy will decrease. We will try to fit the data with two different scenarios.

Assuming that the cross section follows the three-body phase space and that there is a narrow resonance at around 4.2 GeV, we fit the cross sections with the coherent sum of two amplitudes, a constant and a constant width relativistic Breit-Wigner (BW) function; that is,

$$\sigma(m) = |c \cdot \sqrt{PS(m)} + e^{i\phi} BW(m) \sqrt{PS(m)}/PS(M)|^2,$$

where $PS(m)$ is the 3-body phase space factor,

$$BW(m) = \frac{\sqrt{12\pi\Gamma_{e^+e^-}\mathcal{B}(\pi^+\pi^-h_c)\Gamma_{\text{tot}}}}{m^2 - M^2 + iM\Gamma_{\text{tot}}},$$

is the Breit-Wigner (BW) function for a vector state, with mass M , total width Γ_{tot} , electron partial width $\Gamma_{e^+e^-}$, and the branching fraction to $\pi^+\pi^-h_c$, $\mathcal{B}(\pi^+\pi^-h_c)$, keep in mind that from the fit we can only extract the product $\Gamma_{e^+e^-}\mathcal{B}(\pi^+\pi^-h_c)$. The constant term c and the relative phase, ϕ , between the two amplitudes are also free parameters in the fit, together with the resonant parameters of the BW function.

The fit indicates the existence of a resonance (called Y(4220) hereafter) with a mass of (4216 ± 7) MeV/ c^2 and a width of (39 ± 17) MeV, and the goodness-of-fit is $\chi^2/\text{ndf} = 11.04/9$, corresponding to a confidence level of 27%. There are two solutions for the $\Gamma_{e^+e^-} \times \mathcal{B}(Y(4220) \rightarrow \pi^+\pi^-h_c)$, which are (3.2 ± 1.5) eV and (6.0 ± 2.4) eV. Here, all of the errors are from the fit only. Fitting the cross sections without the Y(4220) results in a very bad fit, $\chi^2/\text{ndf} = 72.75/13$, corresponding to a confidence level of 2.5×10^{-10} . The statistical significance of the Y(4220) is calculated to be 7.1σ comparing




Fig. 2. The fit to the cross sections of $e^+e^- \rightarrow \pi^+\pi^-h_c$ from BESIII and CLEO-c (dots with error bars). The solid curves show the best fits, and the dashed ones are individual components. (a) is the fit with the coherent sum of a phase space amplitude and a BW function, and (b) is the coherent sum of two BW functions.

the two χ^2 s obtained above and taking into account the change of the number-of-degree-of-freedom. Fig. 2(a) shows the final fit with the Y(4220).

Assuming that the cross section decreases at high energy, we fit the cross sections with the coherent sum of two constant width relativistic BW functions; that is,

$$\sigma(m) = |BW_1(m) \cdot \sqrt{PS(m)/PS(M_1)} + e^{i\phi} BW_2(m) \cdot \sqrt{PS(m)/PS(M_2)}|^2,$$

where both BW_1 and BW_2 take the same form as $BW(m)$ above but with different resonant parameters.

The fit indicates the existence of the Y(4220) with a mass of (4230 ± 10) MeV/ c^2 and a width of (12 ± 36) MeV, as well as a broad resonance, the Y(4290), with a mass of (4293 ± 9) MeV/ c^2 and width of (222 ± 67) MeV. The goodness-of-the-fit is $\chi^2/\text{ndf} = 1.81/7$, corresponding to a confidence level of 97%, which is an almost perfect fit. There are two solutions for the $\Gamma_{e^+e^-} \times \mathcal{B}[Y(4220)/Y(4290) \rightarrow \pi^+\pi^-h_c]$, which are (0.07 ± 0.07) eV/ (16.1 ± 2.2) eV and (2.7 ± 4.9) eV/ (19.0 ± 5.9) eV. Again, here the errors are from fit only. Fitting the cross sections without the Y(4220) results in a much worse fit, $\chi^2/\text{ndf} = 30.65/11$, corresponding to a confidence level of 1.3×10^{-3} . The statistical significance of the Y(4220) is calculated to be 4.5σ , comparing the two χ^2 s obtained above and taking into account the change of the number-of-degree-of-freedom. Fig. 2(b) shows the final fit with the Y(4220) and Y(4290).

From the two fits shown above, we conclude that it is very likely that there is a narrow structure at around 4.22 GeV/ c^2 , although we are not sure if there is a broad resonance at 4.29 GeV/ c^2 . We try to average the results from the fits to give the best estimation of the resonant parameters. For the Y(4220), we obtain

$$\begin{aligned} M(Y(4220)) &= (4216 \pm 18) \text{ MeV}/c^2, \\ \Gamma_{\text{tot}}(Y(4220)) &= (39 \pm 32) \text{ MeV}, \\ \Gamma_{e^+e^-}^{Y(4220)} \times \mathcal{B}[Y(4220) \rightarrow \pi^+\pi^-h_c] &= (4.6 \pm 4.6) \text{ eV}. \end{aligned}$$

While for the Y(4290), we obtain

$$\begin{aligned} M(Y(4290)) &= (4293 \pm 9) \text{ MeV}/c^2, \\ \Gamma_{\text{tot}}(Y(4290)) &= (222 \pm 67) \text{ MeV}, \\ \Gamma_{e^+e^-}^{Y(4290)} \times \mathcal{B}[Y(4290) \rightarrow \pi^+\pi^-h_c] &= (18 \pm 8) \text{ eV}. \end{aligned}$$

Here, the errors include both statistical and systematic

errors. The results from the two solutions and the two fit scenarios are covered by enlarged errors, the common systematic error in the cross section measurement is included in the error of the $\Gamma_{e^+e^-}$.

It is noticed that the uncertainties of the resonant parameters of the Y(4220) are large, this is due to two important facts: one is the lack of data at CM energies above 4.42 GeV, which may discriminate which of the two above scenarios is correct, the other is the lack of high precision measurements around the Y(4220) peak, especially between 4.23 and 4.26 GeV. The two-fold ambiguity in the fits is a natural consequence of the coherent sum of two amplitudes [16]. Although high precision data will not resolve the problem, they will reduce the errors in $\Gamma_{e^+e^-}$ from the above fits. Since the fit with a phase space amplitude predicts rapidly increasing cross section at high energy, it is very unlikely to be true, so the results from the fit with two resonances is more likely to be true. More measurements from the BESIII experiments at CM energies above 4.42 GeV and more precise data at around the Y(4220) peak will also be crucial to settle down all of these problems.

There are thresholds of $\bar{D}D_1$ [17], $\omega\chi_{cJ}$ [18, 19], $D_s^{*+}D_s^{*-}$ [9] at the Y(4220) mass region, which make the identification of the nature of this structure very complicated. The fits described in this paper supply only one possibility of interpreting the data. In Ref. [20], the BESIII measurements [11] were described with the presence of one relative S -wave $\bar{D}D_1 + c.c.$ molecular state Y(4260) and a non-resonant background term; while in Ref. [21], the BESIII data [11] were fitted with a model where the Y(4260) and Y(4360) are interpreted as the mixture of two hadrocharmonium states. It is worth pointing out that various QCD calculations indicate that the charmonium-hybrid lies in the mass region of these two Y states [22] and the $c\bar{c}$ tend to be in a spin-singlet state. Such a state may couple to a spin-singlet charmonium state such as h_c strongly, this makes the Y(4220) and/or Y(4290) good candidates for the charmonium-hybrid states.

In summary, we fit $e^+e^- \rightarrow \pi^+\pi^-h_c$ cross sections measured by BESIII and CLEO-c experiments. Evidence for a narrow structure at around 4.22 GeV, as well as a wide one at 4.29 GeV, is observed. More high precision measurements at above 4.42 GeV and around 4.22 GeV are desired to better understand these structures.

References

- 1 Aubert B et al. (BaBar collaboration). Phys. Rev. Lett., 2005, **95**: 142001
- 2 YUAN C Z et al. (Belle collaboration). Phys. Rev. Lett., 2007, **99**: 182004
- 3 Lees J P et al. (BaBar collaboration). Phys. Rev. D, 2012, **86**: 051102(R)
- 4 LIU Z Q et al. (Belle collaboration). Phys. Rev. Lett., 2013, **110**: 252002
- 5 Aubert B et al. (BaBar collaboration). Phys. Rev. Lett., 2007, **98**: 212001
- 6 WANG X L et al. (Belle collaboration). Phys. Rev. Lett., 2007, **99**: 142002
- 7 Aubert B et al. (BaBar collaboration). arXiv:1211.6271
- 8 Brambilla N et al. Eur. Phys. J. C, 2011, **71**: 1534
- 9 Beringer J et al. (Particle Data Group). Phys. Rev. D, 2012, **86**: 010001
- 10 Ablikim M et al. (BESIII collaboration). Nucl. Instrum. Methods Phys. Res. Sect. A, 2010, **614**: 345
- 11 Ablikim M et al. (BESIII collaboration). Phys. Rev. Lett., 2013, **111**: 242001
- 12 Ablikim M et al. (BESIII collaboration). Phys. Rev. Lett., 2010, **104**: 132002
- 13 Pedlar T K et al. (CLEO collaboration). Phys. Rev. Lett., 2011, **107**: 041803
- 14 Ablikim M et al. (BESIII collaboration). Phys. Rev. Lett., 2013, **110**: 252001
- 15 For the three low statistics energy points, the χ^2 is not well defined. We take the central values listed in Table 1 as nominal values, and vary the central values and statistical errors in a wide range to estimate the possible bias in this assumption. The bias is found to be small and is considered as systematic error of the results
- 16 ZHU K, MO X H, YUAN C Z, WANG P. Int. J. Mod. Phys. A, 2011, **26**: 4511
- 17 WANG Q, Hanhart C, ZHAO Q. Phys. Rev. Lett., 2013, **111**: 132003
- 18 DAI L Y, SHI M, TANG G Y, ZHENG H Q. arXiv:1206.6911 [hep-ph]
- 19 YUAN C Z, WANG P, MO X H. Phys. Lett. B, 2006, **634**: 399
- 20 Cleven M, WANG Q, GUO F K, Hanhart C, Meißner U G, ZHAO Q. arXiv:1310.2190 [hep-ph]
- 21 LI X, Voloshin M B. arXiv:1309.1681 [hep-ph]
- 22 Barnes T, Close F E, Swanson E S. Phys. Rev. D, 1995, **52**: 5242; GUO P, Szczepaniak A P, Galata G, Vassallo A, Santopinto E. Phys. Rev. D, 2008, **78**: 056003; Dudek J J, Rrapaj E. Phys. Rev. D, 2008, **78**: 094504