

# The NUBASE2020 evaluation of nuclear physics properties\*\*

F.G. Kondev<sup>1,\*</sup>, M. Wang (王猛)<sup>2,3,\*</sup>, W.J. Huang (黄文嘉)<sup>2,4,5,6</sup>, S. Naimi<sup>7</sup>, G. Audi (欧乔治)<sup>6</sup>

<sup>1</sup> Physics Division, Argonne National Laboratory, Lemont, IL 60439, USA

<sup>2</sup> Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China

<sup>3</sup> School of Nuclear Science and Technology, University of Chinese Academy of Sciences, Beijing 100049, China

<sup>4</sup> Advanced Energy Science and Technology Guangdong Laboratory, Huizhou 516003, China

<sup>5</sup> Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany

<sup>6</sup> Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France

<sup>7</sup> RIKEN Nishina Center, Wako, Saitama 351-0198, Japan

**Abstract:** The NUBASE2020 evaluation contains the recommended values of the main nuclear physics properties for all nuclei in their ground and excited, isomeric ( $T_{1/2} \geq 100$  ns) states. It encompasses all experimental data published in primary (journal articles) and secondary (mainly laboratory reports and conference proceedings) references, together with the corresponding bibliographical information. In cases where no experimental data were available for a particular nuclide, trends in the behavior of specific properties in neighboring nuclei were examined and estimated values are proposed. Evaluation procedures and policies that were used during the development of this evaluated nuclear data library are presented, together with a detailed table of recommended values and their uncertainties.

**Keywords:** NUBASE2020 evaluation, nuclear properties, atomic masses, isomers, excitation energy of isomers, spin and parity, half-life, year of discovery, decay modes

**DOI:** 10.1088/1674-1137/abddae

## 1 Introduction

NUBASE2020 is an evaluated nuclear data library that contains the recommended values of the main nuclear physics properties: masses, excitation energies (for excited isomers), half-lives, spins and parities and decay modes, and their intensities, for all known nuclei in their ground and excited, isomeric ( $T_{1/2} \geq 100$  ns) states. It also includes information for yet unobserved nuclides that is based on systematic trends of nuclear properties in neighboring nuclei. The present publication includes updated results for these properties, which were reported in previous versions of this library [1–4]. The recommended data are presented in Table I.

The information included in NUBASE2020 represents the fundamental building blocks of the modern nuclear physics, and specifically of the nuclear structure and nuclear astrophysics research. One of the main applications of NUBASE2020 is the “Atomic Mass Evaluation” (AME2020 - the second and third articles included in this issue) where it is imperative to have an unambiguous identification of all states involved in a particular decay, reaction or mass-spectrometry measurement. This is the main reason for coupling the two evaluations together in the present issue. Furthermore, with

the advances of modern mass-spectrometry techniques and the availability of intense stable and rare-isotope beams, a large number of short-lived nuclei can be produced in a single experiment and their masses can be measured with a high precision. Thus, NUBASE2020 can be a trusted source of information in future mass measurements, where an unambiguous identification of specific nuclides in complex mass-spectrometry data would be required.

NUBASE2020 also serves nuclear structure research, astrophysics network calculations, and theoretical studies of nuclear properties, where complete, up-to-date and reliable data for all known nuclei are needed. It can be particularly useful in present and future studies of nuclei and their properties at the major nuclear physics facilities around the world, such as FAIR, ISOLDE and SPIRAL2 (Europe), ATLAS and FRIB (USA), HIAF (China), RIBF at RIKEN (Japan), ISAC and ARIEL (Canada), and elsewhere.

Furthermore, the evaluated data included in NUBASE2020 are a valuable source of information for specialists in a number of applied nuclear fields, such as safeguards, nuclear forensics, reactor engineering, waste management, material analysis, medical diagnostics and radiotherapy, and elsewhere, where one needs to access reliable

Received 19 January 2021, Accepted 20 January 2021, Published online 5 March 2021

\*\*This work has been undertaken with the endorsement of the IUPAP Commission on Symbols, Units, Nomenclature, Atomic Masses and Fundamental Constants (C2).

\* kondev@anl.gov (F.G. Kondev); corresponding author

\* wangm@impcas.ac.cn (M. Wang); corresponding author



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

nuclear physics information for any nuclide.

The recommended data included in NUBASE2020 fulfill several user-demanded requirements, namely that they are: a) *complete* – include all measured quantities and their uncertainties, b) *up-to-date* – include results from all recent publications, c) *credible and reliable* – identify and resolve contradictory results that exist in the scientific literature, as well as in other nuclear physics databases, d) *properly referenced* – provide comprehensive bibliographical information for all included properties.

In general, NUBASE2020 was updated via three different routes: a) directly from the literature by compiling and evaluating data that were published in *primary* (nuclear physics journals) and *secondary* (abstracts, conference proceedings, private communications, theses, arXiv publications and laboratory reports) references; b) by consulting, and when merited by adopting, recommendations made in topical evaluations that include nuclear properties covered by NUBASE (see the text for details); c) by consulting, and when merited by adopting, recommended values from the “Evaluated Nuclear Structure Data File” (ENSDF) database [5].

It is important to note that data presented in ENSDF and in other topical evaluations were carefully examined, and only results that were found to be *valid* and *up-to-date* were considered. In general, the content of ENSDF is very large, since it encompasses the complex nuclear structure and decay properties for all nuclei and all excited nuclear states. Maintenance of this library requires an enormous effort and it is not surprising that occasionally some older data are missing or misrepresented, and that some recent data are not included. When such cases were identified, the data were re-evaluated and the corresponding conclusions were added as comments in Table I.

The content of NUBASE2020, together with the adopted policies that were used during the development of this nuclear physics data library, is described below. All experimental data available to the authors by October 30, 2020 were considered.

## 2 Content of NUBASE2020

The NUBASE2020 evaluation contains recommended properties for the ground state of 3340 nuclides and for 1938 excited isomeric ( $T_{1/2} \geq 100$  ns) states, derived from all available experimental data. It also includes information for yet unobserved nuclei (218 in their ground state and 45 excited isomers) whose properties were estimated by following the systematic trends in neighboring nuclei (TNN, see section 3.1).

For each nuclide and for each state (ground or isomeric), the following properties were compiled and, when necessary, evaluated: mass excess, excitation energy of the excited isomeric state, half-life, spin and parity, decay modes and their intensities, isotopic abundance (for stable nuclides), year of discovery and the corresponding bibliographical information

for all experimental values of the above items.

### 2.1 Mass excess

In general, the knowledge of atomic masses can provide valuable information on the lifetimes of nuclear states and their decay modes, and in particular on the  $\beta$ -delayed particle decay probabilities for nuclei far from the line of stability.

The mass-excess values and their uncertainties that are presented in Table I were adopted from the latest edition of the Atomic Mass Evaluation, AME2020, as described in the second and third articles of the present issue. Figure 1 displays the uncertainties of the mass-excess values as a function of  $N$  and  $Z$ .

### 2.2 Isomers

Nuclear isomers are excited, intrinsic (single-particle in nature) states with lifetimes ranging from nanoseconds (or even shorter) to years. There are several recent compilations and review articles, where the physics of nuclear isomers was discussed in detail and the reader is referred to Refs. [6, 7] and references therein.

Following the NUBASE2003 publication [2], the present evaluation includes isomeric states with half-lives longer than 100 ns. Although this limit is somewhat arbitrary, the main reason for this choice was to include all short-lived isomers that can be directly produced at the present and future accelerator beam facilities and that can survive the time-of-flight path of the employed recoil mass separator, and as a consequence, their decay properties and/or masses can be directly measured.

Figure 2 shows a compilation of all such known isomers as a function of  $N$  and  $Z$ .

Isomers are listed in Table I in the order of increasing excitation energy and they are identified by the letters ‘ $m$ ’, ‘ $n$ ’, ‘ $p$ ’, ‘ $q$ ’, or ‘ $r$ ’ which are appended to the nuclide name, e.g.  $^{90}\text{Nb}$  for the ground state,  $^{90}\text{Nb}^m$  for the first excited isomer,  $^{90}\text{Nb}^n$ ,  $^{90}\text{Nb}^p$ ,  $^{90}\text{Nb}^q$ , and  $^{90}\text{Nb}^r$  for the second, third, fourth and fifth ones, respectively. In four cases, namely  $^{98}\text{Y}$ ,  $^{174}\text{Lu}$ ,  $^{179}\text{Ta}$  and  $^{214}\text{Ra}$ , a sixth isomer is presented and they were labeled with the letter ‘ $x$ ’ (see the Explanation of Table I for details).

The excitation energy of an isomeric states is determined by different experimental methods, which are generally attributed to the category of either *internal* or *external* relations. A typical *internal* relation involves the  $\gamma$ -ray energy, or the energies of a cascade of  $\gamma$  rays, associated with the isomer decay. The most-accurate values for the excitation energies of isomers that are deduced by this approach can be determined by a least-squares fit to the energies of all  $\gamma$  rays observed along the decay path of a particular isomer. In cases where *internal* relations cannot be established, connections to other nuclides (*external* relations) can be used to deduce the mass difference (excitation energy) between the ground state and isomers, and the excitation energies are taken from AME2020.




Fig. 1. Nuclear chart displaying the mass-excess uncertainties for all nuclei in their ground state.




Fig. 2. Nuclear chart displaying isomeric states with  $T_{1/2} \geq 100$  ns. For a given isotope where multiple isomers exist, only the longest-lived state is plotted.

The method used to establish the *external* relation for a particular isomer (the origin) is indicated by a two-letter code in Table I, next to the isomer excitation energy (see the Explanation of Table I for details). For *internal* relations, the origin field is left blank and the numerical values are taken either from ENSDF or from literature updates, where a least-squares fit to the measured  $\gamma$ -ray energies was performed, whenever possible.

There are also cases where the energy difference between the isomer and the ground state can be obtained by both the *internal* and one, or more, *external* relations with comparable accuracies. In such cases, the excitation energy of the isomer is taken from AME2020. For example, the mass of  $^{178}\text{Lu}^m$  is determined in AME2020 at 66% from  $E_x(\text{IT})=120(3)$  keV [1993Bu02] and at 34% from  $^{176}\text{Lu}(\text{t,p})^{178}\text{Lu}^m=4482(5)$  keV [1981Gi01], resulting in an adjusted excitation energy of 123.8(2.6) keV for the isomer.

In contrast to ENSDF, where the isomer excitation energies may not be quantified and are often given as floating levels with ‘+X’, ‘+Y’, etc., estimated values are always provided in NUBASE2020, based on theoretical considerations or TNN. In such cases, the reported excitation energies are considered as a non-experimental quantity and the values are flagged with the symbol ‘#’.

When the existence of an isomer is uncertain and it is still under discussion, it is flagged with ‘EU’ (“Existence Uncertain”) in the origin field. A comment is usually added to indicate why the existence of this state is questioned or where this issue is discussed in more detail. Eleven isomers, namely  $^{138}\text{Pm}^m$ ,  $^{142}\text{Nd}^m$ ,  $^{144}\text{Cs}^n$ ,  $^{152}\text{Pm}^n$ ,  $^{156}\text{Tm}^m$ ,  $^{162}\text{Lu}^n$ ,  $^{174}\text{W}^m$ ,  $^{174}\text{W}^n$ ,  $^{185}\text{Bi}^n$ ,  $^{190}\text{Tl}^n$ , and  $^{273}\text{Ds}^m$  are treated in this way in the present evaluation. Nevertheless, the mass excess and excitation energy values are given for all of them, except for  $^{138}\text{Pm}^m$ ,  $^{142}\text{Nd}^m$ ,  $^{144}\text{Cs}^n$ ,  $^{152}\text{Pm}^n$ ,  $^{174}\text{W}^m$ ,  $^{174}\text{W}^n$ , and  $^{190}\text{Tl}^n$ , where the existence is strongly doubted.

When an isomer was initially reported as “discovered”, but later this was proven to be an error, such a case is flagged with ‘RN’ (“Reported Non-existent”) in the origin field. Nine isomers, namely  $^{76}\text{Cu}^m$ ,  $^{84}\text{Ga}^m$ ,  $^{84}\text{As}^m$ ,  $^{85}\text{Nb}^n$ ,  $^{86}\text{Nb}^n$ ,  $^{117}\text{La}^m$ ,  $^{181}\text{Pb}^m$ ,  $^{196}\text{Pb}^m$ , and  $^{197}\text{Bi}^n$  are treated in this way and no mass excess or excitation energy values are given. Similarly to the ‘EU’ cases, a “non-exist” label is also added. The use of the two flags, ‘EU’ and ‘RN’, was extended to cases where the discovery of a nuclide is questioned (e.g.  $^{260}\text{Fm}$  or  $^{289}\text{Lv}$  or  $^{293}\text{Og}$ ). However, an estimate for the ground state mass, derived from Trends from the Mass Surface (TMS), is always given in AME2020 and NUBASE2020.

Sometimes, upper and lower limits are known for the excitation energy of the isomeric state. Such cases are treated with uniform probability distribution, as explained in section 3.2. For example, there is solid experimental evidence [1974De47] that the excitation energy of the  $^{162}\text{Tm}^m$  isomer is between the 66.9 keV and 192.0 keV levels and this information is presented (after rounding) in Table I as  $E_x =$

130(40) keV.

When it is not clear which state is the ground state and which one is the isomer, the flag ‘\*’ is added in Table I. Similarly, when the uncertainty of the isomer excitation energy,  $\Delta E_x$ , is relatively large compared to  $E_x$ , e.g.  $\Delta E_x > E_x/2$ , the assignment of the level as a ground or isomeric state is also considered to be uncertain and it is flagged with the symbol ‘\*’, as well.

Based on new experimental mass information, the ordering of several ground and excited isomeric states was reversed in the present work, when compared to the recommendations in ENSDF, and such cases are flagged with the symbol ‘&’ in Table I. In a few other instances, evidence was found for a state that is located below the adopted in ENSDF ground state and such results were also flagged with the symbol ‘&’ in Table I. It is worth noting that because of the coupling between NUBASE2020 and AME2020 all changes in the ordering of nuclear levels are firmly established and synchronized.

### 2.2.1 Isobaric analog states

NUBASE2020 contains information for 205 Isobaric Analog States (IAS), which are labeled in Table I with the isospin multiplet value,  $T$ . Their excitation energies were determined via either the “*internal*” or “*external*” relation. The IAS nuclides are generally marked with the  $i$  or  $j$  superscripts, except for eight excited isomers,  $^{16}\text{N}^m$ ,  $^{26}\text{Al}^m$ ,  $^{34}\text{Cl}^m$ ,  $^{38}\text{K}^m$ ,  $^{46}\text{V}^m$ ,  $^{50}\text{Mn}^m$ ,  $^{54}\text{Co}^m$ , and  $^{70}\text{Br}^m$ . The isospin value is not given for most nuclei in their ground state, since they have  $T = |T_z| = \frac{1}{2} |N - Z|$ . However, it is included for the ground state of the  $N = Z$ , odd-odd  $^{34}\text{Cl}$ ,  $^{42}\text{Sc}$ ,  $^{46}\text{V}$ ,  $^{50}\text{Mn}$ ,  $^{54}\text{Co}$ ,  $^{62}\text{Ga}$ ,  $^{66}\text{As}$ ,  $^{70}\text{Br}$ , and  $^{74}\text{Rb}$  ( $T = 1$ ) and  $^{30}\text{P}$ ,  $^{38}\text{K}$ , and  $^{58}\text{Cu}$  ( $T = 0$ ) nuclides.

Detailed experimental information about IAS was recently compiled in Refs. [8, 9].

### 2.3 Half-life

The lifetime is a fundamental property of a nuclear level. It is related to the total decay width,  $\Gamma$ , a linear sum of all partial decay widths ( $\gamma$  ray, conversion electrons,  $\alpha$  decay,  $\beta$  decay, fission, etc.), through the uncertainty relationship:

$$\Gamma = \frac{\hbar}{\tau} \quad (1)$$

where  $\tau = T_{1/2} / \ln(2)$  is the level mean life and  $T_{1/2}$  is the half-life.

Figures 3 displays the ground-state half-life as a function of  $N$  and  $Z$  for all nuclei included in Table I.

Some light nuclei ( $A < 30$ ) that are located beyond the particle drip-lines are known to exist for a very short time before disintegrating by particle emissions. In such cases only the total level width can be measured and the half-life is deduced by means of equation 1 (in convenient units):

$$T_{1/2} [\text{s}] \simeq 4.562 \times 10^{-22} / \Gamma [\text{MeV}] \quad (2)$$

where  $\Gamma$  is the total width in the center of mass frame. The




Fig. 3. Nuclear chart displaying the ground-state half-lives for all nuclei.

heaviest nuclide whose half-life is determined by this approach is  $^{29}\text{Cl}$ .

The following units are used in NUBASE2020 for a convenient display of half-lives: seconds (s) and its sub-units, minutes (m), hours (h), days (d) and years (y), and its sub-units (see Explanation of Table I for details). While several definitions can be used to convert values between years and days or seconds, such as Julian year, Gregorian year, Sidereal year, Tropical year and others, the conversion via Tropical year is adopted in NUBASE2020:

$$1 \text{ y} = 365.2422 \text{ d} = 31556926 \text{ s}$$

When more than one value is known for the half-life of a particular level, a statistical analysis was performed in accordance with the policies outlined in section 3.2. Experimental half-lives are sometimes given in the literature with most probable lower and upper limits. Such cases are treated with uniform probability distribution, as explained in section 3.2. For example, the half-life of  $^{97}\text{In}^m$  is given as  $1.2 \mu\text{s} \leq T_{1/2} \leq 230 \mu\text{s}$  in Ref. [2018Pa20] and the recommended value in NUBASE2020 is  $T_{1/2} = 120(70) \mu\text{s}$ . Half-lives with asymmetric uncertainties are also frequently reported in the literature. Since it is envisioned that NUBASE2020 will be used by specialists in various application fields, such values are symmetrized prior to performing any statistical analyses, as described in section 3.3.

In experiments where rare events were detected, for example in studies of super-heavy nuclei, the half-lives reported in different publications were not directly averaged. Instead, when the information presented in the literature was sufficient, the time information associated with the individual events was combined and analyzed, as prescribed by Schmidt *et al.* [1984Sc13]. In recent review articles that deal with properties of super-heavy nuclei [2014Kh04, 2016Fo10, 2016Ho09, 2017Og01] events from several experiments were combined together in order to determine the best values. We have adopted these half-life values, rather than averaging the individual results.

In cases of long-lived nuclides that are of importance to metrology and other applications, all available experimental data were carefully examined, including values published by various metrology laboratories over many years. As a policy, we adopted the latest reported value by a particular laboratory, including the latest results published by Unterweger and Fitzgerald [10], which superseded the earlier assessment made by the same authors [11].

An upper or a lower limit for the half-life value is given in Table I for nuclides identified using a time-of-flight technique. The following policies were implemented: a) for *observed* nuclides, the lower limit for the half-life is given in the place of the uncertainty field. However, such a value should be used with caution, since it may be far shorter than the actual level half-life. In order to avoid confusion, a somewhat more realistic estimate, derived from TNN and flagged with

#, is also given (see for example the data entry for  $^{44}\text{Si}$ ). The same notation is also used for half-life limits of very long-lived (stable) nuclei (see for example the data entry for  $^{188}\text{Os}$ ); b) for nuclides that were looked for, but *not observed*, the upper limit is given in the place of the uncertainty field. For example, upper limits were estimated for a number of unobserved nuclides by F. Pougheon [1993Po.A], based on the time-of-flight information and the production yields expected from TNN (see for example the data entry for  $^{21}\text{Al}$ ).

In the course of this work it was found that half-lives for double  $\beta$ -decaying nuclides were not always consistently given in ENSDF. Since the two-neutrino ground-state-to-ground-state transition is the dominant decay mode, only those experimental half-life values, or their upper-limits, are included in NUBASE2020. In a few cases, other partial lifetime data are also compiled and these are given as comments in Table I. No attempt was made to convert the half-life values given by different authors to the same statistical confidence level (CL). The compilations by Barabash [2020Ba.A, 2011Ba28] were consulted in covering such decays.

For nuclei in their ground or excited isomeric state whose half-lives were not directly measured, values from TNN were estimates and included in Table I, whenever possible. Such cases are flagged with the symbol ‘#’.

## 2.4 Spin and parity

Spin and parity values are presented with or without parentheses, based on “weak” or “strong” arguments, respectively, as adopted in ENSDF [12], but with one important exception. Since, it is a policy of NUBASE2020 to make a clear distinction between experimental and non-experimental information, parentheses are used only when the so-called “weak” arguments are based on experimental observations. In cases where the assignments are based on theoretical predictions or TNN, the values are presented without parentheses and they are flagged with the symbol ‘#’. This is in contrast to ENSDF, where values determined from theory or systematics are given in parentheses, and as a consequence, it is not possible to distinguish these tentative values from ones determined from experimental data. It should also be noted, that despite well-defined evaluation policies [12], there are a number of inconsistencies in ENSDF regarding the spin and parity assignments. Often, the proposed spins and parities reflect the interpretation of a particular ENSDF evaluator, rather than that of firm policy rules. As a result, assignments to similar states in neighboring nuclei are put in parenthesis by one evaluator, but not by another, although similar experimental information is available.

There is a large amount of recent experimental data on directly measured spins for nuclei far from the line of stability, where the “in-source” (e.g. RILIS at ISOLDE (CERN) and TRILIS at ISAC (TRIUMF)) and “collinear” (e.g. CRIS at ISOLDE (CERN)) laser spectroscopy techniques were de-




Fig. 4. Nuclear chart displaying the measured ground-state spins.






Fig. 5. Nuclear chart displaying the measured isomeric-state spins.

ployed. In the present work, we compiled the experimentally measured spins for 1062 states (827 ground states and 235 isomers) and the corresponding values are flagged in Table I with the symbol ‘\*’. We have consulted previous compilations by Fuller [13], Otten [1989Ot.A] and McDonald [14], as well as recently measured values in the literature. Figures 4 and 5 show plots of the directly measured spins for the ground and isomeric states, respectively, as a function of  $N$  and  $Z$ .

The experimental methods that are used for spin determination do not provide direct information about the parity of a given state. However, we have used additional spectroscopy data, such as  $l$  value in transfer reactions, hindrance factors in  $\alpha$  decay, measured magnetic moments, as well as other spectroscopic information, in order to make such assignments in Table I.

### 2.5 Decay modes and their intensities

Figure 6 displays the main ground-state decay modes for all nuclei included in Table I.

The most important policy in assembling the information about the decay modes and their intensities was to unambiguously establish whether a particular decay is energetically allowed, but not experimentally observed (represented by a question mark alone ‘?’, e.g. ‘IT?’ or ‘ $\alpha$ ?’, the question mark refers to the decay mode), and whether the decay is actually observed, but its intensity is not determined (represented by ‘=?’, e.g. ‘IT=?’ or ‘ $\alpha=?$ ’, the question mark refers to the branching intensity).

In cases of multiple decay modes, normalization of primary intensities to 100% was made only when the competing decays were experimentally observed. Otherwise, no such corrections were made.

Similarly to previous versions of NUBASE [1–4],  $\beta^+$  denotes a decay process that includes both electron capture,  $\varepsilon$ , and positron emission,  $e^+$ , decays, so that one can symbolically write  $\beta^+ = \varepsilon + e^+$ . It should be made clear that this notation is *not* the same as that used in ENSDF, where the combination of both modes is labeled as  $\varepsilon + \beta^+$ . When the available decay energy is below  $2m_e \simeq 1022$  keV, only electron capture decay mode is allowed, while above that value the two processes are in competition. In the latter case, the separated intensities are not always experimentally available and they are frequently deduced from model calculations. Following one of the general policies of NUBASE that experimental information is exclusively used whenever possible, only measured values for  $\beta^+$ ,  $\varepsilon$  and  $e^+$  are included in Table I. By the same token, both electron capture-delayed fission ( $\varepsilon$ SF) and positron-delayed fission ( $e^+$ SF) are given with the same symbol  $\beta^+$ SF.

For  $\beta$ -delayed particle decays, intensity relations were carefully considered. By definition, the intensity of a specific  $\beta$ -delayed particle decay is taken as a percentage of the main  $\beta$ -decay mode. For example, if the decay of the  $(A, Z)$

nuclide is described as ‘ $\beta^- = 100; \beta^- n = 20$ ’, this means that for 100 decays of the parent, 80  $(A, Z+1)$  and 20  $(A-1, Z+1)$  daughter nuclei are produced and that 100 electrons and 20 delayed neutrons are emitted.

This notation also holds for more complex  $\beta$ -delayed particle emissions. For example, a decay described by ‘ $\beta^- = 100; \beta^- n = 30; \beta^- 2n = 20; \beta^- \alpha = 10$ ’ corresponds to the emission of 100 electrons, 70  $(30+2 \times 20)$  delayed-neutrons and 10 delayed- $\alpha$  particles; and in terms of residual nuclides, to 40  $(A, Z+1)$ , 30  $(A-1, Z+1)$ , 20  $(A-2, Z+1)$  and 10  $(A-4, Z-1)$ , respectively.

In general, the number of neutrons emitted per 100  $\beta^-$  decays,  $P_n$ , can be written as:

$$P_n = \sum_i i \times \beta_m^-;$$

and similar expressions can be written for  $\beta^-$ -delayed  $\alpha$  and proton emissions. The number of residual daughter nuclides  $(A, Z+1)$  populated via  $\beta^-$  decay is then:

$$\beta^- - \sum_i \beta_m^- - \sum_j \beta_{j\alpha}^- - \dots$$

Sometimes, the primary (parent)  $\beta$  decay can populate several excited states in the daughter nuclide, which can further decay by particle emission. However, in a case where the ground state of the daughter nuclide decays also by the same particle emission, some authors included its decay in the value for the corresponding  $\beta$ -delayed particle intensity. It is a policy of NUBASE2020 to not use such an approach for two main reasons: a) the energies of delayed particles emitted from excited states are generally much higher compared to those emitted from the ground state, thus implying different subsequent processes; b) the characteristic decay times from excited states are related to the parent, whereas decays from the daughter’s ground state are connected to the daughter nuclide itself. For example,  ${}^9\text{C}$  decays via  $\beta^+$  emission to the ground state of the proton-unbound  ${}^9\text{B}$  nuclide (feeding intensity of 54.1(1.5)% [2001Be51]) and to several excited states that are proton and/or  $\alpha$  unbound. If one takes the  $\beta^+$  intensities to the excited states in  ${}^9\text{B}$  from Ref. [2000Ge09] and renormalizes them to per 100 decays of the parent, then  $\beta^+ p = 7.5(0.6)\%$  and  $\beta^+ \alpha = 38.4(1.6)\%$  can be determined for  ${}^9\text{C}$ . In a slightly different example,  ${}^8\text{B}$  decays via  $\beta^+$  emission only to two excited,  $\alpha$ -unbound states in  ${}^8\text{Be}$ , but not to the  ${}^8\text{Be}$  ground state. Thus, one may write  $\beta^+ = 100\%$  and  $\beta^+ \alpha = 100\%$ , and therefore, no net population of the  ${}^8\text{Be}$  ground state.

It should be pointed out that the percentages given in the Table I are related to 100 decays of the parent nuclei, rather than to the primary decay mode fraction. For example, the delayed-fission probability in the decay of  ${}^{228}\text{Np}$  is given in the original article as 0.020(9)% [1994Kr13], but this value is relative to the  $\varepsilon$  process, which has an intensity of 60(7)%. Thus, the renormalized delayed-fission intensity is  $0.020(9)\% \times 0.60(7) = 0.012(6)\%$  of the total decay intensity.




Fig. 6. Nuclear chart displaying the main decay mode for nuclei in their ground state.

In addition to applying direct updates from the literature, partial evaluations completed by other authors were also considered in the evaluation of delayed particle data. For example, in compiling data for delayed proton- and  $\alpha$ -branching intensities, the work of Hardy and Hagberg [1989Ha.A], Jonson and Riisager [15], Blank and Borge [16] and Pfützner *et al.* [17], where the corresponding physics was reviewed, as well as the recent compilation of Batchelder [18], were consulted. Similarly, data on  $\beta$ -delayed neutron emission probabilities that were recommended by a recent IAEA Coordinated Research Project [19] were also consulted.

## 2.6 Isotopic abundances

Isotopic abundances are given in the decay field of Table I with the symbol  $IS$  and the values were taken from the most-recent publication of Meija *et al.* [20]. In several cases the  $IS$  values are listed in Ref. [20] as an interval  $[a, b]$ , but in Table I they are given as  $IS = (a + b)/2$  with a variance  $\sigma^2 = (b - a)^2/12$  (see section 3.2).

## 2.7 Year of discovery

Similarly to the previous version of NUBASE [3, 4], Table I includes information about the year of discovery for each nuclide in its ground or isomeric state. For the former, recommendations by Thoennessen [21] were adopted. Similar criteria were used when assigning the year of discovery for isomeric states.

## 2.8 References and Dissemination

The year of the ENSDF archival file that was consulted during the development of NUBASE2020 is given in Table I. The entry is left blank when information for a particular nuclide was not available in ENSDF.

The bibliographical information used in NUBASE2020 is referenced by means of the “Nuclear Science Reference” (NSR) database [22] keynumber style. However, references quoted in Table I are abbreviated with the first two digits of the year of publication being omitted from the NSR-style keynumbers. They are followed by up to three one-letter codes which specify the added or modified physics quantities (see the Explanation of Table I).

In cases where more than one reference was needed to describe a particular update, they were included as comments in Table I. No references were given for estimated values.

The initials of the former and present evaluators, e.g. GAU (G. AUDI), HWJ (W. HUANG), FGK (F. KONDEV), MMC (M. MACCORMICK), SAR (S. NAIMI) WGM (M. WANG), AHW (A. WAPSTRA), were used as reference keys where it may not be clear that the re-interpretation of data was made by the NUBASE evaluators.

In cases of directly measured spins, references are provided only to papers that were not included in the most-recent compilation of McDonald *et al.* [14].

The complete reference list is given at the end of this issue

(see AME2020, Part II), together with the references used in AME2020.

The recommended data for the basic nuclear physics properties are also made available as an ASCII-formatted file (**nubase.mas20**) at the dissemination websites of the collaboration [23].

## 3 Policies of NUBASE2020

### 3.1 Trends in neighboring nuclei (TNN)

In general, NUBASE2020 contains numerical and bibliographical information for all known nuclei for which at least one property is experimentally known. However, it also includes results on yet unobserved nuclides, as well as data on properties (mostly excitation energy for isomers, half-lives and spins and/or parities) that are not yet measured. Such values are estimated from the systematics trends of a particular property in neighboring nuclei by ensuring a continuity in  $N$ , in  $Z$ ,  $A$ , and in  $N - Z$ . This approach allowed to follow the behavior of a particular property as a function of  $N$  and  $Z$  in a consistent way and it proved beneficial in deducing values for other relevant properties. Similarly to AME2020, where masses estimated from Trends from the Mass Surface (TMS) are flagged with ‘#’, the same symbol is used in NUBASE2020 to indicate non-experimental information inferred from TNN. It should be pointed out, however, that deviations from TNN are expected when nuclear structure effects, such as deformation and/or shape changes, occur. Such data were taken into account to the best knowledge of the present authors.

### 3.2 Averaging procedure and uncertainties

It is a policy of NUBASE2020 to use one standard deviation as a representation of uncertainties associated with the recommended values. Unfortunately, authors of research articles do not always clarify the meaning of their reported uncertainties and, under such circumstances, these values are assumed to be one standard deviation. In several instances, uncertainties are not given at all and in such cases they were estimated by the evaluators, considering the limitations of the employed experimental method. When both the statistical and systematic uncertainties were reported in the literature, they were combined in a quadrature by the NUBASE2020 evaluators.

Sometimes lower ( $l$ ) and upper ( $u$ ) limits of a particular quantity,  $q$ , are reported in the literature, e.g.  $q \in [l, u]$ . A policy of NUBASE2020 is that uniform probability distribution is assumed in such cases, which yields a mean value of  $m = (l + u)/2$  and a standard deviation of  $\sigma = (u - l)/\sqrt{12} \approx 0.29 \times (u - l)$ .

When results from two or more independent measurements were reported in the literature, the corresponding values were weighted by their reported uncertainties and averaged. The weighted average value and its uncertainty are cal-

culated as:

$$\bar{x} \pm \Delta\bar{x} = \sum_{i=1}^N w_i x_i / \sum_{i=1}^N w_i \pm \sqrt{1 / \sum_{i=1}^N w_i} \quad (3)$$

where  $w_i = 1/\Delta x_i^2$  and  $x_i$  and  $\Delta x_i$  are the value and its uncertainty reported in the  $i^{\text{th}}$  experiment, and the summation is over all  $N$  experiments. For each average value the NORMALIZED CHI,  $\chi_n$  (or ‘consistency factor’ or ‘Birge ratio’), defined as:

$$\chi_n = \sqrt{\frac{1}{N-1} \sum_{i=1}^N w_i (x_i - \bar{x})^2} \quad (4)$$

is also calculated.

It is a policy of NUBASE2020 to use the weighted average result (equation 3) when  $\chi_n$  is smaller or equal to 2.5. In cases where  $\chi_n$  is larger than 2.5, but less or equal to 4, departure from the statistical result (equation 3) is allowed and the external uncertainty for the average value is adopted:

$$\bar{x} \pm \Delta\bar{x} = \sum_{i=1}^N w_i x_i / \sum_{i=1}^N w_i \pm \chi_n \times \sqrt{1 / \sum_{i=1}^N w_i} \quad (5)$$

In rare cases when  $\chi_n$  is larger than 4, all individual uncertainties are considered to be irrelevant and the arithmetic (unweighted) average is adopted:

$$\bar{x} \pm \Delta\bar{x} = \frac{1}{N} \sum_{i=1}^N x_i \pm \sqrt{\frac{1}{N(N-1)} \sum_{i=1}^N (x_i - \bar{x})^2} \quad (6)$$

The values used in the statistical analysis of a particular quantity are given as comments in Table I. When contradictory (discrepant) results were identified in the literature, a great deal of attention was focused on establishing the reason for such discrepancies, and consequently, on rejecting (or correcting) the corresponding unreliable data prior to performing the statistical analysis. The reasons for such decisions are given as comments in Table I.

### 3.3 Symmetrization of asymmetric uncertainties

Experimental results are sometimes reported in the literature with asymmetric uncertainties, e.g.  $X_{-b}^{+a}$ , and it is a policy of NUBASE to symmetrize these uncertainties.

Similarly to the previous version of NUBASE [1–4], the asymmetric uncertainty is associated with a two-piece normal distribution (sometimes called “split-normal distribution” or “Fechner distribution”),  $TN(X, a, b)$ , and the symmetrization is achieved by mapping this distribution into a normal (symmetric) distribution,  $N(\mu, \sigma)$ , where  $\mu$  is the mean value and  $\sigma$  is the standard deviation.

The probability density function of a two-piece normal distribution is given as:

$$f(x) = \begin{cases} A \times \exp[-(x-X)^2/2a^2] & \text{if } x > X, \\ A \times \exp[-(x-X)^2/2b^2] & \text{if } x < X \end{cases} \quad (7)$$

It has a modal (most probable) value of  $x = X$ , a standard deviation  $b$  for  $x < X$  and a standard deviation  $a$  for  $x > X$  (see Figure 7), with  $A = (\sqrt{\pi/2} \times (a+b))^{-1}$ . This distribution is formed by taking the left half of a normal distribution  $N(X, b)$  and the right half of a normal distribution  $N(X, a)$  and scaling them to give a common value of  $f(x)=A$  at the mode  $X$  (see Figure 7). The mean value and the variance of this distribution can be determined as [24]:

$$\mu = X + \sqrt{2/\pi} \times (a-b) \quad (8)$$

$$\sigma^2 = (1-2/\pi) \times (a-b)^2 + a \times b \quad (9)$$

The median value  $m$ , which divides the distribution into two equal areas is then:

$$m = \begin{cases} X + a\sqrt{2} \times \text{erf}^{-1}\left(\frac{a-b}{2a}\right) & \text{if } a > b, \\ X + b\sqrt{2} \times \text{erf}^{-1}\left(\frac{a-b}{2b}\right) & \text{if } b > a. \end{cases} \quad (10)$$

If one takes  $\text{erf}^{-1}(z) \simeq \sqrt{\pi}z/2$  then

$$m - X \simeq \sqrt{\pi/8} \times (a-b) \simeq 0.6267 \times (a-b) \quad (11)$$

In order to allow for a small non-linearity that appears for high values of  $m - X$ , equation 11 is modified to:

$$m \simeq X + 0.64 \times (a-b) \quad (12)$$

Following the above approach, the two-piece normal distribution  $TN(X, a, b)$  is mapped into an equivalent normal (symmetric) distribution  $N(m, \sigma)$  (see Figure 7) that have a mean value equal to the median value  $m$  (equation 12) and variance  $\sigma$  (equation 9). As a consequence,  $X_{-b}^{+a}$  is symmetrized to  $m \pm \sigma$  and the latter is adopted in NUBASE2020.

### 3.4 Rounding policy

In general, values for properties presented in NUBASE2020 and their uncertainties are rounded off, even if unrounded ones were given in the literature or in ENSDF. However, for some very precise data, as well as for data that were deemed essential for traceability purposes (e.g. isotopic abundances), the precisions quoted in the original publications were retained.

In cases where the two furthest-left significant digits in the uncertainty were larger than a given limit (set to 30 for the mass excess and excitation energy of isomers in order to be consistent with AME, and set to 25 for half-lives and branching ratios, as generally used in ENSDF), the adopted values and corresponding uncertainties were rounded off accordingly.

## 4 Conclusions and outlook

The NUBASE2020 evaluated nuclear data library contains the recommended values for the basic nuclear physics properties for all known nuclei, such as mass excess, excitation




Fig. 7. (dashed black) original Normal distributions,  $N(X, b)$  (left on  $X$ ) and  $N(X, a)$  (right on  $X$ ), associated with the measured quantity  $X_{-b}^{+a}$ ; (solid blue) a two-piece normal distribution  $TN(X, a, b)$ ; (solid red) the equivalent Normal (symmetric) distribution,  $N(m, \sigma)$ ; see section 3.3 for details.

energy of the excited isomeric state, half-life, spin and parity, decay modes and their intensities, isotopic abundance (for stable nuclides), year of discovery, as well as the corresponding bibliographical information. It also contains information for yet unobserved nuclei whose properties were estimated by following the systematic trends in neighboring nuclei.

One of the main requirements in the development of NUBASE2020 was to cover the available experimental data as completely as possible and to provide proper references to all experimental results, especially for cases that are not included in ENSDF or in other topical evaluations. Such a traceability would allow any user to promptly review the recommended data and, if necessary, to undertake a re-evaluation.

NUBASE2020 is an integral part of AME2020 and the synchronization of these two libraries allows better homogeneity of all experimental data to be achieved. Furthermore, assignments of isomeric states and determination of their excitation energies were put on a firm basis and the data were improved.

In the future development of NUBASE, it is envisioned to include additional nuclear properties, such as magnetic and quadrupole moments, charge-radii and isotope shifts, cross sections of importance to nuclear astrophysics applications, as well as additional decay properties of relevance to energy and non-energy applications, in order to better serve the broader nuclear physics community.

## 5 Acknowledgments

This work was supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Contract No. DE-AC02-06CH11357 (ANL) and in part by the National Key Research and Development Program of China (Grant No. 2016YFA0400504) and the Strategic Priority Research Program of Chinese Academy of Sciences (CAS, Grant No. XDB34000000) (IMP). W.J. Huang acknowledges the financial support by the Max-Planck-Society. S. Naimi acknowledges the support of the RIKEN Pioneering Project Funding.

We thank many colleagues who responded to our queries about the experimental data, as well as those who sent us preprints of their work prior publication. We also thank D. Lunney and B. Kay for the careful reading of the manuscript and useful comments.

## References

- 1 G. Audi, O. Bersillon, J. Blachot and A.H. Wapstra, Nucl. Phys. A **624** (1997) 1.
- 2 G. Audi, O. Bersillon, J. Blachot and A.H. Wapstra, Nucl. Phys. A **729** (2003) 3.
- 3 G. Audi, F.G. Kondev, M. Wang, B. Pfeiffer, X. Sun, J. Blachot and M. MacCormick, Chin. Phys. C **36** (2012) 12.
- 4 G. Audi, F.G. Kondev, M. Wang, W.J. Huang, S. Naimi,

- Chin. Phys. **C 41** (2017) 03001.
- 5 T.W. Burrows, Nucl. Instrum. Meth. **286** (1990) 595; <http://www.nndc.bnl.gov/ensdf/>
  - 6 F.G. Kondev, G.D. Dracoulis, T. Kibédi, At. Data Nucl. Data Tables **103-104** (2015) 50; F.G. Kondev, G.D. Dracoulis, T. Kibédi, At. Data Nucl. Data Tables **105-106** (2015) 105 (erratum)
  - 7 G.D. Dracoulis, P.M. Walker, F.G. Kondev, Rep. Prog. Phys. **79** (2016) 076301.
  - 8 M. MacCormick and G. Audi, Nucl. Phys. **A 925** (2014) 61.
  - 9 Y.H. Lama, B. Blank, N.A. Smirnova, J.B. Bueb, and M.S. Antony, At. Data Nucl. Data Tables **99** (2013) 680.
  - 10 M.P. Unterweger and R. Fitzgerald, Appl. Radiat. Isot. **159** (2020) 108976.
  - 11 M.P. Unterweger and R. Fitzgerald, Appl. Radiat. Isot. **87** (2014) 92.
  - 12 General Policies, Nucl. Data Sheets **164** (2020) 478.
  - 13 G.H. Fuller, J. Phys. Chem. Ref. Data **5** (1976) 835.
  - 14 A. MacDonald, B. Karamy, K. Setoodehnia, B. Singh, Nucl. Data Sheets **114** (2013) 397.
  - 15 B. Jonson and K. Riisager, Nucl. Phys. **A 693** (2001) 77.
  - 16 B. Blank and M. Borge, Prog. Part. Nucl. Phys. **60** (2008) 403.
  - 17 M. Pfützner, M. Karny, L.V. Grigorenko, K. Riisager, Rev. Mod. Phys. **84** (2012) 567.
  - 18 J.C. Batchelder, At. Data Nucl. Data Tables **132** (2020) 101323.
  - 19 IAEA Coordinated Research Project on a reference database for beta-delayed neutron emission; <https://www-nds.iaea.org/beta-delayed-neutron/>;
  - 20 M. Birch, B. Singh, I. Dillmann, D. Abriola, T.D. Johnson, E.A. McCutchan, and A.A. Sonzogni, Nucl. Data Sheets **128** (2015) 131; J. Liang, B. Singh, E.A. McCutchan, I. Dillmann, M. Birch, A.A. Sonzogni, X. Huang, M. Kang, J. Wang, G. Mukherjee, K. Banerjee, D. Abriola, A. Algora, A.A. Chen, T.D. Johnson, and K. Miernik, Nucl. Data Sheets **168** (2020) 1.
  - 21 J. Meija, T.B. Coplen, M. Berglund, W.A. Brand, P. De Bièvre, M. Gröning, N.E. Holden, J. Irrgeher, R.D. Loss, T. Walczyk and T. Prohaska, Pure Appl. Chem. **88(3)** (2016) 293.
  - 22 M. Thoennessen, “The Discovery of Isotopes: A Complete Compilation”, Springer International Publishing Switzerland 2016; <https://people.nsl.msui.edu/thoennes/isotopes/>
  - 23 B. Pritychenko, E. Běták, M.A. Kellett, B. Singh and J. Totans, Nucl. Instrum. Methods Phys. Res. **A640** (2011) 213; <http://www.nndc.bnl.gov/nsr/>
  - 24 The NUBASE2020 files in electronic-distribution format and additional complementary information can be retrieved from the main dissemination websites of the present collaboration: IMP at <http://amdc.impcas.ac.cn/>; ANL at <https://www.anl.gov/phy/atomic-mass-data-resources>; IAEA at <https://www-nds.iaea.org/amdc/>
  - 25 S. John, Commun. Statist. -Theor. Meth. **11(8)** (1982) 879.

**Table I. The NUBASE2020 table of nuclear and decay properties**

## EXPLANATION OF TABLE I

Data are presented in groups ordered by increasing mass number,  $A$ .

Nuclide	Nuclide name: mass number $A = N + Z$ and element symbol. The superscript suffixes ‘ $m$ ’, ‘ $n$ ’, ‘ $p$ ’, ‘ $q$ ’, ‘ $r$ ’ and ‘ $x$ ’ indicate assignments to excited isomeric states with a half-life greater than 100 ns. Suffixes ‘ $p$ ’ and ‘ $q$ ’ can also indicate non-isomeric levels, which are used in AME2020. Suffix ‘ $r$ ’ can also indicate a state from a proton resonance occurring in $(p,\gamma)$ reactions (e.g. $^{28}\text{Si}^r$ ). Suffix ‘ $x$ ’ can also indicate a mixture of levels with a relative ratio, $R$ , given in the ‘Half-life’ column. They occur in spallation reactions or fission and are labeled as ‘spmix’ or ‘fsmix’ in the ‘ $J^\pi$ ’ column, respectively. Suffixes ‘ $i$ ’ and ‘ $j$ ’ indicate Isobaric Analog States.																		
Mass excess	<p>Mass excess <math>[M(\text{in u}) - A]</math> and its uncertainty (one standard deviation) in keV, as recommended in AME2020.</p> <p>Rounding policy: in cases where the furthest-left significant digit in the uncertainty is larger than 3, values are rounded-off, but not to more than tens of keV. (Examples: <math>2345.67 \pm 2.78 \rightarrow 2345.7 \pm 2.8</math>, <math>2345.67 \pm 4.68 \rightarrow 2346 \pm 5</math>, but <math>2346.7 \pm 468.2 \rightarrow 2350 \pm 470</math>).</p> <p># indicates that the Mass excess value and its uncertainty are not derived from experimental data, but at least partly from the Trends from the Mass Surface (see the AME2020 publication for details).</p>																		
Excitation energy	<p>The energy difference between the excited isomer and the ground state, and its uncertainty (one standard deviation) in keV. The rounding policy is the same as for the mass excess (see above), with the exception of the very precise values for the <math>^{229}\text{Th}^m</math> and <math>^{235}\text{U}^m</math> isomers, which are given in the comments.</p> <p># indicates that the excitation energy and its uncertainty are not derived from experimental data, but from the Trends in Neighboring Nuclei (TNN) (see section 3.1)</p> <p>When the excitation energy is determined by an <i>external</i> relation, it is followed by one or two-letters code (the origin code) that indicates the method used to establish such a relation. The field is left blank when the value is derived from <math>\gamma</math>-ray spectroscopy data (<i>internal</i> relation):</p> <table border="0" style="margin-left: 20px;"> <tr><td>MD</td><td>mass doublet</td></tr> <tr><td>RQ</td><td>reaction <math>Q</math>-value</td></tr> <tr><td>AD</td><td><math>\alpha</math> energy difference</td></tr> <tr><td>BD</td><td><math>\beta</math>-decay end-point energy data</td></tr> <tr><td>p, 2p</td><td>one-, two-proton decay</td></tr> <tr><td>IT</td><td>combination of AME and <math>\gamma</math>-ray spectroscopy data</td></tr> <tr><td>Nm</td><td>estimated value derived using the Nilsson model</td></tr> </table> <p>When the existence of a nuclide or an isomer is questionable the following codes are used:</p> <table border="0" style="margin-left: 20px;"> <tr> <td>EU</td> <td>the existence is under discussion (e.g. <math>^{185}\text{Bi}^n</math>). If the existence is strongly doubted, no excitation energy and mass excess values are given, and they are replaced by the keyword “non-exist” (e.g. <math>^{138}\text{Pm}^m</math>).</td> </tr> <tr> <td>RN</td> <td>the isomer has been proven not to exist (e.g. <math>^{181}\text{Pb}^m</math>). Excitation energy and mass excess values are replaced by the keyword “non-exist”.</td> </tr> </table> <p><i>Remark:</i> codes EU and RN are also used when the discovery of a nuclide is questioned (e.g. <math>^{260}\text{Fm}</math> and <math>^{289}\text{Lv}</math>). In this case, a mass excess value derived from the Trends from the Mass Surface (see the AME2020 publication for details) is always given for the ground state.</p>	MD	mass doublet	RQ	reaction $Q$ -value	AD	$\alpha$ energy difference	BD	$\beta$ -decay end-point energy data	p, 2p	one-, two-proton decay	IT	combination of AME and $\gamma$ -ray spectroscopy data	Nm	estimated value derived using the Nilsson model	EU	the existence is under discussion (e.g. $^{185}\text{Bi}^n$ ). If the existence is strongly doubted, no excitation energy and mass excess values are given, and they are replaced by the keyword “non-exist” (e.g. $^{138}\text{Pm}^m$ ).	RN	the isomer has been proven not to exist (e.g. $^{181}\text{Pb}^m$ ). Excitation energy and mass excess values are replaced by the keyword “non-exist”.
MD	mass doublet																		
RQ	reaction $Q$ -value																		
AD	$\alpha$ energy difference																		
BD	$\beta$ -decay end-point energy data																		
p, 2p	one-, two-proton decay																		
IT	combination of AME and $\gamma$ -ray spectroscopy data																		
Nm	estimated value derived using the Nilsson model																		
EU	the existence is under discussion (e.g. $^{185}\text{Bi}^n$ ). If the existence is strongly doubted, no excitation energy and mass excess values are given, and they are replaced by the keyword “non-exist” (e.g. $^{138}\text{Pm}^m$ ).																		
RN	the isomer has been proven not to exist (e.g. $^{181}\text{Pb}^m$ ). Excitation energy and mass excess values are replaced by the keyword “non-exist”.																		



## Isomer assignment:

- \* when the available experimental information is insufficient to unambiguously determine which state is the ground state and which one is the excited isomer, as well as in cases where the uncertainty ( $\Delta E_x$ ) of the excitation energy ( $E_x$ ) is greater than half the excitation energy value ( $\Delta E_x > E_x/2$ ), these quantities are followed by an asterisk (see for example  $^{102}\text{Y}$  and  $^{102}\text{Y}^m$ ).
- & when the ordering of the ground state and the excited isomer is reversed in comparison to the assignment made in ENSDF, the ampersand sign is added in the table (see for example  $^{100}\text{Y}$  and  $^{100}\text{Y}^m$ ).

## Half-life

Half-life value (see section 2.3).

s = seconds; m = minutes; h = hours; d = days; y = years; 1 y = 365.2422 d = 31 556 926 s.

STABLE = stable nuclide or nuclide for which no finite half-life value was established.

# indicate non-experimental value estimated from Trends in Neighboring Nuclei (TNN) (see section 3.1).

subunits:

ms	:	$10^{-3}$	s	millisecond	ky	:	$10^3$	y	kiloyear
$\mu\text{s}$	:	$10^{-6}$	s	microsecond	My	:	$10^6$	y	megayear
ns	:	$10^{-9}$	s	nanosecond	Gy	:	$10^9$	y	gigayear
ps	:	$10^{-12}$	s	picosecond	Ty	:	$10^{12}$	y	terayear
fs	:	$10^{-15}$	s	femtosecond	Py	:	$10^{15}$	y	petayear
as	:	$10^{-18}$	s	attosecond	Ey	:	$10^{18}$	y	exayear
zs	:	$10^{-21}$	s	zeptosecond	Zy	:	$10^{21}$	y	zettayear
ys	:	$10^{-24}$	s	yoctosecond	Yy	:	$10^{24}$	y	yottayear

 $J^\pi$ 

Spin and parity (see section 2.4):

- () uncertain spin and/or parity based on *weak* experimental arguments.
- \* directly measured spin (see section 2.4).
- # non-experimental value estimated from Trends in Neighboring Nuclei (TNN) (see section 3.1) or from theoretical predictions.
- high high spin.
- low low spin.
- am same  $J^\pi$  as the  $\alpha$ -decay parent nuclide.
- $T$  isospin multiplet value for Isobaric Analog States (see section 2.2.1).

## Ens

Year of the ENSDF file archive. In order to reduce the width of the table, the two century digits are omitted.

## Reference

Reference key-numbers (see section 2.8). In order to reduce the width of the table, the two century digits are omitted from the NSR reference key. The complete references list and associated NSR reference key-numbers are given in the second AME publication in the present volume.

10Cr02	updates derived from a <i>primary</i> (journal article) reference with the keynumber taken from the “Nuclear Science Reference” (NSR) database [22] (see section 2.8). When the keynumber was not available, the style 12Ma.1 was provisionally adopted.
12Dr.A	updates derived from a <i>secondary</i> (abstract, preprint, private communication, not-refereed conference proceeding, thesis or laboratory report) reference.
AHW	(or GAU, HWJ, FGK, MMC, SAR, WGM), re-interpretation by one of the NUBASE evaluators.
Mirror	deduced from mirror nuclide properties.
Imme	deduced from Isobaric Multiplet Mass Equation.

The reference key-numbers are followed by codes having up to three letters that indicates which physics quantity was added or modified:

M	mass excess
E	isomer excitation energy
T	half-life
J	spin and/or parity
D	decay mode and/or its intensity
I	identification

Year of  
discovery

Year of discovery assigned for the ground and excited isomeric states (see section 2.7).

Decay modes  
and intensities

Decay modes followed by their intensities and associated uncertainties, both in % (see section 2.5). The ordering is according to decreasing intensities. The uncertainties are given by the ENSDF-style format, e.g.  $\alpha=25.9\ 23$  stands for  $\alpha=25.9\% \pm 2.3\%$ . The notation  $1.8e-12$  stands for  $1.8 \times 10^{-12}$ .

$\alpha ?$  means that the  $\alpha$ -decay mode is energetically allowed, but not experimentally observed

$\alpha=?$  means that the  $\alpha$ -decay is observed, but its intensity is not experimentally known

$\alpha$	$\alpha$ emission
p 2p	proton emission      2-proton emission
n 2n	neutron emission      2-neutron emission
$\epsilon$	electron capture
$e^+$	positron emission
$\beta^+$	$\beta^+$ decay      ( $\beta^+ = \epsilon + e^+$ )
$\beta^-$	$\beta^-$ decay
$2\beta^-$	double $\beta^-$ decay
$2\beta^+$	double $\beta^+$ decay
$\beta^-n$	$\beta^-$ -delayed neutron emission
$\beta^-2n$	$\beta^-$ -delayed 2-neutron emission
$\beta^-3n$	$\beta^-$ -delayed 3-neutron emission
$\beta^+p$	$\beta^+$ -delayed proton emission
$\beta^+2p$	$\beta^+$ -delayed 2-proton emission
$\beta^+3p$	$\beta^+$ -delayed 3-proton emission
$\beta^-\alpha$	$\beta^-$ -delayed $\alpha$ emission
$\beta^+\alpha$	$\beta^+$ -delayed $\alpha$ emission
$\beta^-d$	$\beta^-$ -delayed deuteron emission
$\beta^-t$	$\beta^-$ -delayed triton emission
IT	internal transition

SF	spontaneous fission
$\beta^+$ SF	$\beta^+$ -delayed fission
$\beta^-$ SF	$\beta^-$ -delayed fission
$^{24}\text{Ne}$	heavy cluster emission

For stable or long-lived nuclides:

IS	Isotopic abundance taken from Ref. [20] (see section 2.6).
----	--

- \* Indicates a comment to a nuclide, which is given below the block of data corresponding to the same A. The asterisk symbol is also included at the end of the data line. The comment starts with a letter code, similar to the one that follows the reference key-number (see above), indicating to which physics quantity the remark is applied. It contains: (i) information explaining how a specific value was derived; (ii) reasons for changing a value or its uncertainty that were reported by the original authors, or for rejecting it; (iii) complementary references to updated data; (iv) individual values used in the statistical analysis of data.

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^1_0\text{n}$	8071.3181	0.0004			609.8 s 0.6	$1/2^+*$	06 19Ma40 T	1932	$\beta^-$ =100	*
$^1_1\text{H}$	7288.9710	0.0001			STABLE	$1/2^+*$	06 11Be53 D	1920	IS=99.9855 78	*
* $^1_0\text{n}$	T : from the world average in 19Ma40=609.8(0.6) ( $\tau$ =879.7(0.8) s)									**
* $^1_1\text{H}$	M : rounded from 7288.971064(0.000013) keV									**
$^2_0\text{H}$	13135.7229	0.0001			STABLE	$1^+*$	03	1932	IS=0.0145 78	*
* $^2_0\text{H}$	M : rounded from 13135.722895(0.000015) keV									**
$^3_0\text{H}$	14949.8109	0.0001			12.32 y 0.02	$1/2^+*$	00	1934	$\beta^-$ =100	*
$^3_1\text{He}$	14931.2188	0.0001			STABLE	$1/2^+*$	98	1934	IS=0.0002 2	*
$^3_2\text{Li}$	28670#	2000#			p-unstable	$3/2^- \#$	98		p ?	*
* $^3_0\text{H}$	M : rounded from 14949.81090(0.00008) keV									**
* $^3_1\text{He}$	M : rounded from 14931.21888(0.00006) keV									**
* $^3_2\text{Li}$	I : identification in 69Wi13 not accepted									**
$^4_0\text{H}$	24620	100			139 ys 10	$2^-$	98 03Me11 T	1981	n=100	*
$^4_1\text{He}$	2424.9158	0.0001			STABLE	$0^+$	98	1908	IS=99.9998 2	*
$^4_2\text{Li}$	25320	210			91 ys 9	$2^-$	98 65Ce02 T	1965	p=100	*
* $^4_0\text{H}$	T : width=3.28(0.23) MeV; other 91Go19=4.7(1.0) outweighed, not used									**
* $^4_1\text{He}$	M : rounded from 2424.91587(0.00015) keV									**
$^5_0\text{H}$	32890	90			86 ys 6	$(1/2^+)$	19 17Wu03 T	1987	2n=100	*
$^5_1\text{He}$	11231	20			602 ys 22	$3/2^-$	02	1937	n=100	*
$^5_2\text{Li}$	11680	50			370 ys 30	$3/2^-$	02	1941	p=100	*
$^5_3\text{Be}$	37140#	2000#			p-unstable	$1/2^+ \#$	18		p ?	*
* $^5_0\text{H}$	T : from width=5.3(0.4) MeV in 17Wu03									**
* $^5_1\text{He}$	J : from angular distribution data consistent with $l = 0$ in 01Ko52									**
* $^5_2\text{Li}$	T : from width=758(28) keV, average 12Lu01=767(10) keV									**
* $^5_3\text{Be}$	T : 09Ak03=670 (12, stat) (30, syst) keV; Birge ratio=2.9									**
$^6_0\text{H}$	41880	250			294 ys 67	$2^- \#$	19	1984	n ?; 3n ?	*
$^6_1\text{He}$	17592.10	0.05			806.92 ms 0.24	$0^+$	02 15Pf01 D	1936	$\beta^-$ =100; $\beta^-$ -d=0.000278 18	*
$^6_2\text{Li}$	14086.8804	0.0014			STABLE	$1^+*$	02	1921	IS=4.85 171	*
$^6_3\text{Li}^i$	17649.76	0.10	3562.88	0.10	56 as 14	$0^+ T=1$	02 81Ro02 E	1981	IT=100	*
$^6_4\text{Be}$	18375	5			5.0 zs 0.3	$0^+$	02	1958	2p=100	*
$^6_5\text{B}$	47320#	2000#			p-unstable	$2^- \#$			2p ?	*
* $^6_0\text{H}$	T : from width=1.55(0.35) MeV, average 84Al08=1.8(5) MeV 86Be35=1.3(5) MeV									**
* $^6_1\text{He}$	D : other $\beta^-$ -d 09Ra33=1.65(0.10)e-6, but with 525-keV threshold									**
* $^6_2\text{Li}$	T : symmetrized from 12Kn01=806.89(0.11, stat)(+0.23-0.19, syst)									**
$^7_0\text{H}$	49140#	1000#			652 ys 558	$1/2^+ \#$	17 08Ca22 T	2003	2n ?	*
$^7_1\text{He}$	26073	8			2.51 zs 0.07	$(3/2)^-$	03 12Ca05 T	1967	n=100	*
$^7_2\text{Li}$	14907.105	0.004			STABLE	$3/2^-*$	03	1921	IS=95.15 171	*
$^7_3\text{Li}^i$	26150	30	11250	30	RQ	$3/2^- T=3/2$	03			*
$^7_4\text{Be}$	15769.00	0.07			53.22 d 0.06	$3/2^-$	03	1938	$\epsilon$ =100	*
$^7_5\text{Be}^i$	26750	30	10980	30	RQ	$3/2^- T=3/2$	03		p ?; $^3\text{He}$ ?; $\alpha$ ?	*
$^7_6\text{B}$	27677	25			570 ys 14	$(3/2)^-$	14 11Ch32 T	1967	p=100	*
* $^7_0\text{H}$	T : symmetrized from 08Ca22=0.09(+94-6) MeV									**
* $^7_1\text{He}$	T : from width=182(5) keV in 12Ca05; others 09Ak03=190(30)									**
* $^7_2\text{Li}$	T : 08De29=125(+40-15) 02Me07=150(80) 69St02=160(30) (outweighed)									**
* $^7_3\text{Li}$	T : from width=801(20) keV in 11Ch32									**
$^8_0\text{H}$	31609.68	0.09			119.5 ms 1.5	$0^+$	05 15Bi05 TD	1965	$\beta^-$ =100; $\beta^-$ -n=16 1;	*
$^8_1\text{Li}$	20945.80	0.05			838.7 ms 0.3	$2^+$	05 10Fl01 T	1935	$\beta^-$ -t=0.9 1	*
$^8_2\text{Li}^i$	31768	5	10822	5	RQ	$0^+ T=2$	05		$\beta^-$ =100; $\beta^-$ - $\alpha$ =100	*

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
$^8\text{Be}$	4941.67	0.04				81.9 as 3.7	$0^+$	05		1932	$\alpha=100$	
$^8\text{Be}^i$	21568	3	16626	3			$2^+$ frg. T=1	04Ti06	E	2004	$\alpha\approx 100$	
$^8\text{Be}^j$	32436.0	2.0	27494.3	2.0	RQ		$0^+$ T=2	05			n=39.4;d=27.0; $^3\text{H}=11.7$ ; $\alpha=7.9$ ;p=6.9; $^3\text{He}=6.6$ ; IT=0.60	
$^8\text{B}$	22921.6	1.0				771.9 ms 0.9	$2^+$	05 20Vi03	T	1950	$\beta^+=100$ ; $\beta^+\alpha=100$	
$^8\text{B}^i$	33546	8	10624	8	RQ		$0^+$ T=2	05		1975		
$^8\text{C}$	35064	18				3.5 zs 1.4	$0^+$	18		1974	2p=100	
$^8\text{He}$	D : $\% \beta^-$ t intensity from $^8\text{Be}$ 041											
$^8\text{Li}$	D : $\beta^-$ decay to first $2^+$ state in $^8\text{Be}$ , which decays 100% by $2\alpha$											
$^8\text{Li}$	T : average 10Fl01=838.40(0.36) 03Wi17=839.60(1.06) 90Sa16=840.3(0.9)											
$^8\text{B}$	T : average 20Vi03=771.9(1.7),773.9(1.8),770.9(1.7), uncertainty in the											
$^8\text{B}$	T : last value from priv. comm. with the authors (from 2.7 to 1.7),											
$^8\text{B}$	T : $^7\text{McZ}W=772(4)$ $^7\text{Wi}05=762(5)$ $^6\text{Ma}35=774(4)$											
$^8\text{C}$	T : from width=130(50) keV in $^{11}\text{C}32$											
$^9\text{He}$	40940	50				2.5 zs 2.3	$1/2^+(^+)$	06 16Ub01	J	1987	n=100	
$^9\text{Li}$	24954.91	0.19				178.2 ms 0.4	$3/2^-$	06 15Bi05	TD	1951	$\beta^-=100$ ; $\beta^-n=50.5$ 10	
$^9\text{Be}$	11348.45	0.08				STABLE	$3/2^-*$	06		1921	IS=100	
$^9\text{Be}^i$	25738.8	1.7	14390.3	1.7	RQ	1.25 as 0.10	$3/2^-$ T=3/2	06		1976		
$^9\text{B}$	12416.5	0.9				800 zs 300	$3/2^-$	06		1940	p=100	
$^9\text{B}^i$	27071.0	2.3	14654.5	2.5	RQ		$3/2^-$ T=3/2	06				
$^9\text{C}$	28911.0	2.1				126.5 ms 0.9	$3/2^-$	06 01Be51	D	1964	$\beta^+=100$ ; $\beta^+p=7.5$ 6; $\beta^+\alpha=38.4$ 16	
$^9\text{He}$	T : from width=180(100) keV in $^{13}\text{Al}14$ ; other width=100(60) keV in $^{99}\text{Bo}26$											
$^9\text{C}$	D : $\% \beta^+p$ from $\% \beta^+$ (to $^9\text{B}$ gs)=54.1(1.5) from 01Be51 and $\% \beta^+$ (to $^9\text{B}$ exc)											
$^9\text{C}$	D : from 00Ge09, but renormalized in order to have $\% \beta^+(^9\text{C})=100$ , and											
$^9\text{C}$	D : $\% p/\% \alpha$ from 00Ge09; $\% \beta^+\alpha=100 - \% \beta^+$ (to $^9\text{B}$ gs) - $\% \beta^+$											
$^9\text{C}$	J : from 04Ti06											
$^{10}\text{He}$	49200	90				260 ys 40	$0^+$	07		1994	2n=100	
$^{10}\text{Li}$	33053	13				2.0 zs 0.5	$(1^-, 2^-)$	07 94Yo01	TJ	1975	n=100	
$^{10}\text{Li}^m$	33250	40	200	40	RQ	3.7 zs 1.5	$1^+$	07 97Zi04	T	1994	IT=100	
$^{10}\text{Li}^n$	33530	40	480	40	RQ	1.35 zs 0.24	$(2^+)$	07 94Yo01	T	1993	IT=100	
$^{10}\text{Be}$	12607.49	0.08				1.387 My 0.012	$0^+$	07 10Ch18	T	1935	$\beta^-=100$	
$^{10}\text{Be}^i$	33787	21	21179	21	RQ		$(2^-)$ T=2	07			n ?; p ?; $^3\text{H}$ ?	
$^{10}\text{B}$	12050.611	0.015				STABLE	$3^+*$	07		1920	IS=19.65 44	
$^{10}\text{B}^i$	13790.66	0.04	1740.05	0.04			$0^+$ T=1	07			IT=100	
$^{10}\text{C}$	15698.67	0.07				19.3011 s 0.0015	$0^+$	07 16Du10	T	1949	$\beta^+=100$	
$^{10}\text{N}$	38800	400				143 ys 36	$1^-, 2^-$	17 17Ho10	TJ	2002	p ?	
$^{10}\text{He}$	D : most probably 2 neutron emitter from S2n=-1440(90) keV											
$^{10}\text{He}$	T : from width=1.76(0.27) MeV, average 10Ko43=1.8(4) and 10Jo06=1.73(0.36),											
$^{10}\text{He}$	T : the latter average of 1.11(0.76), assuming a single narrow resonance,											
$^{10}\text{He}$	T : and 1.91(0.41), assuming two overlapping resonances; others:											
$^{10}\text{He}$	T : width=2 MeV in $^{12}\text{Si}07$ , 100-500 keV in $^{94}\text{Os}04$ , <1.2 MeV in $^{94}\text{Ko}16$											
$^{10}\text{Li}^m$	T : from average width=120(+100-50) keV in $^{97}\text{Zi}04$ and 100(70) keV in $^{94}\text{Yo}01$											
$^{10}\text{Li}^n$	T : from average width=358(23) keV in $^{94}\text{Yo}01$ , 150(70) keV in $^{93}\text{Bo}03$ ;											
$^{10}\text{Li}^n$	T : Birge ratio=2.8											
$^{10}\text{Be}$	T : average 10Ch18=1.386(0.016) 10Ko19=1.388(0.018)											
$^{10}\text{C}$	T : average 16Du10=19.2969(0.0074), 19.3009(0.0017) 09Ba06=19.282(0.011)											
$^{10}\text{C}$	T : 08Ia01=19.310(0.004) 90Ba02=19.295(0.015) 74Az01=19.280(0.020)											
$^{10}\text{C}$	T : $^7\text{Ro}21=19.150(0.030)$											
$^{10}\text{N}$	T : from width=3.1(+0.9-0.7) MeV for J=2- in $^{17}\text{Ho}10$ ; other:											
$^{10}\text{N}$	T : width=2.5(+2.0-1.5) MeV for J=1- in $^{17}\text{Ho}10$ .											
$^{11}\text{Li}$	40728.3	0.6				8.75 ms 0.06	$3/2^-*$	12 12Ke01	D	1966	$\beta^-=100$ ; $\beta^-n=86.3$ 9; $\beta^-2n=4.1$ 4; $\beta^-3n=1.9$ 2; $\beta^-\alpha=1.7$ 3; $\beta^-d=0.0130$ 13; $\beta^-t=0.0093$ 8	
$^{11}\text{Be}$	20177.17	0.24				13.76 s 0.07	$1/2^+*$	12 19Re03	D	1958	$\beta^-=100$ ; $\beta^-\alpha=3.3$ 1;	

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{11}\text{Be}^i$	41336	20	21158	20	RQ	0.93 zs 0.13	$3/2^-$ T=5/2	MMC162J	1997	$\beta^-$ p=0.0013 3; $\beta^-$ n ?
$^{11}\text{B}$	8667.708	0.012				STABLE	$3/2^-$ *	12	1920	IT=80.35 44
$^{11}\text{B}^i$	21228	9	12560	9	RQ	T=3/2	$1/2^+, (3/2^+)$	12	1963	
$^{11}\text{B}^j$	42230	80	33570	80	2p		$3/2^-$ T=5/2	MMC146J		
$^{11}\text{C}$	10649.40	0.06				20.3402 m 0.0053	$3/2^-$ *	12 18Va04	T 1934	$\beta^+$ =100
$^{11}\text{C}^i$	22810	40	12160	40	RQ		$1/2^+$ T=3/2	12 71Wa21	D 1971	p=?
$^{11}\text{N}$	24366	5				585 ys 7	$1/2^+$	12 19We11	T 1974	p=100
$^{11}\text{N}^m$	25110	60	740	60		690 ys 80	$1/2^-$	12 96Ax01	ETJ 1974	p=100
$^{11}\text{O}$	47740	60				198 ys 12	$(3/2^-)$	20 20We08	TJ 2019	2p=100
* $^{11}\text{Li}$	T : average 97Mo35=8.99(0.10) 96Mu19=8.2(0.2) 95Re.A=8.4(0.2)									
* $^{11}\text{Li}$	T : 81Bj01=8.83(0.12) and 74Ro31=8.5(0.2)									
* $^{11}\text{Be}$	D : % $\beta^-$ - $\alpha$ from 19Re03=3.30(0.10); other 81A03=2.9(4)									
* $^{11}\text{Be}$	D : % $\beta^-$ -p from 19Ay03=0.0013(0.0003); others (indirect) 14Ri01=0.00083(9)									
* $^{11}\text{Be}$	D : 20Ri02<0.00022%									
* $^{11}\text{Be}$	J : 14Ta10=1/2									
* $^{11}\text{Be}^i$	T : from width=490(70) keV in 97Te07									
* $^{11}\text{C}$	T : from 18Va04 using world data									
* $^{11}\text{N}$	T : from width=780(10) keV in 19We11									
* $^{11}\text{O}$	T : from width=2.31 (0.14) MeV in 20We08; other width=2.46 MeV in 19Fo10									
$^{12}\text{Li}$	49010	30					$(1^-, 2^-)$	17 74Bo05	I 2008	n ?
$^{12}\text{Be}$	25077.8	1.9				21.46 ms 0.05	$0^+$	17	1966	$\beta^-$ =100; $\beta^-$ n=0.50 3
$^{12}\text{Be}^m$	27328.8	2.1	2251	1		233 ns 7	$0^+$	17	2007	IT=100
$^{12}\text{B}$	13369.4	1.3				20.20 ms 0.02	$1^+*$	17	1935	$\beta^-$ =100; $\beta^-$ - $\alpha$ =0.60 2
$^{12}\text{B}^i$	26088	19	12719	19	RQ		$0^+$ T=2	17 08Ch28	J 1919	
$^{12}\text{C}$	0.0	0.0				STABLE	$0^+$	17	1919	IS=98.94 6
$^{12}\text{C}^i$	15108	3	15108	3	RQ		$1^+$ T=1	17		IT=?; $\alpha$ ?
$^{12}\text{C}^j$	27595.0	2.4	27595.0	2.4	RQ		$0^+$ T=2	17		
$^{12}\text{N}$	17338.1	1.0				11.000 ms 0.016	$1^+*$	17 09Hy01	D 1949	$\beta^+$ =100; $\beta^+$ - $\alpha$ =1.93 4
$^{12}\text{N}^i$	29580	4	12242	4	2p	> 5 zs	$(0^+)$ T=2	17 MMC142J		
$^{12}\text{O}$	32013	12				8.9 zs 3.3	$0^+$	17 19We11	T 1978	2p=100
* $^{12}\text{Be}^m$	T : average 07Sh34=229(8) 13Jo06=247(15); other 18Ch31=270(+12-120)									
* $^{12}\text{N}$	T : from 78A101; other 20Bi15=10.92(0.11,stat)(0.01,syst)									
* $^{12}\text{N}^i$	T : from width<100 keV in 19We11									
* $^{12}\text{O}$	T : from width=51(19) keV in 19We11; others 12Ja11<72 keV									
* $^{12}\text{O}$	T : 09Su14=600(500) keV 95Kr03=578(205)keV									
$^{13}\text{Li}$	56980	70				3.3 zs 1.2	$3/2^-$ #	08Ak03	D 2008	2n=100
$^{13}\text{Be}$	33659	10				1.0 zs 0.7	$(1/2^-)$	19Co02	J 1983	n ?
$^{13}\text{Be}^p$	35160	50	1500	50	RQ		$(5/2^+)$		1992	
$^{13}\text{B}$	16561.9	1.0				17.16 ms 0.18	$3/2^-$	00 15Bi05	TD 1956	$\beta^-$ =100; $\beta^-$ n=0.266 36
$^{13}\text{C}$	3125.0093	0.0002				STABLE	$1/2^-$ *	01	1929	IS=1.06 6
$^{13}\text{C}^i$	18233.8	1.1	15108.8	1.1	RQ		$3/2^-$ T=3/2	00		IT=0.82 7;n ?; $\alpha$ ?
$^{13}\text{N}$	5345.48	0.27				9.965 m 0.004	$1/2^-$ *	00	1934	$\beta^+$ =100
$^{13}\text{N}^i$	20410.59	0.18	15065.1	0.3	RQ		$3/2^-$ T=3/2	00		IT=4.9 3;p ?; $\alpha$ ?
$^{13}\text{O}$	23115	10				8.58 ms 0.05	$(3/2^-)$	00 70Es03	D 1963	$\beta^+$ =100; $\beta^+$ -p=10.9 2
$^{13}\text{F}$	42030#	500#					$1/2^+*$			p ?
* $^{13}\text{Li}$	T : from width=125(60-40) keV in 13Ko03									
* $^{13}\text{Be}$	T : from width=450(30) keV in 10Ko17; other width=300(200) keV in 95Pe12									
* $^{13}\text{Be}$	J : from 10Ko17,19Co02; others J=1/2+ in 01Th01,08Ch07,13Ak02,14Ra07									
$^{14}\text{Be}$	39950	130				4.53 ms 0.27	$0^+$	01 15Bi05	TD 1973	$\beta^-$ =100; $\beta^-$ n=86 6; $\beta^-$ 2n=5 2; $\beta^-$ t=0.02 1; $\beta^-$ - $\alpha$ <0.004
$^{14}\text{Be}^p$	41470	60	1520	150	RQ		$(2^+)$	95Bo10	I 1995	
$^{14}\text{B}$	23664	21				12.36 ms 0.29	$2^-$	01 15Bi05	TD 1966	$\beta^-$ =100; $\beta^-$ n=6.04 23; $\beta^-$ 2n ?
$^{14}\text{B}^i$	40728	20	17065	29	RQ	4.15 zs 1.9	$0^+$ T=3	MMC162J	2001	IT ?
$^{14}\text{C}$	3019.893	0.004				5.70 ky 0.03	$0^+$	01	1936	$\beta^-$ =100
$^{14}\text{C}^i$	25120	100	22100	100			$(2^-)$ T=2	01	1989	IT=100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>14</sup> N	2863.4168	0.0002			STABLE	1 <sup>+</sup> *	01	1920	IS=99.6205 247	
<sup>14</sup> N <sup>i</sup>	5176.007	0.010	2312.590	0.010		0 <sup>+</sup> T=1	01 01Ba06	E 1963	IT=100	
<sup>14</sup> O	8007.781	0.025			70.621 s 0.011	0 <sup>+</sup>	01 13La23	T 1949	$\beta^+$ =100	
<sup>14</sup> F	31960	40			500 ys 60	2 <sup>-</sup>	14 10Go16	TJ 2010	p ?	
* <sup>14</sup> Be	D : % $\beta^-$ -t, % $\beta^-$ - $\alpha$ from 02Je11									
* <sup>14</sup> B <sup>i</sup>	T : from width=110(50) keV in 01Ta23									
* <sup>14</sup> O	T : average 13La23=70.610(0.030),70.632(0.094) 12Ta.B=70.623(0.053)									
* <sup>14</sup> O	T : 06Bu06=70.696(0.052) 04Ba78=70.641(0.020) 01Ga59=70.560(0.049)									
* <sup>14</sup> O	T : 78Be61=70.684(0.077) 78Wi04=70.613(0.025) 73Cl12=70.588(0.028);									
* <sup>14</sup> O	T : others (outweighed): 74Az01=70.43(0.18) 72Al01=70.48(0.15)									
* <sup>14</sup> O	T : 72Si50=70.32(0.12)									
* <sup>14</sup> F	T : from width=910(100) keV in 10Go16									
<sup>15</sup> Be	49830	170			790 ys 270	(5/2 <sup>+</sup> )	15 13Sn02	TD 2013	n=100	
<sup>15</sup> B	28957	21			10.18 ms 0.35	3/2 <sup>-</sup>	02 15Bi05	TD 1966	$\beta^-$ =100; $\beta^-$ -n=98.7 10; $\beta^-$ -2n<1.5	
<sup>15</sup> C	9873.1	0.8			2.449 s 0.005	1/2 <sup>+</sup>	02	1950	$\beta^-$ =100	
<sup>15</sup> N	101.4381	0.0006			STABLE	1/2 <sup>-</sup> *	02	1929	IS=0.3795 247	
<sup>15</sup> N <sup>i</sup>	11717	4	11615	4	RQ	1/2 <sup>+</sup> T=3/2	02		n ?;p ?;IT=0.00523 19	
<sup>15</sup> O	2855.6	0.5			122.266 s 0.043	1/2 <sup>-</sup> *	02 20Bu02	T 1934	$\beta^+$ =100	
<sup>15</sup> O <sup>i</sup>	14020#	40#	11165#	35#		(1/2 <sup>+</sup> ) T=3/2	02 Imme	E	p=100	
<sup>15</sup> F	16567	14			1.1 zs 0.3	1/2 <sup>+</sup>	16 04Go15	J 1978	p=100	
<sup>15</sup> Ne	40220	70		2p	770 ys 300	(3/2 <sup>-</sup> )	14 14Wa09	JTD 2014	2p=100	
* <sup>15</sup> Be	T : from width=575(200) keV in 13Sn02									
* <sup>15</sup> B	D : % $\beta^-$ -2n symmetrized from 91Ha25=99.68(+0.08-1.58); other 95Re.A=93.6(1.2)									
* <sup>15</sup> O	T : average 20Bu02=122.308(49) 77Az01=122.23(0.23) 60Ja12=122.1(0.1)									
* <sup>15</sup> F	T : from width=370(70)(+200-0) keV in 16De15									
* <sup>15</sup> Ne	T : from width=590(230) keV									
<sup>16</sup> Be	57450	170			650 ys 130	0 <sup>+</sup>	15 12Sp01	TD 2012	2n=100	
<sup>16</sup> B	37112	25			> 4.6 zs	0 <sup>-</sup> #	16	2000	n ?	
<sup>16</sup> C	13694	4			750 ms 6	0 <sup>+</sup>	99	1961	$\beta^-$ =100; $\beta^-$ -n=99.0 3	
<sup>16</sup> N	5683.9	2.3			7.13 s 0.02	2 <sup>-</sup>	99 18Ki12	D 1933	$\beta^-$ =100; $\beta^-$ - $\alpha$ =0.00154 5	
<sup>16</sup> N <sup>m</sup>	5804.3	2.3	120.42	0.12	5.25 $\mu$ s 0.06	0 <sup>-</sup> T=1	99 83Mi20	D 1957	IT $\approx$ 100; $\beta^-$ =0.000389 25	
<sup>16</sup> N <sup>i</sup>	15613	7	9929	7	RQ	0 <sup>+</sup> T=2	99			
<sup>16</sup> O	-4737.0021	0.0003			STABLE	0 <sup>+</sup>	99	1919	IS=99.757 11	
<sup>16</sup> O <sup>p</sup>	8231.60	0.27	12968.6	0.27		2 <sup>-</sup>	99 64Bo13	E	p=78 4; $\alpha$ =22 4;IT=0.28 3	
<sup>16</sup> O <sup>i</sup>	8059	4	12796	4	RQ	0 <sup>-</sup> T=1	99		IT=100	
<sup>16</sup> O <sup>j</sup>	17984	4	22721	4	RQ	0 <sup>+</sup> T=2	99			
<sup>16</sup> F	10675	5			21 zs 5	0 <sup>-</sup>	99 18Ch25	T 1964	p=100	
<sup>16</sup> Ne	23987	20			> 5.7 zs	0 <sup>+</sup>	99 14Br19	T 1977	2p=100	
* <sup>16</sup> Be	T : from width=0.8(+0.1-0.2) MeV									
* <sup>16</sup> C	T : average 01Gr06=753(8) 76Al02=747(8)									
* <sup>16</sup> N	D : % $\beta^-$ - $\alpha$ average 18Ki12=0.00159(6) 16Re01=0.00149(5stat)(+0-10sys)									
* <sup>16</sup> N <sup>m</sup>	D : % $\beta^-$ - average 83Mi20=4.35(0.50)e-4 5.10(0.65)e-4 83Ga18=3.42(0.37)e-4,									
* <sup>16</sup> N <sup>m</sup>	D : supersedes 82Ga05=3.13(0.43)e-4, 75Pa01=3.3(0.7)e-4									
* <sup>16</sup> F	T : from width=21.3(5.1) keV									
* <sup>16</sup> Ne	T : from width<80 keV (3 sigma upper limit) in 14Br19									
<sup>17</sup> B	43720	200			5.08 ms 0.05	(3/2 <sup>-</sup> )	99 88Du09	D 1973	$\beta^-$ =100; $\beta^-$ -n=63 1; $\beta^-$ -2n=12 2; $\beta^-$ -3n=3.5 7; $\beta^-$ -4n=0.4 3	
<sup>17</sup> C	21032	17			193 ms 6	3/2 <sup>+</sup>	17	1968	$\beta^-$ =100; $\beta^-$ -n=28.4 13; $\beta^-$ -2n ?	
<sup>17</sup> N	7870	15			4.173 s 0.004	1/2 <sup>-</sup>	99 94Do08	D 1949	$\beta^-$ =100; $\beta^-$ -n=95.1 7; $\beta^-$ - $\alpha$ =0.0025 4	
<sup>17</sup> O	-808.7642	0.0006			STABLE	5/2 <sup>+</sup> *	99	1925	IS=0.03835 96	
<sup>17</sup> O <sup>i</sup>	10270.02	0.17	11078.78	0.17	RQ	1/2 <sup>-</sup> T=3/2	99		$\beta^-$ ?;n ?;IT=0.42 14	
<sup>17</sup> F	1951.70	0.25			64.370 s 0.027	5/2 <sup>+</sup>	99 16Br01	T 1934	$\beta^+$ =100	
<sup>17</sup> F <sup>i</sup>	13144.7	1.9	11193.0	1.9	RQ	1/2 <sup>-</sup> T=3/2	99			

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{17}\text{Ne}$	16500.5	0.4			109.2 ms 0.6	$1/2^-*$	99 02Mo19 D	1963	$\beta^+=100; \beta^+p=94.4\ 29;$ $\beta^+\alpha=3.51\ 1; \beta^+p\alpha=0.014\ 4$ $p=100$
$^{17}\text{Na}$	34720	60				$(1/2^+)$	17 17Br07 IJD	2017	
* $^{17}\text{N}$	D : $\% \beta^-n$ from 76Al02								**
* $^{17}\text{F}$	T : average 16Br01=64.402(0.042) 15Gr14=64.347(0.035)								**
$^{18}\text{B}$	51790	200			<26 ns	$(2^-)$	16 10Sp02 IJ	2010	$n=100$
$^{18}\text{C}$	24920	30			92 ms 2	$0^+$	17	1969	$\beta^-=100; \beta^-n=31.5\ 15;$ $\beta^-2n?$
$^{18}\text{N}$	13113	19			619.2 ms 1.9	$1^-$	96 05Li60 TD	1964	$\beta^-=100; \beta^-n=7.0\ 15;$ $\beta^-\alpha=12.2\ 6; \beta^-2n?$ $IS=0.2045\ 102$
$^{18}\text{O}$	-782.8163	0.0006			STABLE	$0^+$	96	1929	
$^{18}\text{O}^i$	15495	20	16278	20		$1^- T=2$	AHWe		*
$^{18}\text{F}$	873.1	0.5			109.734 m 0.008	$1^+$	96 FGK204 T	1937	$\beta^+=100$
$^{18}\text{F}^m$	1994.5	0.5	1121.36	0.15	162 ns 7	$5^+$	96		IT=100
$^{18}\text{F}^i$	1914.7	0.5	1041.55	0.08		$0^+ T=1$	96		IT=100
$^{18}\text{Ne}$	5317.6	0.4			1664.20 ms 0.47	$0^+$	96 15La19 T	1954	$\beta^+=100$
$^{18}\text{Na}$	25040	90			1.3 zs 0.4	$1^- \#$	15 04Ze05 TD	2004	$p=?$
* $^{18}\text{N}$	D : $\% \beta^- \alpha$ from 89Zn04								**
* $^{18}\text{N}$	D : other $\% \beta^-n$ 94Sc01=2.2(0.4) 95Re.A=10.9(0.9) 91Re02=14.3(2.0)(same group)								**
* $^{18}\text{N}$	T : average 05Li60=619(2) 99Og03=620(14) 82O101=624(12) 64Ch19=630(30)								**
* $^{18}\text{O}^i$	E : assuming 16399(5), 17025(10) levels to be IAS's of 114.90(0.18), 747(10)								**
* $^{18}\text{O}^i$	E : levels in $^{18}\text{N}$ (see 95Ti07)								**
* $^{18}\text{Ne}$	T : average 15La19=1664.00(+0.57-0.48) 13Gr03=1664.8(1.1), supersedes								**
* $^{18}\text{Ne}$	T : 07Gr18=1665.6(1.9); others (outweighed): 75Al27=1669(4) 75Ha21=1687(9)								**
$^{19}\text{B}$	59770	530			2.92 ms 0.13	$(3/2^-)$	18 03Yo02 TD	1984	$\beta^-=100; \beta^-n=71\ 9;$ $\beta^-2n=17\ 5; \beta^-3n<9.1$
$^{19}\text{C}$	32410	100			46.2 ms 2.3	$1/2^+$	17 88Du09 TD	1974	$\beta^-=100; \beta^-n=47\ 3; \beta^-2n=7\ 3$
$^{19}\text{N}$	15856	16			336 ms 3	$1/2^-$	96 06Su12 TJD	1968	$\beta^-=100; \beta^-n=41.8\ 9$
$^{19}\text{O}$	3332.9	2.6			26.470 s 0.006	$5/2^+$	96 13Uj01 T	1936	$\beta^-=100$
$^{19}\text{F}$	-1487.4451	0.0008			STABLE	$1/2^+*$	96	1920	IS=100
$^{19}\text{F}^i$	6052.2	0.9	7539.6	0.9		$5/2^+ T=3/2$	96		IT=100
$^{19}\text{Ne}$	1752.05	0.16			17.2569 s 0.0019	$1/2^+*$	96 17Fo24 T	1939	$\beta^+=100$
$^{19}\text{Ne}^i$	9253	9	7501	9 RQ		$(5/2^+) T=3/2$	96 MMC127J		*
$^{19}\text{Na}$	12929	11			> 1 as	$(5/2^+)$	15 10Mu12 T	1969	$p=100$
$^{19}\text{Mg}$	31840	60			5 ps 3	$1/2^- \#$	14 09Mu17 TD	2007	$2p=100$
* $^{19}\text{B}$	D : $\% \beta^-n, \% \beta^-2n$ symmetrized from 03Yo02=71.8(+8.3-9.1), 16.0(+5.6-4.8)								**
* $^{19}\text{C}$	T : average 88Du09=49(4) 95Re.A=44(4) 95Oz02=45.5(4.0)								**
* $^{19}\text{C}$	J : from 01Ma08, 99Na27 and 95Ba28								**
* $^{19}\text{O}$	T : average 13Uj01=26.476(0.009) 94It.A=26.464(0.009)								**
* $^{19}\text{Ne}$	T : average 17Fo24=17.2569(0.0021) 13Uj01=17.254(0.005) 12Tr06=17.262(0.007);								**
* $^{19}\text{Ne}$	T : others (outliers) 14Br06=17.283(0.008) 94Ko.A=17.296(0.005)								**
* $^{19}\text{Ne}$	T : 92Ge08=18.5(0.6) for $q=10+$ (bare ion)								**
* $^{19}\text{Ne}^i$	J : possible IAS of $^{19}\text{O}$ gs ( $J=5/2^+$ )								**
* $^{19}\text{Na}$	T : from width <40 keV in 10Mu12, dominated by resolution of <1 eV								**
* $^{19}\text{Mg}$	T : symmetrized from 09Mu17=6(+2-4); supersedes 07Mu15=4.0(1.5)								**
$^{20}\text{B}$	69400	550			> 912.4 ys	$(1^-, 2^-)$	19 18Le18 TJ	2018	$n=100; \beta^-n?; \beta^-2n?$
$^{20}\text{C}$	37500	230			16 ms 3	$0^+$	17 90Mu06 TD	1981	$\beta^-=100; \beta^-n=70\ 11;$ $\beta^-2n<18.6$
$^{20}\text{N}$	21770	80			136 ms 3	$(2^-)$	18 06Su12 TD	1969	$\beta^-=100; \beta^-n=42.9\ 14;$ $\beta^-2n?$
$^{20}\text{O}$	3796.2	0.9			13.51 s 0.05	$0^+$	98	1959	$\beta^-=100$
$^{20}\text{F}$	-17.463	0.030			11.0062 s 0.0080	$2^+$	98 19Bu01 T	1935	$\beta^+=100$
$^{20}\text{F}^i$	6503	3	6521	3 RQ		$0^+ T=2$	98		*
$^{20}\text{Ne}$	-7041.9322	0.0015			STABLE	$0^+$	98	1913	IS=90.48 3
$^{20}\text{Ne}^i$	3230.5	2.0	10272.5	2.0 RQ		$2^+ T=1$	98		IT=100
$^{20}\text{Ne}^j$	9690.9	2.8	16732.8	2.8 RQ		$0^+ T=2$	98		IT=100
$^{20}\text{Na}$	6850.5	1.1			447.9 ms 2.3	$2^+*$	98 89Cl02 D	1950	$\beta^+=100; \beta^+\alpha=25.0\ 4$



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{20}\text{Na}^i$	13348.9	1.2	6498.4	0.5	p	$0^+ T=2$	98	1979	p=100
$^{20}\text{Mg}$	17477.7	1.9			90.4 ms 0.5	$0^+$	16 17Su05 T	1974	$\beta^+=100; \beta^+p=30.3$ 12
* $^{20}\text{B}$	T : from width < 0.5 MeV								**
* $^{20}\text{C}$	D : % $\beta^-$ -n average 03Yo02=65(+19-18) 90Mu06=72(14)								**
* $^{20}\text{C}$	T : average 90Mu06=14(+6-5), supersedes 89Le16=16(+14-7) same group,								**
* $^{20}\text{C}$	T : 95Re.A=16.7(3.5); other 03Yo02=21.8(+15.0-7.4)								**
* $^{20}\text{F}$	T : evaluated in 19Bu01 using the world data								**
* $^{20}\text{Mg}$	T : average 17Su05=90.0(0.6) 16Lu13=91.4(1.0)								**
$^{21}\text{B}$	78380	560			> 760 ys	$(3/2^-)$	19 18Le18 TJI	2018	2n=100
$^{21}\text{C}$	45640#	600#			<30ns	$1/2^+\#$	15 93Po.A I		n ?
$^{21}\text{N}$	25230	130			85 ms 5	$(1/2^-)$	15 09Li51 TD	1970	$\beta^-=100; \beta^-n=87$ 3; $\beta^-2n$ ?
$^{21}\text{O}$	8062	12			3.42 s 0.10	$(5/2^+)$	15	1968	$\beta^-=100; \beta^-n$ ?
$^{21}\text{F}$	-47.6	1.8			4.158 s 0.020	$5/2^+$	15	1955	$\beta^-=100$
$^{21}\text{Ne}$	-5731.78	0.04			STABLE	$3/2^+*$	15	1928	IS=0.27 1
$^{21}\text{Ne}^i$	3129.0	0.3	8860.8	0.3		$5/2^+ T=3/2$	15		
$^{21}\text{Na}$	-2184.86	0.04			22.4550 s 0.0054	$3/2^+*$	15 18Sh27 T	1940	$\beta^+=100$
$^{21}\text{Na}^i$	6790	4	8975	4	p	$5/2^+ T=3/2$	15		
$^{21}\text{Mg}$	10903.9	0.8			120.0 ms 0.4	$5/2^+*$	15 15Lu13 D	1963	$\beta^+=100; \beta^+p=20.1$ 21; $\beta^+ \alpha=0.116$ 18; $\beta^+p \alpha=0.016$ 3
$^{21}\text{Al}$	27090#	600#			<35ns	$5/2^+\#$	15 93Po.A I		p ?
* $^{21}\text{B}$	T : from width < 0.6 MeV								**
* $^{21}\text{N}$	T : average 09Li51=83(8), supersedes 08Lo06=82.9(1.9), 07Su05=85(14)								**
* $^{21}\text{N}$	T : 90Mu06=95(+15-11) 95Re.A=83.6(6.7), supersedes 91Re02=61(23)								**
* $^{21}\text{N}$	D : % $\beta^-$ -n average 09Li51=90.5(4.2) 90Mu06=84(9) 95Re.A=78(7), supersedes								**
* $^{21}\text{N}$	D : 91Re02=76(15)								**
* $^{21}\text{Na}$	T : average 18Sh27=22.4615(0.0040) 17Fi07=22.4056(0.0033); others								**
* $^{21}\text{Na}$	T : 15Gr05=22.422(0.010) 75Az01=22.47(0.03) 74Al03=22.55(0.10)								**
* $^{21}\text{Mg}$	T : average 15Lu12=118.6(0.5) 18Wa20=121.9(0.6) 92Go10=120(5)								**
* $^{21}\text{Mg}$	T : 73Se08=123(3) 65Ha20=121(5)								**
* $^{21}\text{Mg}$	D : % $\beta^+p$ average 18Wa20=19.2(30) 15Lu13=21.0(3.0)								**
$^{22}\text{C}$	53610	230			6.2 ms 1.3	$0^+$	15 03Yo02 TD	1986	$\beta^-=100; \beta^-n=61$ 14; $\beta^-2n<37$
$^{22}\text{N}$	31760	210			23 ms 3	$0^-\#$	15	1979	$\beta^-=100; \beta^-n=34$ 3; $\beta^-2n=12$ 3
$^{22}\text{O}$	9280	60			2.25 s 0.09	$0^+$	15	1969	$\beta^-=100; \beta^-n<22$
$^{22}\text{F}$	2793	12			4.23 s 0.04	$(4^+)$	15	1965	$\beta^-=100; \beta^-n<11$
$^{22}\text{Ne}$	-8024.716	0.018			STABLE	$0^+$	15	1913	IS=9.25 3
$^{22}\text{Ne}^i$	5855	10	13880	10		$4^+ T=2$	15 04Go03 E		*
$^{22}\text{Na}$	-5181.39	0.13			2.6019 y 0.0006	$3^+*$	15 FGK204 T	1935	$\beta^+=100; e^+=90.57$ 8; $\epsilon=9.43$ 6
$^{22}\text{Na}^m$	-4598.34	0.16	583.05	0.10	243 ns 2	$1^+$	15	1958	IT=100
$^{22}\text{Na}^i$	-4524.39	0.19	657.00	0.14	19.6 ps 0.7	$0^+ T=1$	15		IT=100
$^{22}\text{Mg}$	-399.99	0.16			3.8745 s 0.0007	$0^+$	15 17Du11 T	1961	$\beta^+=100$
$^{22}\text{Mg}^i$	13644	6	14044	6	p	$(4)^+ T=2$	15 MMC12 J		$\alpha=?; p=?$
$^{22}\text{Al}$	18200#	400#			91.1 ms 0.5	$(4)^+$	15	1982	$\beta^+=100; \beta^+p=55$ 3; $\beta^+2p=1.10$ 11; $\beta^+ \alpha=0.038$ 17
$^{22}\text{Si}$	33640#	500#			28.7 ms 1.1	$0^+$	15 20Le16 TD	1987	$\beta^+=100; \beta^+p=62$ 5; $\beta^+2p=0.7$ 3
* $^{22}\text{C}$	T : symmetrized from 03Yo02=6.1(+1.4-1.2)								**
* $^{22}\text{C}$	D : % $\beta^-$ -n symmetrized from 03Yo02=61(+14-13)								**
* $^{22}\text{Ne}^i$	E : from 16Ma.A, but T=2 assignment is not firm								**
* $^{22}\text{Na}$	D : from 71GoYM								**
* $^{22}\text{Mg}$	T : average 17Du11=3.87400(0.00079) 03Ha20=3.8755(0.0012)								**
* $^{22}\text{Mg}^i$	J : IAS of $^{22}\text{Al}$ gs [J=(4)+]								**
* $^{22}\text{Si}$	T : average 20Le16=28.6(1.4), supersedes 17Xu01=27.8(3.5), 96B111=29(2)								**
* $^{22}\text{Si}$	D : % $\beta^+p$ from 20Le16, based on %I(p)=5.3(1.0), 43.0(4.6) and 13.5(2.1);								**
* $^{22}\text{Si}$	D : % $\beta^+2p$ from 17Xu01								**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
<sup>23</sup> C	64170#	1000#				3/2 <sup>+</sup> #			n ?
<sup>23</sup> N	36720	420			13.9 ms 1.4	1/2 <sup>-</sup> #	21 03Yo02	TD 1985	$\beta^- = 100; \beta^- n = 42.6; \beta^- 2n = 8.4; \beta^- 3n < 3.4$ *
<sup>23</sup> O	14620	120			97 ms 8	1/2 <sup>+</sup>	21 07Su05	TD 1970	$\beta^- = 100; \beta^- n = 7.2$
<sup>23</sup> F	3290	30			2.23 s 0.14	5/2 <sup>+</sup>	21 95Re.A	D 1970	$\beta^- = 100; \beta^- n < 14$
<sup>23</sup> Ne	-5154.05	0.10			37.15 s 0.03	5/2 <sup>+</sup> *	21 15La19	T 1936	$\beta^- = 100$ *
<sup>23</sup> Na	-9529.8535	0.0018			STABLE	3/2 <sup>+</sup> *	21	1921	IS=100 *
<sup>23</sup> Na <sup>i</sup>	-1638.7	0.3	7891.2	0.3		5/2 <sup>+</sup> T=3/2	21		IT=100
<sup>23</sup> Na <sup>j</sup>	10060.6	2.0	19590.4	2.0	240 zs 120	T=5/2	21 85Ev01	T	*
<sup>23</sup> Mg	-5473.67	0.03			11.3039 s 0.0032	3/2 <sup>+</sup> *	21 17Yo05	J 1939	$\beta^+ = 100$
<sup>23</sup> Mg <sup>i</sup>	2329.3	0.6	7803.0	0.6		5/2 <sup>+</sup> T=3/2	21 00Pe28	D 1981	IT≈100; p=0.17 8
<sup>23</sup> Al	6748.1	0.3			446 ms 6	5/2 <sup>+</sup>	21 06Ia03	T 1969	$\beta^+ = 100; \beta^+ p = 1.22 5$ *
<sup>23</sup> Al <sup>i</sup>	18470	40	11720	40		(5/2 <sup>+</sup> ) <sup>+</sup> T=5/2	21	1997	p=0.10 5; 2p=3.6 4
<sup>23</sup> Si	23950#	500#			42.3 ms 0.4	3/2 <sup>+</sup> #	21 97Bl04	TD 1986	$\beta^+ = 100; \beta^+ p \approx 88; \beta^+ 2p = 3.6 3$ *
* <sup>23</sup> N	T : symmetrized from 03Yo02=14.1(+1.2-1.5) **								
* <sup>23</sup> N	D : % $\beta^- n$ and % $\beta^- 2n$ symmetrized from 03Yo02=42.2(+6.3-6.5) and 8.0(+3.8-3.4) **								
* <sup>23</sup> Ne	T : average 15La19=37.148(0.032) 07Gr18=37.11(0.06) 74Al03=37.24(0.12) **								
* <sup>23</sup> Na	J : 00Ke09=5/2 **								
* <sup>23</sup> Na <sup>j</sup>	T : from width=1.9(0.8) keV **								
* <sup>23</sup> Mg	T : average 77Az01=11.317(0.011) 17Ma18=11.3027(0.0033) **								
* <sup>23</sup> Al	D : from 11Sa15 **								
* <sup>23</sup> Si	T : also 18Wa05=40.17(1.86) for all delayed proton event > 300 keV **								
<sup>24</sup> N	46940#	400#			<52ns		07 93Po.A	I	n ?
<sup>24</sup> O	18500	160			77.4 ms 4.5	0 <sup>+</sup>	07 15Ca09	TD 1970	$\beta^- = 100; \beta^- n = 43.4$ *
<sup>24</sup> F	7540	100			384 ms 16	3 <sup>+</sup>	07 07Su05	T 1970	$\beta^- = 100; \beta^- n < 5.9$ *
<sup>24</sup> Ne	-5951.6	0.5			3.38 m 0.02	0 <sup>+</sup>	07	1956	$\beta^- = 100; [gs=0, m=100]$
<sup>24</sup> Na	-8417.901	0.017			14.9560 h 0.0015	4 <sup>+</sup> *	07 FGK204	T 1934	$\beta^- = 100$ *
<sup>24</sup> Na <sup>m</sup>	-7945.694	0.017	472.2074	0.0008	20.18 ms 0.10	1 <sup>+</sup>	07	1961	IT≈100; $\beta^- = 0.05$
<sup>24</sup> Na <sup>i</sup>	-2450.53	0.13	5967.37	0.13		0 <sup>+</sup> T=2	07		
<sup>24</sup> Mg	-13933.578	0.013			STABLE	0 <sup>+</sup>	07	1920	IS=78.965 49
<sup>24</sup> Mg <sup>i</sup>	-4417.30	0.04	9516.28	0.04		(4 <sup>+</sup> ) T=1	07		
<sup>24</sup> Mg <sup>j</sup>	1502.8	0.6	15436.4	0.6		0 <sup>+</sup> T=2	07		
<sup>24</sup> Al	-48.81	0.23			2.053 s 0.004	4 <sup>+</sup>	07	1953	$\beta^+ = 100; \beta^+ \alpha = 0.035 6; \beta^+ p = 0.0016 3$
<sup>24</sup> Al <sup>m</sup>	376.99	0.25	425.8	0.1	130 ms 3	1 <sup>+</sup>	07	1968	IT=82.5 30; $\beta^+ = 17.5 30; \beta^+ \alpha = 0.028 6$
<sup>24</sup> Al <sup>i</sup>	5900	3	5949	3		0 <sup>+</sup> T=2	07		
<sup>24</sup> Si	10745	19			143.2 ms 2.1	0 <sup>+</sup>	07 15Su15	T 1979	$\beta^+ = 100; \beta^+ p = 34.5 14$ *
<sup>24</sup> P	34020#	500#				1 <sup>+</sup> #			p ?; $\beta^+ ?; \beta^+ p ?$
* <sup>24</sup> O	T : average 15Ca09=80(5) 01Pe14=67(10); other 90Mu06=61(+32-19) **								
* <sup>24</sup> F	J : 15Ca09=3 **								
* <sup>24</sup> Na	J : 00Ke09=4 **								
* <sup>24</sup> Si	T : average 15Su15=143.3(2.2) 97Cz02=140(8) **								
* <sup>24</sup> Si	D : % $\beta^+ p$ average 98Cz01= 37.6(2.5) 11Ic06=33.3(1.6) **								
<sup>25</sup> N	55980#	500#			<260ns	1/2 <sup>-</sup> #	09 99Sa06	I	n ?; 2n ?; $\beta^- ?$ *
<sup>25</sup> O	27330	170			5.18 zs 0.35	3/2 <sup>+</sup> #	09 16Ko11	T 2008	n=100 *
<sup>25</sup> F	11330	100			80 ms 9	(5/2 <sup>+</sup> )	09	1970	$\beta^- = 100; \beta^- n = 23.1 45; \beta^- 2n ?$
<sup>25</sup> Ne	-2036	29			602 ms 8	1/2 <sup>+</sup> *	09	1970	$\beta^- = 100$
<sup>25</sup> Na	-9357.8	1.2			59.1 s 0.6	5/2 <sup>+</sup> *	09	1943	$\beta^- = 100$ *
<sup>25</sup> Mg	-13192.78	0.05			STABLE	5/2 <sup>+</sup> *	09	1920	IS=10.011 13 *
<sup>25</sup> Mg <sup>i</sup>	-5405.8	0.3	7787.0	0.3		5/2 <sup>+</sup> T=3/2	09		
<sup>25</sup> Al	-8915.97	0.06			7.1666 s 0.0023	5/2 <sup>+</sup>	09 17Lo09	T 1953	$\beta^+ = 100$ *
<sup>25</sup> Al <sup>i</sup>	-1014.9	1.8	7901.1	1.8		5/2 <sup>+</sup> T=3/2	09 20Su.1	E	
<sup>25</sup> Si	3827	10			220.6 ms 1.0	5/2 <sup>+</sup>	09 20Su.1	T 1963	$\beta^+ = 100; \beta^+ p = 35 2$ *
<sup>25</sup> P	20190#	400#			<30ns	1/2 <sup>+</sup> #	09 93Po.A	I	p ?
* <sup>25</sup> N	I : 240 <sup>25</sup> N events expected, but none observed in 99Sa06 **								
* <sup>25</sup> O	T : from width=88(6) keV in 16Ko11; other width=20(+60-20) keV in 13Ca18 **								

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>25</sup> Na	J : 00Ke09=5/2								**
* <sup>25</sup> Mg	J : also 17Yo05,19Yo06=5/2								**
* <sup>25</sup> Al	T : average 17Lo09=7.1657(0.0024) 75Az01=7.174(0.007)								**
* <sup>25</sup> Si	T : symmetrized from 20Su.1=220.9(+0.8-1.2)								**
<sup>26</sup> O	34660	160			4.2 ps 3.3	0 <sup>+</sup>	16 13Ko10 T	2012	2n=100
<sup>26</sup> F	18670	110			8.2 ms 0.9	1 <sup>+</sup>	16	1979	$\beta^-$ =100; $\beta^-$ n=13.5 40; $\beta^-$ 2n ?
<sup>26</sup> F <sup>m</sup>	19310	110	643.4	0.1	2.2 ms 0.1	(4 <sup>+</sup> )	16	2013	IT=82 11; $\beta^-$ =?; $\beta^-$ n=12 8
<sup>26</sup> Ne	481	18			197 ms 2	0 <sup>+</sup>	16	1970	$\beta^-$ =100; $\beta^-$ n=0.13 3
<sup>26</sup> Na	-6861	4			1071.28 ms 0.25	3 <sup>+</sup> *	16	1958	$\beta^-$ =100
<sup>26</sup> Na <sup>m</sup>	-6779	4	82.4	0.04	4.35 $\mu$ s 0.16	1 <sup>+</sup>	16 14NiZZ ET	1987	IT=100
<sup>26</sup> Mg	-16214.544	0.029			STABLE	0 <sup>+</sup>	16	1920	IS=11.025 38
<sup>26</sup> Al	-12210.14	0.07			717 ky 24	5 <sup>+</sup>	16	1934	$\beta^+$ =100
<sup>26</sup> Al <sup>m</sup>	-11981.83	0.07	228.306	0.013 MD	6346.0 ms 0.5	0 <sup>+</sup> T=1	16 13Ch51 T	1934	$\beta^+$ =100
<sup>26</sup> Si	-7141.00	0.11			2.2453 s 0.0007	0 <sup>+</sup>	16 10Ia01 T	1960	$\beta^+$ =100
<sup>26</sup> Si <sup>i</sup>	5874	4	13015	4 p		(3 <sup>+</sup> ) T=2	16		
<sup>26</sup> P	10970#	200#			43.6 ms 0.3	(3 <sup>+</sup> ) <sup>+</sup>	16 17Ja05 D	1983	$\beta^+$ =100; $\beta^+$ p=35.1 14; $\beta^+$ 2p=1.99 21
<sup>26</sup> P <sup>m</sup>	11130#	200#	164.4	0.1	115 ns 8	(1 <sup>+</sup> )	16 17Pe09 ET	2014	IT=100
<sup>26</sup> S	27680#	600#			<79ns	0 <sup>+</sup>	16 11Fo08 IT		2p ?
* <sup>26</sup> O	T : symmetrized from 13Ko10=4.5(+1.1-1.5 stat)(3 syst)								**
* <sup>26</sup> Na	J : 00Ke09=1								**
* <sup>26</sup> Na <sup>m</sup>	T : also 87DuZU=9(2)								**
* <sup>26</sup> Al <sup>m</sup>	T : average 13Ch51=6345.30(0.90) 11Fi01=6346.54(0.46,stat)(0.60,syst)								**
* <sup>26</sup> Al <sup>m</sup>	T : 11Sc22=6347.8(2.5) 83Ko22=6346.2(2.6) 77Al11=6339.5(4.5)								**
* <sup>26</sup> Al <sup>m</sup>	T : 75Az01=6346(5) 72Ha82=6351(10) 69Fr08=6346(5)								**
* <sup>26</sup> Si	T : other 08Ma39=2.2283(0.0027), discrepant; see discussions in 10Ia01								**
* <sup>26</sup> P	D : % $\beta^+$ 2p average 17Ja05=1.5(0.4) 04Th09=2.16(0.24);								**
* <sup>26</sup> P	D : % $\beta^+$ p + % $\beta^+$ 2p average 17Ja05=35(2) 04Th09=39(2)								**
* <sup>26</sup> P	T : average 20Li06=43.6(0.3) 07Th09=43.7(0.6);								**
* <sup>26</sup> P	T : others: 17Ja05=50(+23-12) 83Ca06,84Ca29=20(+35-15)								**
* <sup>26</sup> P <sup>m</sup>	T : average 14NiZZ=120(9) 17Pe09=104(14)								**
<sup>27</sup> O	44670#	500#			<260ns	3/2 <sup>+</sup> #	99Sa06 I		n ?;2n ?
<sup>27</sup> F	25130	120			5.0 ms 0.2	5/2 <sup>+</sup> #	11	1981	$\beta^-$ =100; $\beta^-$ n=77 21; $\beta^-$ 2n ?
<sup>27</sup> Ne	7050	90			30.9 ms 1.1	(3/2 <sup>+</sup> )	11 17Ha23 T	1977	$\beta^-$ =100; $\beta^-$ n=2.0 5; $\beta^-$ 2n ?
<sup>27</sup> Na	-5518	4			301 ms 6	5/2 <sup>+</sup> *	11	1968	$\beta^-$ =100; $\beta^-$ n=0.098 24
<sup>27</sup> Mg	-14586.59	0.05			9.435 m 0.027	1/2 <sup>+</sup> *	11 15ZaZY T	1934	$\beta^-$ =100
<sup>27</sup> Al	-17196.86	0.05			STABLE	5/2 <sup>+</sup> *	11	1922	IS=100
<sup>27</sup> Al <sup>i</sup>	-10383.1	0.7	6813.8	0.7		1/2 <sup>+</sup> T=3/2	11		IT=100
<sup>27</sup> Si	-12384.51	0.11			4.117 s 0.014	5/2 <sup>+</sup>	11 17Ma18 T	1939	$\beta^+$ =100
<sup>27</sup> Si <sup>i</sup>	-5759.5	2.3	6625.0	2.3 RQ		1/2 <sup>+</sup> T=3/2	11	1977	IT ?
<sup>27</sup> P	-659	9			260 ms 80	1/2 <sup>+</sup>	11	1977	$\beta^+$ =100; $\beta^+$ p $\approx$ 0.07
<sup>27</sup> P <sup>i</sup>	12010	30	12670	30 p		5/2 <sup>+</sup> T=5/2	11	1991	IT ?
<sup>27</sup> S	17490#	400#			16.3 ms 0.2	(5/2 <sup>+</sup> )	11 19Su14 T	1986	$\beta^+$ =100; $\beta^+$ p=61 3; $\beta^+$ 2p=3.0 6
* <sup>27</sup> F	T : average 99Re16=6.5(1.1) 97Ta22=5.3(0.9) 99Di01=5.2(0.3) 98NoZW=4.9(0.2)								**
* <sup>27</sup> F	D : % $\beta^-$ n symmetrized from 99Re16=90(+10-30)								**
* <sup>27</sup> Ne	T : average 17Ha23=29.3(2.1) 06Tr02=31.5(1.3)								**
* <sup>27</sup> Na	J : 00Ke09=5/2								**
* <sup>27</sup> Na	D : % $\beta^-$ n average 84Gu19=0.13(0.04) 74Ro31=0.08(0.03)								**
* <sup>27</sup> Mg	T : average 15ZaZY=9.408 (0.012) 70Re13=9.462 (0.012); Birge ratio=3.18								**
* <sup>27</sup> Si	T : average 17Ma18=4.1117(0.0020) 75Az01=4.109 (0.004) 76Ge06=4.206(0.008);								**
* <sup>27</sup> Si	T : Birge ratio=8.19								**
* <sup>27</sup> S	T : others 17Ja05=15.5(1.6) 01Ca60=15.5(1.5)								**
* <sup>27</sup> S	D : % $\beta^+$ p deduced from % $\beta^+$ p + % $\beta^+$ 2p = 64(3) and % $\beta^+$ 2p=3.0(0.6)								**
<sup>28</sup> O	52080#	700#			<100ns	0 <sup>+</sup>	13 98Po.A I		2n ?; $\beta^-$ =0
<sup>28</sup> F	33400	120			46 zs	(4 <sup>-</sup> )	13 20Re06 JD	2012	n=100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{28}\text{Ne}$	11300	130			18.8 ms 0.2	$0^+$	13 17Ha23 T	1979	$\beta^- = 100; \beta^- n = 12.1;$ $\beta^- 2n = 3.7.5$	*
$^{28}\text{Na}$	-988	10			33.1 ms 1.3	$1^+*$	13 17Ha23 T	1969	$\beta^- = 100; \beta^- n = 0.58.12$	*
$^{28}\text{Mg}$	-15019.95	0.26			20.915 h 0.009	$0^+$	13	1953	$\beta^- = 100$	
$^{28}\text{Al}$	-16850.72	0.05			2.245 m 0.005	$3^+*$	13 20He.A J	1934	$\beta^- = 100$	
$^{28}\text{Al}^i$	-10858.14	0.11	5992.58	0.10		$0^+ T=2$	13			
$^{28}\text{Si}$	-21492.7971	0.0005			STABLE	$0^+$	13	1920	IS=92.2545.37	
$^{28}\text{Si}^r$	-8951.75	0.05	12541.04	0.05	RQ	$(3^+)$	13			
$^{28}\text{Si}^i$	-12176.88	0.10	9315.92	0.10		$3^+ T=1$	13			
$^{28}\text{Si}^j$	-6265.8	1.0	15227	1		$(0^+) T=2$	13 68Mc12 D	1968	$\alpha = 90.11; p = 10.11$	
$^{28}\text{P}$	-7147.9	1.1			270.3 ms 0.5	$3^+$	13 79Ho27 D	1953	$\beta^+ = 100; \beta^+ p = 0.0013.4;$ $\beta^+ \alpha = 0.00086.25$	
$^{28}\text{P}^i$	-1261	20	5887	20	p	$0^+ T=2$	13			
$^{28}\text{S}$	4070	160			125 ms 10	$0^+$	13	1982	$\beta^+ = 100; \beta^+ p = 20.7.19$	
$^{28}\text{Cl}$	28270#	500#			>100ns	$1^+\#$	18Mu18 TD	2018	p=100	
* $^{28}\text{O}$	I: also 11 and 37 $^{28}\text{O}$ events expected in 97Ta22 and 99Sa06,									**
* $^{28}\text{O}$	I: respectively, but none observed									**
* $^{28}\text{Ne}$	T: average 17Ha23=19.2(0.6) 15Le17=18.6(0.2) 06Tr02=20.0(0.5)									**
* $^{28}\text{Na}$	T: unweighted average 17Ha23=34.6(1.0) 84Gu19=34.1(0.6) 74Ro31=30.5(0.4);									**
* $^{28}\text{Na}$	T: Birge ratio=4.06									**
* $^{28}\text{Na}$	J: 00Ke09=1									**
$^{29}\text{F}$	40150	530			2.5 ms 0.3	$(5/2^+)$	12 17Ma77 J	1989	$\beta^- = 100; \beta^- n = 60.40; \beta^- 2n ?$	*
$^{29}\text{Ne}$	18400	150			14.7 ms 0.4	$(3/2^-)$	12 05Tr13 T	1985	$\beta^- = 100; \beta^- n = 28.5; \beta^- 2n = 4.1$	*
$^{29}\text{Na}$	2680	7			43.2 ms 0.4	$3/2^+*$	12 17Ha23 T	1969	$\beta^- = 100; \beta^- n = 22.3; \beta^- 2n ?$	*
$^{29}\text{Mg}$	-10612.4	0.3			1.30 s 0.12	$3/2^+*$	12	1971	$\beta^- = 100$	*
$^{29}\text{Al}$	-18207.8	0.3			6.56 m 0.06	$5/2^+*$	12 20He.A J	1939	$\beta^- = 100$	
$^{29}\text{Si}$	-21895.0815	0.0006			STABLE	$1/2^+*$	12	1920	IS=4.672.16	
$^{29}\text{Si}^i$	-13605	5	8290	5		$5/2^+ T=3/2$	12		IT=100	
$^{29}\text{P}$	-16952.8	0.4			4.102 s 0.004	$1/2^+$	12 20Lo01 T	1941	$\beta^+ = 100$	*
$^{29}\text{P}^i$	-8571.0	2.5	8381.8	2.4	RQ	$5/2^+ T=3/2$	12	1969	IT=100	
$^{29}\text{S}$	-3094	13			188 ms 4	$5/2^+\#$	12 79Vi01 D	1964	$\beta^+ = 100; \beta^+ p = 46.4.10$	
$^{29}\text{Cl}$	14020#	190#			5.4 zs 1.9	$(1/2^+)$	16 15Mu13 I	1993	p=100	*
$^{29}\text{Ar}$	37970#	440#			>100ns	$5/2^+\#$	18Mu18 TD	2018	2p=100	
* $^{29}\text{F}$	D: % $\beta^- n$ from 99DI01, 01Pe14=100(80)									**
* $^{29}\text{Ne}$	T: average 05Tr13=13.8(0.5) 06Tr02=15.1(2.6) 97No.A=15.6(0.5); others:									**
* $^{29}\text{Ne}$	T: 06Tr02=16.4(1.3) 01Pe14=15(3) 99DI01=15(4) 99Re16=19(9) 97Ta22=15(3)									**
* $^{29}\text{Ne}$	J: 16Ko05=(3/2)									**
* $^{29}\text{Ne}$	D: % $\beta^- n$ average 06Tr02=29(7) 99Re16, 99DI01=27(9) 01Pe14=27(9);									**
* $^{29}\text{Ne}$	D: other 01Be53=17.5									**
* $^{29}\text{Na}$	D: % $\beta^- n$ average 95Re.A=27.1(1.6) 84La03=21.5(3.0) 74Ro31=15.1(1.8)									**
* $^{29}\text{Na}$	D: 79De02=21(4); Birge ratio=2.88									**
* $^{29}\text{Na}$	T: average 17Ha23=42.8(0.5) 84Gu19=44.9(1.2) 84La03=44(1) 74Ro31=42.9(1.5)									**
* $^{29}\text{Na}$	J: 00Ke09=3/2									**
* $^{29}\text{Mg}$	J: also 19Yo06=5/2									**
* $^{29}\text{P}$	T: average 20Lo01=4.1055(0.0044) 80Wi13=4.084(0.022) 75Az01=4.083(0.012);									**
* $^{29}\text{P}$	T: other (not used) 73Ta04=4.149(0.005)									**
* $^{29}\text{Cl}$	T: from width=85(30) keV in 16Go.1									**
$^{30}\text{F}$	48960#	500#			<260ns		10 99Sa06 I		n ?	
$^{30}\text{Ne}$	23280	250			7.22 ms 0.18	$0^+$	10 15St14 T	1985	$\beta^- = 100; \beta^- n = 13.4;$ $\beta^- 2n = 8.9.23$	*
$^{30}\text{Na}$	8475	5			45.9 ms 0.7	$2^+*$	10 17Ha23 T	1969	$\beta^- = 100; \beta^- n = 28.6.22;$ $\beta^- 2n = 1.24.19;$ $\beta^- \alpha = 5.5e-5.2$	*
$^{30}\text{Mg}$	-8881.4	1.3			317 ms 4	$0^+$	10 84La03 D	1971	$\beta^- = 100; \beta^- n < 0.06$	*
$^{30}\text{Al}$	-15864.1	1.9			3.62 s 0.06	$3^+*$	10 20He.A J	1961	$\beta^- = 100$	
$^{30}\text{Si}$	-24432.962	0.022			STABLE	$0^+$	10	1924	IS=3.0735.21	
$^{30}\text{P}$	-20200.86	0.07			2.5000 m 0.0017	$1^+* T=0$	10 18Ia01 T	1934	$\beta^+ = 100$	*
$^{30}\text{P}^i$	-19523.85	0.08	677.01	0.03		$0^+ T=1$	10			
$^{30}\text{S}$	-14059.25	0.21			1.1798 s 0.0003	$0^+$	10 18Ia01 T	1961	$\beta^+ = 100$	*

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
<sup>30</sup> Cl	4675	24			>100ns	3 <sup>+</sup> #	10 18Mu18	TD 2018	p=100
<sup>30</sup> Ar	22070#	180#			< 10 ps	0 <sup>+</sup>	16 15Mu13	IDT 2015	2p=100
* <sup>30</sup> Ne	T : average 15St14=7.18(0.22) 07Tr08=7.3(0.3);								**
* <sup>30</sup> Ne	T : others 01Pe14=7(2) 99Dl01=7.5(1.5)								**
* <sup>30</sup> Na	T : average 17Ha23=44.1(0.8) 84La02=48(2) 84Gu19=50(3) 79De02=53(3)								**
* <sup>30</sup> Na	T : 74Ro01=53(3) 99Dl01=50(4) 97Ta22=48(5)								**
* <sup>30</sup> Na	J : 00Ke09=2								**
* <sup>30</sup> Na	D : Pn=32.2(2.6), average 84La03=33(5) 84Gu19=30(5) 74Ro01=33.1(3.8);								**
* <sup>30</sup> Na	D : %β <sup>-</sup> 2n average 80De26=1.15(0.25) and 1.35(28), from Pn=32.2(2.6) and								**
* <sup>30</sup> Na	D : P2n/Pn=0.042(0.008) in 80De26. %β <sup>-</sup> n average 79De02=26(4) and 29.7(2.6)								**
* <sup>30</sup> Na	D : from Pn-2*β <sup>-</sup> 2n=32.2(2.6)-2*1.24(0.19); %β <sup>-</sup> α from 83De23								**
* <sup>30</sup> Mg	T : average 17Ha23=311(8) 16Ol06=335(10) 08Hi05=314(5)								**
* <sup>30</sup> P	T : average 18Ia01=2.501(0.002) 80Wi13=2.498(0.004) 63Mc02=2.497(0.005)								**
* <sup>30</sup> S	T : average 18Ia01=1.17992(0.00034) 11So11 = 1.1759(0.0017)								**
<sup>31</sup> F	56840#	540#			2# ms >260ns	5/2 <sup>+</sup> #	13 99Sa06	I 1999	β <sup>-</sup> ?; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>31</sup> Ne	31180	270			3.4 ms 0.8	(3/2 <sup>-</sup> )	13		β <sup>-</sup> =100; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>31</sup> Na	12246	14			16.8 ms 0.3	3/2 <sup>+</sup> *	13 19Ni04	D 1969	β <sup>-</sup> =100; β <sup>-</sup> n=36.0 35; β <sup>-</sup> 2n=0.73 9; β <sup>-</sup> 3n<0.05
<sup>31</sup> Mg	-3122	3			270 ms 2	1/2 <sup>+</sup> *	13 17Ha23	T 1977	β <sup>-</sup> =100; β <sup>-</sup> n=6.2 19
<sup>31</sup> Al	-14950.7	2.2			644 ms 25	5/2 <sup>+</sup> *	13 16He09	J 1971	β <sup>-</sup> =100; β <sup>-</sup> n<1.6
<sup>31</sup> Si	-22949.04	0.04			157.16 m 0.20	3/2 <sup>+</sup>	13 17Da28	T 1934	β <sup>-</sup> =100
<sup>31</sup> P	-24440.5444	0.0007			STABLE	1/2 <sup>+</sup> *	13		IS=100
<sup>31</sup> P <sup>i</sup>	-18059.7	2.0	6380.8	2.0		3/2 <sup>+</sup> T=3/2	13		IT=100
<sup>31</sup> S	-19042.53	0.23			2.5534 s 0.0018	1/2 <sup>+</sup>	13		β <sup>+</sup> =100
<sup>31</sup> S <sup>i</sup>	-12761.9	0.6	6280.60	0.60		3/2 <sup>+</sup> T=3/2	13		
<sup>31</sup> Cl	-7035	3			190 ms 1	3/2 <sup>+</sup>	13		β <sup>+</sup> =100; β <sup>+</sup> p=2.4 2
<sup>31</sup> Cl <sup>i</sup>	5256	3	12291	5 RQ		3/2 <sup>+</sup> T=5/2			
<sup>31</sup> Ar	11330#	200#			15.0 ms 0.3	5/2 <sup>+</sup> *	13 14Ko17	T 1986	β <sup>+</sup> =100; β <sup>+</sup> p=68.3 3; β <sup>+</sup> 2p=9.0 2; β <sup>+</sup> pα<0.38; β <sup>+</sup> 3p=0.07 2; β <sup>+</sup> α<0.03; 2p<0.0006
<sup>31</sup> K	34260#	300#			> 10 ps	3/2 <sup>+</sup> #	19Ko18	IT 2019	3p=100
* <sup>31</sup> Na	D : %β <sup>-</sup> 2n average 19Ni04=0.7(0.1) and 0.86(0.20) from Pn=37.5(3.5), average								**
* <sup>31</sup> Na	D : of 74Ro31=30(8) 84La03=38(6) 19Ni04=40(5), and P2n/Pn=0.023(0.005)								**
* <sup>31</sup> Na	D : in 80De26. %β <sup>-</sup> n from Pn-2*P2n=37.5(3.5)-2*0.73(0.09). P3n from 84Gu19								**
* <sup>31</sup> Na	T : average 17Ha23=16.6(0.4) 84La03=17.0(0.4) 74Ro31=16.9(0.7) 01Pe14=18(2)								**
* <sup>31</sup> Na	J : 00Ke09=3/2								**
* <sup>31</sup> Mg	D : %β <sup>-</sup> n strongly conflicting with earlier 84La03=1.7(0.3)								**
* <sup>31</sup> Al	J : 20He.A, 16He09=5/2								**
* <sup>31</sup> Si	T : other 89Ab05=157.474(0.012), the small uncertainty is not justified								**
* <sup>31</sup> Ar	T : average 14Ko17=15.1(0.3) 00Fy01=14.1(0.7) 92Ba01=15.1(+1.3-1.1)								**
<sup>32</sup> Ne	37000#	500#			3.5 ms 0.9	0 <sup>+</sup>	11		β <sup>-</sup> =100; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>32</sup> Na	18640	40			12.9 ms 0.3	(3 <sup>-</sup> )	11 08Tr04	TJ 1972	β <sup>-</sup> =100; β <sup>-</sup> n=26 6; β <sup>-</sup> 2n=7.6 15
<sup>32</sup> Mg	-829	3			80.4 ms 0.4	0 <sup>+</sup>	11 17Ha23	T 1977	β <sup>-</sup> =100; β <sup>-</sup> n=5.5 5
<sup>32</sup> Al	-11099	7			32.6 ms 0.5	1 <sup>+</sup> *	11 17Ha23	T 1971	β <sup>-</sup> =100; β <sup>-</sup> n=0.7 5
<sup>32</sup> Al <sup>m</sup>	-10142	7	956.6	0.5	200 ns 20	(4 <sup>+</sup> )	11		IT=100
<sup>32</sup> Si	-24077.69	0.30			157 y 7	0 <sup>+</sup>	20		β <sup>-</sup> =100
<sup>32</sup> P	-24304.88	0.04			14.269 d 0.007	1 <sup>+</sup> *	11 FKG204	T 1934	β <sup>-</sup> =100
<sup>32</sup> P <sup>i</sup>	-19232.44	0.07	5072.44	0.06		0 <sup>+</sup> T=2	11		IT=100
<sup>32</sup> S	-26015.5371	0.0013			STABLE	0 <sup>+</sup>	11		IS=94.85 255
<sup>32</sup> S <sup>i</sup>	-19014.1	0.4	7001.4	0.4		1 <sup>+</sup> T=1	11		IT=100
<sup>32</sup> S <sup>j</sup>	-13967.58	0.28	12047.96	0.28		0 <sup>+</sup> T=2	11		IT=100
<sup>32</sup> Cl	-13334.7	0.6			298 ms 1	1 <sup>+</sup>	11		β <sup>+</sup> =100; β <sup>+</sup> α=0.054 8; β <sup>+</sup> p=0.026 5
<sup>32</sup> Cl <sup>i</sup>	-8288.4	0.7	5046.3	0.3		0 <sup>+</sup> T=2	11		IT=100
<sup>32</sup> Ar	-2200.4	1.8			98 ms 2	0 <sup>+</sup>	11		β <sup>+</sup> =100; β <sup>+</sup> p=35.58 22
<sup>32</sup> K	21990#	400#				1 <sup>+</sup> #			p ?
<sup>32</sup> K <sup>m</sup>	22940#	410#	950#	100#		4 <sup>+</sup> #	Mirror	I	p ?

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>32</sup> Na	T : average 08Tr04=13.1(0.5) and 11.5(1.2) 84La03=13.2(0.4) 98No.A=11.5(0.8)								**
* <sup>32</sup> Na	D : % $\beta^-$ n average 84Gu19=32(13) 84La03=24(7), from Pn=39(6) and								**
* <sup>32</sup> Na	D : % $\beta^-$ 2n=7.6(1.5); other 80De26=10(4); % $\beta^-$ 2n average 84Gu19=8(3)								**
* <sup>32</sup> Na	D : 84La03=9.4(2.4) 80De26=5.5(2.5)								**
* <sup>32</sup> Al	T : average 17Ha23=31.7(0.3) 05Ue01=33.0(0.2); Birge ratio=3.6								**
* <sup>32</sup> Al	J : 18Xu05,20He.A=1								**
<sup>33</sup> Ne	46130#	600#			<260ns	7/2 <sup>-</sup> #	11 02No11 I		n ? *
<sup>33</sup> Na	23780	450			8.2 ms 0.4	(3/2 <sup>+</sup> )	11	1972	$\beta^-$ =100; $\beta^-$ n=47 6; $\beta^-$ 2n=13 3
<sup>33</sup> Mg	4962.9	2.7			92.0 ms 1.2	3/2 <sup>-</sup> *	11 17Ha23 T	1979	$\beta^-$ =100; $\beta^-$ n=14 2; $\beta^-$ 2n ? *
<sup>33</sup> Al	-8497	7			41.46 ms 0.09	5/2 <sup>+</sup> *	11 16He09 J	1971	$\beta^-$ =100; $\beta^-$ n=8.5 7 *
<sup>33</sup> Si	-20514.3	0.7			6.18 s 0.18	3/2 <sup>+</sup> *	11	1971	$\beta^-$ =100
<sup>33</sup> P	-26337.4	1.1			25.35 d 0.11	1/2 <sup>+</sup>	20	1951	$\beta^-$ =100
<sup>33</sup> S	-26585.8583	0.0013			STABLE	3/2 <sup>+</sup>	11	1926	IS=0.763 20
<sup>33</sup> Si	-21106.07	0.13	5479.79	0.13		1/2 <sup>+</sup> T=3/2	11		IT=100
<sup>33</sup> Cl	-21003.3	0.4			2.5038 s 0.0022	3/2 <sup>+</sup>	11 15Gr14 T	1940	$\beta^+$ =100
<sup>33</sup> Cl <sup>i</sup>	-15454.9	0.5	5548.4	0.4	RQ	1/2 <sup>+</sup> T=3/2	11		IT=100
<sup>33</sup> Ar	-9384.3	0.4			173.0 ms 2.0	1/2 <sup>+</sup>	11 10Ad03 D	1964	$\beta^+$ =100; $\beta^+$ p=38.7 8 *
<sup>33</sup> K	7540#	200#			<25ns	3/2 <sup>+</sup> #	11 93Po.A I		p ?
<sup>33</sup> Ca	31030#	400#				5/2 <sup>+</sup> #			p ?
* <sup>33</sup> Ne	T : estimated partial $\beta^-$ decay half-life of 1# ms								**
* <sup>33</sup> Ne	I : also 02Le.A < 1.5 us								**
* <sup>33</sup> Mg	T : average 17Ha23=93.9(1.8) 02Mo29=90.5(1.6); other 84La03=90(20)								**
* <sup>33</sup> Mg	J : also 19Yo06=3/2								**
* <sup>33</sup> Al	T : average 17Ha23=41.4(0.1) 02Mo29=41.7(0.2); also 95Re.A=40.5(2.8)								**
* <sup>33</sup> Ar	D : % $\beta^+$ p average 10Ad03=38.8(1.3) 87Bo21=38.7(1.0)								**
<sup>34</sup> Ne	52840#	510#			2# ms >1.5us	0 <sup>+</sup>	12 02Le.A I	2002	$\beta^-$ ?; $\beta^-$ 2n ?; $\beta^-$ n ?
<sup>34</sup> Na	31680	600			5.5 ms 1.0	1 <sup>+</sup>	12 GAu03 D	1983	$\beta^-$ =100; $\beta^-$ 2n $\approx$ 50; $\beta^-$ n $\approx$ 15 *
<sup>34</sup> Mg	8323	7			44.9 ms 0.4	0 <sup>+</sup>	12 17Li03 TD	1979	$\beta^-$ =100; $\beta^-$ n=21 7; $\beta^-$ 2n<0.1
<sup>34</sup> Al	-2997.6	2.1			53.73 ms 0.13	4 <sup>-</sup>	12 19Li41 T	1977	$\beta^-$ =100; $\beta^-$ n=26 4; $\beta^-$ 2n ?
<sup>34</sup> Al <sup>m</sup>	-2951.1	2.1	46.47	0.17	22.1 ms 0.2	1 <sup>+</sup> *	19Li41 TD	2012	$\beta^-$ $\approx$ 100; $\beta^-$ n=11 4; $\beta^-$ 2n ? *
<sup>34</sup> Si	-19991.7	0.8			2.77 s 0.20	0 <sup>+</sup>	12	1971	$\beta^-$ =100
<sup>34</sup> Si <sup>m</sup>	-15735.6	0.9	4256.1	0.4	< 210 ns	(3 <sup>-</sup> )	12	1989	IT=100
<sup>34</sup> P	-24548.7	0.8			12.43 s 0.10	1 <sup>+</sup>	12	1945	$\beta^-$ =100
<sup>34</sup> S	-29931.69	0.04			STABLE	0 <sup>+</sup>	12	1926	IS=4.365 235
<sup>34</sup> Cl	-24440.09	0.05			1.5267 s 0.0004	0 <sup>+</sup> T=1	12 06Ia05 T	1934	$\beta^+$ =100 *
<sup>34</sup> Cl <sup>m</sup>	-24293.73	0.05	146.360	0.027	MD	3 <sup>+</sup> T=0	12	1965	$\beta^+$ =55.4 6;IT=44.6 6
<sup>34</sup> Ar	-18378.29	0.08			846.46 ms 0.35	0 <sup>+</sup>	12 20Ia01 T	1966	$\beta^+$ =100 *
<sup>34</sup> Ar <sup>i</sup>	-10444	5	7934	5	RQ	1 <sup>+</sup> # T=2	12	1969	IT ?
<sup>34</sup> K	-1220#	200#			<40ns	1 <sup>+</sup> #	12 93Po.A I		p ?
<sup>34</sup> Ca	14890#	300#			<35ns	0 <sup>+</sup>	12 93Po.A I		2p ?
* <sup>34</sup> Na	D : % $\beta^-$ n, % $\beta^-$ 2n estimated from Pn = $\beta^-$ n + 2* $\beta^-$ 2n = 115(20)% in 84La03								**
* <sup>34</sup> Na	D : by assuming $\beta^-$ n/ $\beta^-$ 2n=0.3 from trends in neighboring nuclei								**
* <sup>34</sup> Al <sup>m</sup>	E : from FGK204 using a least-squares fit to data in 17Li03								**
* <sup>34</sup> Al <sup>m</sup>	J : 18Xu05=1								**
* <sup>34</sup> Cl	T : average 06Ia05=1.5268(5) 83Ko22=1.5277(22) 76Wi08=1.5252(11)								**
* <sup>34</sup> Cl	T : 73Ry01=1.526(2) 72Ha82=1.534(3)								**
* <sup>34</sup> Ar	T : others 06Ia05=843.8(0.4) 74Ha26=844.5(3.4)								**
<sup>35</sup> Na	37830#	670#			1.5 ms 0.5	3/2 <sup>+</sup> #	11	1983	$\beta^-$ =100; $\beta^-$ n ?; $\beta^-$ 2n ? *
<sup>35</sup> Mg	15640	270			11.3 ms 0.6	(3/2 <sup>-</sup> , 5/2 <sup>-</sup> )	17 17Mo26 J	1989	$\beta^-$ =100; $\beta^-$ n=52 46; $\beta^-$ 2n ?
<sup>35</sup> Al	-224	7			38.16 ms 0.21	(5/2 <sup>+</sup> , 3/2 <sup>+</sup> )	11 17Ch36 J	1979	$\beta^-$ =100; $\beta^-$ n=35.8 17; $\beta^-$ 2n ? *
<sup>35</sup> Si	-14390	40			780 ms 120	7/2 <sup>-</sup> #	15 95Re.A D	1971	$\beta^-$ =100; $\beta^-$ n<5
<sup>35</sup> P	-24857.8	1.9			47.3 s 0.8	1/2 <sup>+</sup>	11	1971	$\beta^-$ =100
<sup>35</sup> S	-28846.21	0.04			87.37 d 0.04	3/2 <sup>+</sup> *	11	1936	$\beta^-$ =100
<sup>35</sup> S <sup>i</sup>	-19691	10	9155	10	RQ	T=5/2	(1/2 <sup>-</sup> : 9/2 <sup>+</sup> )	11	1975
<sup>35</sup> Cl	-29013.53	0.04			STABLE	3/2 <sup>+</sup> *	11	1919	IS=75.8 2

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{35}\text{Cl}^i$	-23359.05	0.22	5654.48	0.22		$3/2^+$ T=3/2	11		IT=100
$^{35}\text{Ar}$	-23047.3	0.7			1.7756 s 0.0010	$3/2^+$	11	1940	$\beta^+$ =100
$^{35}\text{Ar}^i$	-17474.6	0.7	5572.66	0.15		$3/2^+$ T=3/2	11		IT=100
$^{35}\text{K}$	-11172.9	0.5			175.2 ms 1.9	$3/2^+$	11 06Me04 J	1976	$\beta^+$ =100; $\beta^+$ p=0.37 15
$^{35}\text{K}^i$	-2110	40	9060	40	2p	$3/2^+$ T=5/2			
$^{35}\text{Ca}$	5190#	200#			25.7 ms 0.2	$1/2^+$ #	11 99Tr04 DT	1985	$\beta^+$ =100; $\beta^+$ p=95.8 14; $\beta^+$ 2p=4.2 3
$^{35}\text{Sc}$	27100#	400#				$7/2^-$ #			p ?
* $^{35}\text{Na}$	D : $\beta^-$ n was observed by 83La12, but it was not quantified								
* $^{35}\text{Al}$	T : average 17Ha23=38.4(0.3) 05Ti11=36.8(0.5) 01Nu01=38.6(0.4); others								
* $^{35}\text{Al}$	T : 95Re.A=30(4) 89Le16=170(+90-50) 88Mu08=130(+100-50)								
* $^{35}\text{Al}$	D : % $\beta^-$ n average 05Ti11=38(2) 01Nu01=41(13) 95Re.A=26(4) 89Le16=40(10);								
* $^{35}\text{Al}$	D : other 88Mu08=87(+37-25)								
* $^{35}\text{K}$	T : average 18Sa.A=175(2) 98Sc19=178(8)								
$^{36}\text{Na}$	45900#	690#			<180ns		12 02Le.A I		n ?
$^{36}\text{Mg}$	20380	690			3.9 ms 1.3	$0^+$	12	1989	$\beta^-$ =100; $\beta^-$ n=48 12; $\beta^-$ 2n ?
$^{36}\text{Al}$	5950	150			90 ms 40		12	1979	$\beta^-$ =100; $\beta^-$ n<31; $\beta^-$ 2n ?
$^{36}\text{Si}$	-12440	70			503 ms 2	$0^+$	12 95Re.A D	1971	$\beta^-$ =100; $\beta^-$ n=12 5
$^{36}\text{P}$	-20251	13			5.6 s 0.3	$4^-$	12 15Ch56 J	1971	$\beta^-$ =100; $\beta^-$ n ?
$^{36}\text{S}$	-30664.14	0.19			STABLE	$0^+$	12	1938	IS=0.0158 17
$^{36}\text{Cl}$	-29522.01	0.04			301.3 ky 1.5	$2^+*$	12	1941	$\beta^-$ =98.1 1; $\beta^+$ =1.9 1
$^{36}\text{Cl}^i$	-25222.34	0.04	4299.667	0.014		$(0)^+$ T=2	12		IT=100
$^{36}\text{Ar}$	-30231.542	0.027			STABLE	$0^+$	12	1920	IS=0.3336 210;2 $\beta^+$ ?
$^{36}\text{Ar}^i$	-23620.5	0.3	6611.0	0.3		$2^+$ T=1	12		IT=100
$^{36}\text{Ar}^j$	-19379.4	1.2	10852.2	1.2	RQ	$0^+$ T=2	12		IT=100
$^{36}\text{K}$	-17417.2	0.3			341 ms 3	$2^+*$	12	1967	$\beta^+$ =100; $\beta^+$ p=0.048 14; $\beta^+$ $\alpha$ =0.0034 13
$^{36}\text{K}^i$	-13134.5	2.4	4282.7	2.4	p	$0^+$ T=2	12		p=100
$^{36}\text{Ca}$	-6450	40			100.9 ms 1.3	$0^+$	12 15Su01 T	1977	$\beta^+$ =100; $\beta^+$ p=51.2 10
$^{36}\text{Sc}$	16150#	300#							p ?
* $^{36}\text{Mg}$	D : % $\beta^-$ n from 99YoZW								
* $^{36}\text{Si}$	T : from 17Ha23								
* $^{36}\text{K}^i$	E : Ensdf2012 reports 4281.9(0.8) as IAS of $^{36}\text{Ca}$ gs, but the small								
* $^{36}\text{K}^j$	E : uncertainty is not justified								
* $^{36}\text{Ca}$	T : average 15Su01=100.0(2.4) 07Do17=100.1(2.3) 95Tr02,97Tr05=102(2)								
$^{37}\text{Na}$	53130#	690#			1# ms >1.5us	$3/2^+$ #	12 02Le.A I	2002	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?
$^{37}\text{Mg}$	28210	700			8 ms 4	$(3/2^-)$	12 14Ko14 J	1996	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?
$^{37}\text{Al}$	9810	180			11.4 ms 0.3	$5/2^+$ #	12 19Ab06 TD	1979	$\beta^-$ =100; $\beta^-$ n=52 5; $\beta^-$ 2n>1
$^{37}\text{Si}$	-6570	110			141.0 ms 3.5	$(5/2^-)$	15 19Ab06 TJ	1979	$\beta^-$ =100; $\beta^-$ n=17 13; $\beta^-$ 2n ?
$^{37}\text{P}$	-19000	40			2.31 s 0.13	$(1/2^+)$	12 15Ch56 J	1971	$\beta^-$ =100; $\beta^-$ n ?
$^{37}\text{S}$	-26896.43	0.20			5.05 m 0.02	$7/2^-$	12	1945	$\beta^-$ =100
$^{37}\text{Cl}$	-31761.55	0.05			STABLE	$3/2^+*$	12	1919	IS=24.2 2
$^{37}\text{Cl}^i$	-21539.7	0.3	10221.8	0.3	RQ	$7/2^-$ T=5/2	12	1984	IT=100
$^{37}\text{Ar}$	-30947.68	0.21			35.011 d 0.019	$3/2^+*$	12	1941	$\epsilon$ =100
$^{37}\text{Ar}^i$	-25956	6	4992	6	RQ	$3/2^+$ T=3/2	12	1973	
$^{37}\text{K}$	-24800.20	0.09			1.23651 s 0.00094	$3/2^+*$	12 14Sh25 T	1958	$\beta^+$ =100
$^{37}\text{K}^i$	-19749.9	0.8	5050.3	0.8	RQ	$3/2^+$ T=3/2	12	1973	IT=100
$^{37}\text{Ca}$	-13136.1	0.6			181.0 ms 0.9	$3/2^+*$	12 19Kl06 J	1964	$\beta^+$ =100; $\beta^+$ p=76.8 7
$^{37}\text{Sc}$	3780#	300#				$7/2^-$ #			p ?
$^{37}\text{Ti}$	25170#	400#				$1/2^+$ #			p ?
* $^{37}\text{Al}$	T : average 19Ab06=11.3(0.4) 15St14=11.5(0.4)								
* $^{37}\text{Al}$	D : % $\beta^-$ 2n 15St14>1%								
* $^{37}\text{Si}$	T : average 19Ab06=138(4) 17Ha23=150(7)								
* $^{37}\text{K}$	J : also 14Kr04=3/2								
* $^{37}\text{Ca}$	T : average 15Su01=180.5(2.1) 07Do17=181.7(3.6) 97Tr05=181.1(1.0)								
* $^{37}\text{Ca}$	D : % $\beta^+$ p average 15Su01=70.6(1.8) 07Do17=72.2(4.3) 97Tr05=78.4(0.8)								
* $^{37}\text{Ca}$	D : $^{74}\text{Se}11=75(2)$ . Ensdf2012 gives $^{97}\text{Tr}05=82.1(0.7)$ , which								
* $^{37}\text{Ca}$	D : differs from $^{97}\text{Tr}05=78.4(0.8)$ quoted in the present evaluation								

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)		
<sup>38</sup> Na	61910#	720#	<400 ns		19Ah07	I	n ?	*	
<sup>38</sup> Mg	34070#	500#	2# ms >260ns	0 <sup>+</sup>	17	1997	$\beta^- = 100\#; \beta^- n ?; \beta^- 2n ?$	*	
<sup>38</sup> Al	16470#	150#	9.0 ms 0.7	0 <sup>-</sup> #	17 15St14	T	1989	$\beta^- = 100; \beta^- n = 84 19; \beta^- 2n ?$	*
<sup>38</sup> Si	-4170	100	63 ms 8	0 <sup>+</sup>	17 17Tr02	TD	1979	$\beta^- = 100\#; \beta^- n = 25 10$	
<sup>38</sup> P	-14620	70	640 ms 140	(2 <sup>-</sup> )	17 15Ch56	J	1971	$\beta^- = 100; \beta^- n = 12 5$	
<sup>38</sup> S	-26861	7	170.3 m 0.7	0 <sup>+</sup>	17		1958	$\beta^- = 100$	
<sup>38</sup> Cl	-29798.12	0.10	37.230 m 0.014	2 <sup>-</sup>	17		1940	$\beta^- = 100$	
<sup>38</sup> Cl <sup>m</sup>	-29126.75	0.10	671.365 0.008	5 <sup>-</sup>	17		1954	IT=100	
<sup>38</sup> Cl <sup>i</sup>	-21590	24	8208 24 RQ	0 <sup>+</sup> T=3	17				
<sup>38</sup> Ar	-34714.83	0.19	STABLE	0 <sup>+</sup>	17		1934	IS=0.0629 70	
<sup>38</sup> Ar <sup>i</sup>	-24083.9	0.9	10630.9 0.9	(2 <sup>-</sup> ) T=2	17				
<sup>38</sup> Ar <sup>j</sup>	-15940	30	18780 30 RQ	0 <sup>+</sup> T=3	17				
<sup>38</sup> K	-28800.76	0.20	7.651 m 0.019	3 <sup>+</sup> * T=0	17		1937	$\beta^+ = 100$	*
<sup>38</sup> K <sup>m</sup>	-28670.61	0.20	130.15 0.04 MD	0 <sup>+</sup> * T=1	17 14Pa45	J	1953	$\beta^+ = 99.9670 43;$ IT=0.0330 43	*
<sup>38</sup> K <sup>n</sup>	-25342.66	0.26	3458.10 0.17	(7 <sup>+</sup> )	17		1971	IT=100	
<sup>38</sup> Ca	-22058.50	0.19	443.70 ms 0.25	0 <sup>+</sup>	17 15B102	T	1966	$\beta^+ = 100$	*
<sup>38</sup> Sc	-4250#	200#	<300ns	2 <sup>-</sup> #	17 94B110	I		p ?	
<sup>38</sup> Sc <sup>m</sup>	-3580#	220#	670# 100#	5 <sup>-</sup> #	Mirror	I		IT ?; p ?	
<sup>38</sup> Ti	11370#	300#	<120ns	0 <sup>+</sup>	17 96B121	I		2p ?	
* <sup>38</sup> Na	I : no events observed in 19Ah07							**	
* <sup>38</sup> Al	T : other 04Gr20=7.6(0.6) without $\beta$ - $\gamma$ coin gating							**	
* <sup>38</sup> K	J : 19Ko19,14Kr04,14Pa45=3							**	
* <sup>38</sup> K <sup>m</sup>	T : average 10Ba43=924.46(0.14) 00Bb01=924.4(0.6) 83Ko22=924.15(0.31)							**	
* <sup>38</sup> K <sup>n</sup>	T : 78Th02=928.8(2.0) 78Wi04=921.71(0.65) 76Wi08=922.3(1.1)							**	
* <sup>38</sup> K <sup>m</sup>	T : 72Ha82=929.2(3.5) 75Sq01=925.6(0.7)							**	
* <sup>38</sup> Ca	T : average of 15B102=443.63(0.35) 11Pa38=443.77(0.36) 10B109=443.8(1.9)							**	
<sup>39</sup> Na	69980#	740#	1# ms >400ns	3/2 <sup>+</sup> #	19Ah07	I	2019	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$	*
<sup>39</sup> Mg	42780#	510#	<180ns	7/2 <sup>-</sup> #	18			n ?; $\beta^- ?$	*
<sup>39</sup> Al	21490#	300#	7.6 ms 1.6	5/2 <sup>+</sup> #	18		1989	$\beta^- = 100; \beta^- n = 97 22; \beta^- 2n ?$	*
<sup>39</sup> Si	2320	140	41.2 ms 4.1	(5/2 <sup>-</sup> )	18 19Ab06	TDJ	1979	$\beta^- = 100; \beta^- n = 33 3; \beta^- 2n ?$	*
<sup>39</sup> P	-12770	110	282 ms 24	(1/2 <sup>+</sup> )	18 19Ab06	J	1977	$\beta^- = 100; \beta^- n = 26 8$	*
<sup>39</sup> S	-23160	50	11.5 s 0.5	(7/2 <sup>-</sup> )	18		1971	$\beta^- = 100$	
<sup>39</sup> Cl	-29800.2	1.7	56.2 m 0.6	3/2 <sup>+</sup>	18		1949	$\beta^- = 100$	
<sup>39</sup> Ar	-33242	5	268 y 8	7/2 <sup>-</sup> *	18		1950	$\beta^- = 100$	
<sup>39</sup> Ar <sup>i</sup>	-24161	7	9081 9 RQ	3/2 <sup>+</sup> T=5/2	18 MMC149J				*
<sup>39</sup> K	-33807.195	0.005	STABLE	3/2 <sup>+</sup> *	18		1921	IS=93.2581 44	*
<sup>39</sup> K <sup>i</sup>	-27260.8	1.9	6546.4 1.9	7/2 <sup>-</sup> T=3/2	18			IT=100	
<sup>39</sup> Ca	-27282.7	0.6	860.3 ms 0.8	3/2 <sup>+</sup> *	18 19K106	J	1943	$\beta^+ = 100$	*
<sup>39</sup> Ca <sup>i</sup>	-20917#	9#	6366# 9#	3/2 <sup>+</sup> T=3/2	Imme	E			
<sup>39</sup> Sc	-14173	24		7/2 <sup>-</sup> #	18 94B110	I	1988	p=100	*
<sup>39</sup> Sc <sup>i</sup>	-5050	40	9120 50 2p	(3/2 <sup>+</sup> ) T=5/2	18				
<sup>39</sup> Ti	2500#	200#	28.5 ms 0.9	3/2 <sup>+</sup> #	18 07Do17	TD	1990	$\beta^+ = 100; \beta^+ p = 93.7 28;$ $\beta^+ 2p = ?$ p ?	*
<sup>39</sup> V	22570#	400#		7/2 <sup>-</sup> #					
* <sup>39</sup> Na	I : one event observed in 19Ah07							**	
* <sup>39</sup> Mg	T : estimated partial $\beta^-$ half-life of 1# ms							**	
* <sup>39</sup> Si	T : average 19Ab06=38.6(1.3) 04Gr20=47.5(2.0); Birge ratio=3.7							**	
* <sup>39</sup> P	T : average 04Gr20=250(80) 98Wi.A=320(30) 95Re.A=190(50)							**	
* <sup>39</sup> Ar <sup>i</sup>	J : from IAS appartenance; 3/2+,5/2+ in Ensd12018							**	
* <sup>39</sup> K	J : 19Ko19,14Kr04,14Pa45,13Pa11=3/2							**	
* <sup>39</sup> Ca	T : average 10B109=860.7(1.0) 77Az01=859.4(1.6) 73Al11=860.4(3.0)							**	
* <sup>39</sup> Sc	D : most likely a proton emitter from Sp=-597(24) keV							**	
* <sup>39</sup> Ti	D : % $\beta^+ p$ includes contribution from <sup>39</sup> Sc gs (%p=100); $\beta^+ 2p$ decay was							**	
* <sup>39</sup> Ti	D : observed in 92Mo15, but was not quantified							**	
<sup>40</sup> Mg	49550#	500#	1# ms >170ns	0 <sup>+</sup>	17 14Cr02	I	2007	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$	*
<sup>40</sup> Al	28820#	300#	10# ms >260ns		17		1996	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$	*
<sup>40</sup> Si	5670	120	31.2 ms 2.6	0 <sup>+</sup>	17 17Tr02	TD	1989	$\beta^- = 100; \beta^- n = 38 5; \beta^- 2n ?$	*



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{40}\text{P}$	-8140	80			150 ms 8	$(2^-, 3^-)$	17	1979	$\beta^- = 100; \beta^- n = 15.8 \text{ 21};$ $\beta^- 2n ?$
$^{40}\text{S}$	-22838	4			8.8 s 2.2	$0^+$	17	1971	$\beta^- = 100$
$^{40}\text{Cl}$	-27560	30			1.35 m 0.03	$2^-$	17	1956	$\beta^- = 100$
$^{40}\text{Ar}$	-35039.9000	0.0022			STABLE	$0^+$	17	1920	IS=99.6035 250
$^{40}\text{K}$	-33535.50	0.06			1.248 Gy 0.003	$4^- *$	17	1935	IS=0.0117 1; $\beta^- = 89.28 \text{ 13};$ $\beta^+ = 10.72 \text{ 13}$ *
$^{40}\text{K}^m$	-31891.86	0.06	1643.638	0.011	336 ns 12	$0^+$	17	1968	IT=100
$^{40}\text{K}^i$	-29151.5	0.3	4384.0	0.3		$0^+ \text{ T}=2$	17		IT=100
$^{40}\text{Ca}$	-34846.402	0.020			STABLE >9.9Zy	$0^+$	17 16An14 T	1922	IS=96.941 156; $2\beta^+ ?$
$^{40}\text{Ca}^i$	-27188.22	0.05	7658.18	0.05		$4^- \text{ T}=1$	17 AHW E		IT=100 *
$^{40}\text{Ca}^j$	-22858.4	1.0	11988	1		$0^+ \text{ T}=2$	17		IT=100
$^{40}\text{Sc}$	-20523.4	2.8			182.3 ms 0.7	$4^-$	17	1955	$\beta^+ = 100; \beta^+ p = 0.44 \text{ 7};$ $\beta^+ \alpha = 0.017 \text{ 5}$
$^{40}\text{Sc}^i$	-16164	6	4359	6 RQ		$0^+ \text{ T}=2$	17		IT=100
$^{40}\text{Ti}$	-8990	70			52.4 ms 0.3	$0^+$	17 07Do17 TD	1982	$\beta^+ = 100; \beta^+ p = 95.8 \text{ 13}$
$^{40}\text{V}$	12470#	300#				$2^- \#$			p ?
* $^{40}\text{Mg}$	I : 5 events observed in direct two-proton removal from $^{42}\text{Si}$ **								
* $^{40}\text{Si}$	T : average 17Tr02=27.6(1.4) 04Gr20=33(1); Birge ratio=3.14 **								
* $^{40}\text{K}$	J : also 14Kr04j=4 **								
* $^{40}\text{Ca}^i$	E : originally 7658.23(0.05), recalibrated -0.05 keV for $^{27}\text{Al}+p$ **								
* $^{40}\text{Ca}^i$	E : resonances **								
$^{41}\text{Mg}$	58100#	500#				$3/2^- \#$			$\beta^- ?; \beta^- n ?$
$^{41}\text{Al}$	34590#	400#			6# ms >260ns	$5/2^+ \#$	16 97Sa14 I	1997	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$
$^{41}\text{Si}$	13200#	300#			20.0 ms 2.5	$7/2^- \#$	16	1989	$\beta^- = 100; \beta^- n > 55; \beta^- 2n ?$ *
$^{41}\text{P}$	-4980	120			101 ms 5	$1/2^+ \#$	16	1979	$\beta^- = 100; \beta^- n = 30 \text{ 10}; \beta^- 2n ?$
$^{41}\text{S}$	-19009	4			1.99 s 0.05	$7/2^- \#$	16	1979	$\beta^- = 100; \beta^- n ?$
$^{41}\text{Cl}$	-27310	70			38.4 s 0.8	$(1/2^+)$	16	1971	$\beta^- = 100$
$^{41}\text{Ar}$	-33067.5	0.3			109.61 m 0.04	$7/2^-$	16	1936	$\beta^- = 100$
$^{41}\text{K}$	-35559.549	0.004			STABLE	$3/2^+ *$	16	1921	IS=6.7302 44 *
$^{41}\text{K}^i$	-27210	15	8349	15 RQ		$7/2^- \text{ T}=5/2$	16 75Me10 J	1975	
$^{41}\text{Ca}$	-35137.91	0.14			99.4 ky 1.5	$7/2^- *$	16	1939	$\epsilon = 100$
$^{41}\text{Ca}^i$	-29320.8	0.5	5817.1	0.5	< 28 fs	$3/2^+ \text{ T}=3/2$	16		IT=100
$^{41}\text{Sc}$	-28642.36	0.08			596.3 ms 1.7	$7/2^-$	16	1941	$\beta^+ = 100$
$^{41}\text{Sc}^r$	-25760.01	0.08	2882.35	0.05 RQ		$7/2^+$	16		p=59 2; IT=41 2
$^{41}\text{Sc}^i$	-22704	3	5939	3 RQ		$3/2^+ \text{ T}=3/2$	16		p=100
$^{41}\text{Ti}$	-15698	28			81.9 ms 0.5	$3/2^+$	16 07Do17 D	1964	$\beta^+ = 100; \beta^+ p = 91.1 \text{ 6}$
$^{41}\text{V}$	310#	200#				$7/2^- \#$			p ?
$^{41}\text{Cr}$	20410#	400#				$3/2^+ \#$			p ?
* $^{41}\text{Si}$	D : % $\beta^- n$ from Pn=103(38) in 99YoZW **								
* $^{41}\text{K}$	J : also 14Kr04j=3/2 **								
* $^{41}\text{K}^i$	J : l=3 in ( $^3\text{He}, d$ ); assigned as IAS of $^{41}\text{Ar}$ gs **								
$^{42}\text{Al}$	41990#	500#			3# ms >170ns		16	2007	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$
$^{42}\text{Si}$	16840#	300#			12.5 ms 3.5	$0^+$	16	1990	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$
$^{42}\text{P}$	1090	100			48.5 ms 1.5		16	1979	$\beta^- = 100; \beta^- n = 50 \text{ 20}; \beta^- 2n ?$
$^{42}\text{S}$	-17637.7	2.8			1.016 s 0.015	$0^+$	16 06Wi10 D	1979	$\beta^- = 100; \beta^- n < 1$
$^{42}\text{Cl}$	-24830	60			6.8 s 0.3	$(2^-)$	16	1971	$\beta^- = 100; \beta^- n ?$
$^{42}\text{Ar}$	-34423	6			32.9 y 1.1	$0^+$	16	1952	$\beta^- = 100$
$^{42}\text{K}$	-35022.03	0.11			12.355 h 0.007	$2^- *$	16	1935	$\beta^- = 100$ *
$^{42}\text{K}^i$	-28570	100	6450	100		$(0^+) \text{ T}=3$	16		
$^{42}\text{Ca}$	-38547.29	0.15			STABLE	$0^+$	16	1934	IS=0.647 23
$^{42}\text{Ca}^i$	-28797	10	9750	10		$(2^-) \text{ T}=2$	16		
$^{42}\text{Sc}$	-32121.00	0.15			680.72 ms 0.26	$0^+ \text{ T}=1$	16 97Ko65 T	1955	$\beta^+ = 100$ *
$^{42}\text{Sc}^m$	-31504.19	0.16	616.81	0.06 MD	61.7 s 0.4	$7^+$	16	1963	$\beta^+ = 100$
$^{42}\text{Sc}^r$	-26044.80	0.16	6076.20	0.07 RQ		$(2^+, 3^+, 4^+)$	16		IT=100
$^{42}\text{Ti}$	-25104.35	0.27			208.3 ms 0.4	$0^+$	16 15Mo01 T	1964	$\beta^+ = 100$ *
$^{42}\text{V}$	-7620#	200#			<55ns	$2^- \#$	16 92Bo37 I		p ?
$^{42}\text{Cr}$	7060#	300#			13.3 ms 1.0	$0^+$	16	1996	$\beta^+ \approx 100; \beta^+ p = 94.4 \text{ 50}; 2p ?$
* $^{42}\text{K}$	J : 19Ko19, 14Kr04, 14Pa45=2 **								

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>42</sup> Sc	T : average 97Ko65=680.67(0.28) 76Wi08=680.98(0.62)								**
* <sup>42</sup> Ti	T : average 15Mo01=211.7(1.9), 209.5(5.2) 09Ku19=208.14(0.45)								**
<sup>43</sup> Al	48270#	600#			4# ms >170ns	5/2 <sup>+</sup> #	15	2007	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
<sup>43</sup> Si	24330#	400#			30# ms >260ns	3/2 <sup>-</sup> #	15 02No11 I	2002	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
<sup>43</sup> P	5040#	300#			35.8 ms 1.3	(1/2 <sup>+</sup> )	15 04Gr20 T	1989	$\beta^-$ =100; $\beta^-n$ =100; $\beta^-2n$ ?
<sup>43</sup> S	-12195	5			265 ms 13	3/2 <sup>-</sup>	15 20Mo32 J	1979	$\beta^-$ =100; $\beta^-n$ =40 10
<sup>43</sup> S <sup>m</sup>	-11874	5	320.7	0.5	415.0 ns 2.6	(7/2 <sup>-</sup> )	15 09Ga05 J	2000	IT=100
<sup>43</sup> Cl	-24160	60			3.13 s 0.09	(3/2 <sup>+</sup> )	15 06Wi10 JT	1976	$\beta^-$ =100; $\beta^-n$ ?
<sup>43</sup> Ar	-32010	5			5.37 m 0.06	5/2 <sup>(-)</sup>	15	1969	$\beta^-$ =100
<sup>43</sup> K	-36575.4	0.4			22.3 h 0.1	3/2 <sup>+</sup> *	15 14Kr04 J	1949	$\beta^-$ =100
<sup>43</sup> K <sup>m</sup>	-35837.1	0.4	738.30	0.06	200 ns 5	7/2 <sup>-</sup>	15	1978	IT=100
<sup>43</sup> Ca	-38408.87	0.23			STABLE	7/2 <sup>-</sup> *	15	1934	IS=0.135 10
<sup>43</sup> Ca <sup>i</sup>	-30414	14	7995	14	RQ	(3/2 <sup>+</sup> ) <sup>+</sup> T=5/2	15		
<sup>43</sup> Sc	-36188.1	1.9			3.891 h 0.012	7/2 <sup>-</sup> *	15	1935	$\beta^+$ =100
<sup>43</sup> Sc <sup>m</sup>	-36036.3	1.9	151.79	0.08	438 $\mu$ s 5	3/2 <sup>+</sup>	15 77Mi10 T	1964	IT=100
<sup>43</sup> Sc <sup>n</sup>	-33064.4	1.9	3123.73	0.15	472 ns 3	19/2 <sup>-</sup>	15 08Fe02 T	1978	IT=100
<sup>43</sup> Sc <sup>i</sup>	-31956	3	4232	4	RQ	7/2 <sup>-</sup> T=3/2	15		
<sup>43</sup> Ti	-29316	6			509 ms 5	7/2 <sup>-</sup>	15	1948	$\beta^+$ =100; $\beta^+p$ ?
<sup>43</sup> Ti <sup>m</sup>	-29003	6	313.0	1.0	11.9 $\mu$ s 0.3	(3/2 <sup>+</sup> )	15	1978	IT=100
<sup>43</sup> Ti <sup>n</sup>	-26250	6	3066.4	1.0	556 ns 6	(19/2 <sup>-</sup> )	15	1978	IT=100
<sup>43</sup> Ti <sup>i</sup>	-24610#	50#	4710#	50#		7/2 <sup>-</sup> # T=3/2			
<sup>43</sup> V	-17920	40			79.3 ms 2.4	7/2 <sup>-</sup> #	15 07Do17 D	1987	$\beta^+$ =100; $\beta^+p$ <2.5
<sup>43</sup> Vi	-9705	15	8210	50	RQ	3/2 <sup>+</sup> T=5/2			
<sup>43</sup> Cr	-1970#	200#			21.1 ms 0.3	(3/2 <sup>+</sup> )	15 11Po01 T	1992	$\beta^+$ =100; $\beta^+p$ =79.3 30; $\beta^+2p$ =11.6 10; $\beta^+3p$ =0.13 +18-8; $\beta^+\alpha$ ?
<sup>43</sup> Mn	17370#	400#				5/2 <sup>-</sup> #			p ?
* <sup>43</sup> P	T : average 04Gr20=36.5(1.5) 95So03=33(3)								**
* <sup>43</sup> S <sup>m</sup>	T : average 12Ch16=415(3) 09Ga05=415(5)								**
* <sup>43</sup> Cl	T: 06Wi10, supersedes 98WiZX=3.07(0.07); others 81Vo04=3.3(0.2)								**
* <sup>43</sup> Cl	T: 81HuZT=3.4(0.3)								**
* <sup>43</sup> Ca	J : also 15Ru02=7/2								**
* <sup>43</sup> Sc	J : also 11Av01=7/2								**
* <sup>43</sup> Sc <sup>m</sup>	T : average 77Mi10=438(7) 65De15=470(20) 64Ho14=435(7)								**
* <sup>43</sup> Sc <sup>n</sup>	T : average 08Fe02=481(9) 81Da06=469(4) 78Ha07=473(5)								**
* <sup>43</sup> Cr	T : average 11Po01=20.6(0.9) 07Do17=21.1(0.4) 01Gi01=21.6(0.7)								**
<sup>44</sup> Si	29310#	500#			4# ms >360ns	0 <sup>+</sup>	11	2007	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
<sup>44</sup> P	11110#	400#			18.5 ms 2.5		11	1989	$\beta^-$ =100; $\beta^-n$ ?; $\beta^-2n$ ?
<sup>44</sup> S	-9204	5			100 ms 1	0 <sup>+</sup>	11	1979	$\beta^-$ =100; $\beta^-n$ =18 3
<sup>44</sup> S <sup>m</sup>	-7839	5	1365.0	0.8	2.619 $\mu$ s 0.026	0 <sup>+</sup>	11	2005	IT=100
<sup>44</sup> Cl	-20480	90			562 ms 106	(2 <sup>-</sup> )	11	1979	$\beta^-$ =100; $\beta^-n$ <8
<sup>44</sup> Ar	-32673.3	1.6			11.87 m 0.05	0 <sup>+</sup>	11	1969	$\beta^-$ =100
<sup>44</sup> K	-35781.5	0.4			22.13 m 0.19	2 <sup>-</sup> *	11	1954	$\beta^-$ =100
<sup>44</sup> Ca	-41468.7	0.3			STABLE	0 <sup>+</sup>	11	1922	IS=2.086 110
<sup>44</sup> Ca <sup>i</sup>	-29619	10	11850	10		2 <sup>-</sup> T=3	11 MMC143J		*
<sup>44</sup> Sc	-37816.0	1.8			4.0421 h 0.0025	2 <sup>+</sup> *	11 16Ga24 T	1937	$\beta^+$ =100
<sup>44</sup> Sc <sup>m</sup>	-37748.1	1.8	67.8679	0.0014	154.8 ns 0.8	1 <sup>-</sup>	11	1967	IT=100
<sup>44</sup> Sc <sup>n</sup>	-37669.8	1.8	146.1914	0.0020	51.0 $\mu$ s 0.3	0 <sup>-</sup>	11	1963	IT=100
<sup>44</sup> Sc <sup>p</sup>	-37544.8	1.8	271.240	0.010	58.61 h 0.10	6 <sup>+</sup> *	11	1940	IT=98.80 7; $\beta^+$ =1.20 7
<sup>44</sup> Sc <sup>i</sup>	-35038.3	2.5	2778	3	RQ	0 <sup>+</sup> T=2	11		
<sup>44</sup> Ti	-37548.6	0.7			59.1 y 0.3	0 <sup>+</sup>	11	1954	$\epsilon$ =100
<sup>44</sup> Ti <sup>i</sup>	-30942.2	0.9	6606.4	0.5		2 <sup>+</sup> T=1	11		IT=100
<sup>44</sup> Ti <sup>j</sup>	-28210.6	2.1	9338	2		0 <sup>+</sup> frg. T=2	11		IT=100
<sup>44</sup> V	-23808	7			111 ms 7	(2 <sup>+</sup> )	11	1971	$\beta^+$ =100; $\beta^+\alpha$ =?; $\beta^+p$ ?
<sup>44</sup> V <sup>m</sup>	-23537	5	271	9	MD	(6 <sup>+</sup> )	11	1997	$\beta^+$ =100
<sup>44</sup> V <sup>n</sup>	-23660#	100#	150#	100#		0 <sup>-</sup> #	Mirror I		
<sup>44</sup> Vi	-21119	12	2689	14	p	0 <sup>+</sup> # T=2	92Bo37 D	1992	p=100
<sup>44</sup> Cr	-13420	50			42.8 ms 0.6	0 <sup>+</sup>	11 07Do17 TD	1987	$\beta^+$ =100; $\beta^+p$ =12 2
<sup>44</sup> Mn	7460#	300#			<105ns	2 <sup>-</sup> #	11		p ?

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
* <sup>44</sup> Cl	T : average 99WiZX=650(50) 95So03=434(60); Birge ratio=2.77						**	
* <sup>44</sup> K	J : 19Ko19,14Kr04,14Pa45=2						**	
* <sup>44</sup> Ca <sup>i</sup>	J : from (e,e') in 84Ra04; IAS candidate						**	
* <sup>44</sup> Sc	T : average 16Ga24=4.0420(0.0025) 69Sa34=4.05(0.03). others (not used)						**	
* <sup>44</sup> Sc	T : 69Ra16=3.927(0.008) 66Ta01=4.00(0.02)						**	
* <sup>44</sup> Sc	J : also 11Av01=2						**	
* <sup>44</sup> Sc <sup>p</sup>	J : also 11Av01=6						**	
* <sup>44</sup> Cr	T : others 14Po05=25(+6-4) 92Bo37=53(+4-3)						**	
* <sup>44</sup> Cr	D : %β <sup>+</sup> p average 07Do17=14.0(0.9) 14Po05=10(1); Birge ratio=2.97;						**	
* <sup>44</sup> Cr	D : other 96Fa09>7(3)						**	
<sup>45</sup> Si	37090#	600#	4# ms	3/2 <sup>-</sup> #			β <sup>-</sup> ?;β <sup>-</sup> n ?;β <sup>-</sup> 2n ?	
<sup>45</sup> P	15960#	500#	10# ms >200ns	1/2 <sup>+</sup> #	08	1990	β <sup>-</sup> ?;β <sup>-</sup> n ?;β <sup>-</sup> 2n ?	
<sup>45</sup> S	-3340#	300#	68 ms 2	3/2 <sup>-</sup> #	08	1989	β <sup>-</sup> =100;β <sup>-</sup> n≈54;β <sup>-</sup> 2n ?	
<sup>45</sup> Cl	-18260	140	413 ms 25	(3/2 <sup>+</sup> )	08 12Ri08	J 1979	β <sup>-</sup> =100;β <sup>-</sup> n=24 4	
<sup>45</sup> Ar	-29770.8	0.5	21.48 s 0.15	(5/2 <sup>-</sup> ,7/2 <sup>-</sup> )	08	1974	β <sup>-</sup> =100	
<sup>45</sup> K	-36615.6	0.5	17.8 m 0.6	3/2 <sup>+</sup> *	08 14Kr04	J 1964	β <sup>-</sup> =100	
<sup>45</sup> Ca	-40812.2	0.4	162.61 d 0.09	7/2 <sup>-</sup> *	08	1940	β <sup>-</sup> =100	
<sup>45</sup> Sc	-41072.3	0.7	STABLE	7/2 <sup>-</sup> *	08	1923	IS=100	
<sup>45</sup> Sc <sup>m</sup>	-41059.9	0.7	12.40 0.05	3/2 <sup>+</sup>	08	1964	IT=100	
<sup>45</sup> Sc <sup>i</sup>	-34373	15	6699 15	7/2 <sup>-</sup>	T=5/2 08			
<sup>45</sup> Ti	-39010.3	0.8	184.8 m 0.5	7/2 <sup>-</sup> *	08	1941	β <sup>+</sup> =100	
<sup>45</sup> Ti <sup>m</sup>	-38973.8	0.8	36.53 0.15	3/2 <sup>-</sup>	08	2006	IT=100	
<sup>45</sup> Ti <sup>i</sup>	-34291	3	4719 3	RQ	7/2 <sup>-</sup>	T=3/2 08		
<sup>45</sup> V	-31886.4	0.9	547 ms 6	7/2 <sup>-</sup>	08	1975	β <sup>+</sup> =100	
<sup>45</sup> V <sup>m</sup>	-31829.6	1.1	56.8 0.6	(3/2 <sup>-</sup> )	08 11Ho02	T 1980	IT=100	
<sup>45</sup> Vi	-27090	9	4797 9	RQ	7/2 <sup>-</sup>	T=3/2 08	p=100	
<sup>45</sup> Cr	-19510	40	60.9 ms 0.4	*	7/2 <sup>-</sup> #	08	1974	β <sup>+</sup> =100;β <sup>+</sup> p=34.4 8
<sup>45</sup> Cr <sup>m</sup>	-19400	40	107 1	*	(3/2)	11 11Ho02	ETJ 2011	IT=100
<sup>45</sup> Mn	-4980#	300#	<70ns	5/2 <sup>-</sup> #	08 92Bo37	I	p ?	
<sup>45</sup> Fe	14410#	280#	2.5 ms 0.2	3/2 <sup>+</sup> #	08 07Mi36	TD 1996	2p=70 4;β <sup>+</sup> =30 4; β <sup>+</sup> p=18.9 35;β <sup>+</sup> 2p=7.8 23	
* <sup>45</sup> Ca	J : also 15Ru02=7/2						**	
* <sup>45</sup> Sc	J : also 11Av01=7/2						**	
* <sup>45</sup> V <sup>m</sup>	T : average 11Ho02=468(23) 87Ha.B=430(80) 82Ho11=539(18) 82Al.C=610(80)						**	
* <sup>45</sup> V <sup>m</sup>	T : 80Gr.A=510(50)						**	
* <sup>45</sup> Fe	T : average 07Mi40=2.6(0.2) (2p gated) 07Mi36=2.8(0.4) (β gated)						**	
* <sup>45</sup> Fe	T : 05Do20=1.6(+0.5-0.3) 02Gi09=4.7(+3.4-1.4) 02Pi02=3.2(+2.6-1.0);						**	
* <sup>45</sup> Fe	T : 02Gi09 supersedes 01Gi01=6(+17-3), 5.98(2.49), 4.22(1.88)						**	
* <sup>45</sup> Fe	D : %2p from 07Mi40; other 05Do20=57(10)%.						**	
<sup>46</sup> P	22840#	500#	9# ms >200ns		00 90Le03	I 1990	β <sup>-</sup> ?;β <sup>-</sup> n ?;β <sup>-</sup> 2n ?	
<sup>46</sup> S	640#	400#	50 ms 8	0 <sup>+</sup>	10	1989	β <sup>-</sup> =100;β <sup>-</sup> n ?;β <sup>-</sup> 2n ?	
<sup>46</sup> Cl	-13730	100	232 ms 2	2 <sup>-</sup> #	12	1989	β <sup>-</sup> =100;β <sup>-</sup> n=60 9;β <sup>-</sup> 2n ?	
<sup>46</sup> Ar	-29771.3	2.3	8.4 s 0.6	0 <sup>+</sup>	00	1974	β <sup>-</sup> =100	
<sup>46</sup> K	-35413.9	0.7	96.30 s 0.08	2 <sup>-</sup> *	00 14Ku.A	T 1965	β <sup>-</sup> =100	
<sup>46</sup> Ca	-43139.6	2.2	STABLE	0 <sup>+</sup>	00	1938	IS=0.004 3;2β <sup>-</sup> ?	
<sup>46</sup> Sc	-41761.6	0.7	83.757 d 0.014	4 <sup>+</sup> *	00 FGK204	T 1936	β <sup>-</sup> =100	
<sup>46</sup> Sc <sup>m</sup>	-41709.6	0.7	52.011 0.001	6 <sup>+</sup>	00	1966	IT=100	
<sup>46</sup> Sc <sup>n</sup>	-41619.1	0.7	142.528 0.007	1 <sup>-</sup>	00	1948	IT=100	
<sup>46</sup> Sc <sup>i</sup>	-36748	4	5014 4	RQ	0 <sup>+</sup>	T=3 00		
<sup>46</sup> Ti	-44128.27	0.09	STABLE	0 <sup>+</sup>	00	1934	IS=8.25 3	
<sup>46</sup> Ti <sup>i</sup>	-34962	7	9166 7	RQ	4 <sup>+</sup>	T=2 00		
<sup>46</sup> Ti <sup>i</sup>	-29977	6	14151 6	RQ	0 <sup>+</sup>	T=3 00		
<sup>46</sup> V	-37075.90	0.13	422.62 ms 0.05	0 <sup>+</sup>	T=1 00 12Pa07	T 1952	β <sup>+</sup> =100	
<sup>46</sup> V <sup>m</sup>	-36274.44	0.16	1.02 ms 0.07	3 <sup>+</sup>	T=0 00	1962	IT=100	
<sup>46</sup> Cr	-29472	11	224.3 ms 1.3	0 <sup>+</sup>	10 15Mo01	T 1972	β <sup>+</sup> =100	
<sup>46</sup> Cr <sup>i</sup>	-20328	13	9143 17	RQ	(4 <sup>+</sup> )	T=2 10	p=?	
<sup>46</sup> Mn	-12420	90	36.2 ms 0.4	*	(4 <sup>+</sup> )	10	1987	β <sup>+</sup> =100;β <sup>+</sup> p=57.0 8; β <sup>+</sup> 2p≈18;β <sup>+</sup> α ?
<sup>46</sup> Mn <sup>m</sup>	-12270#	140#	150# 100#	*	1 <sup>-</sup> #		β <sup>+</sup> ?	

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
<sup>46</sup> Mn <sup>i</sup>	-7390	50	5030	100	p		T=3					
<sup>46</sup> Fe	1210#	300#				13.0 ms 2.0	0 <sup>+</sup>	10 07Do17	TD	1992	$\beta^+$ =100; $\beta^+$ p=78.7 38; $\beta^+$ 2p=?	*
<sup>46</sup> K	J : 19Ko19,14Kr04,14Pa45,82To02=2											**
<sup>46</sup> K	T : other 19Po06=96.5(4)											**
<sup>46</sup> Ca	T : 99Be64 : Onu-BB>100 Ey											**
<sup>46</sup> Sc	J : other 11Av01=4											**
<sup>46</sup> V	T : average 12Pa07=422.66(0.06) 97Ko65=422.57(0.13) 77Ba01=422.28(0.23)											**
<sup>46</sup> V	T : 77Al11=422.47(0.39); others 74Ha59=423.4(2.0) 73Al02=425.3(2.0)											**
<sup>46</sup> Cr	T : others (outweighed) 15Mo01=223.9(9.9) 05On03=240(140) 72Zi02=260(60)											**
<sup>46</sup> Mn	T : others 92Bo37=41(+7-6) 01Gi01=34.0(+4.5-3.5)											**
<sup>46</sup> Mn	D : % $\beta^+$ 2p estimated from Pp = % $\beta^+$ p + 2*% $\beta^+$ 2p = 57(1)											**
<sup>46</sup> Fe	T : average 14Po05=16.4(+4.2-2.8) 07Do17=13.0(2.0) 01Gi01=9.7(+3.5-4.3)											**
<sup>46</sup> Fe	D : other % $\beta^+$ p 14Po05=66(4)% 01Gi01=36(20)%; $\beta^+$ 2p, 1 event in 14Po05											**
<sup>47</sup> P	28810#	600#				4# ms >400ns	1/2 <sup>+</sup> #	18Ta17	I	2018	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?	
<sup>47</sup> S	7200#	400#				24# ms >200ns	3/2 <sup>-</sup> #	07 89Gu03	I	1989	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?	
<sup>47</sup> Cl	-9580#	200#				101 ms 5	3/2 <sup>+</sup> #	07		1989	$\beta^-$ =100; $\beta^-$ n<3; $\beta^-$ 2n ?	
<sup>47</sup> Ar	-25367.3	1.2				1.23 s 0.03	(3/2) <sup>-</sup>	07		1985	$\beta^-$ =100; $\beta^-$ n<0.2	
<sup>47</sup> K	-35712.0	1.4				17.38 s 0.03	1/2 <sup>+</sup> *	07 20Sm02	T	1964	$\beta^-$ =100	*
<sup>47</sup> Ca	-42344.7	2.2				4.536 d 0.003	7/2 <sup>-</sup> *	07		1951	$\beta^-$ =100	*
<sup>47</sup> Sc	-44336.8	1.9				3.3492 d 0.0006	7/2 <sup>-</sup> *	07		1945	$\beta^-$ =100	
<sup>47</sup> Sc <sup>m</sup>	-43570.0	1.9	766.83	0.09		272 ns 8	(3/2) <sup>+</sup>	07		1968	IT=100	
<sup>47</sup> Ti	-44937.61	0.08				STABLE	5/2 <sup>-</sup> *	07		1934	IS=7.44 2	
<sup>47</sup> Ti <sup>i</sup>	-37588.6	0.7	7349.0	0.7			7/2 <sup>-</sup> T=5/2	07				
<sup>47</sup> V	-42007.07	0.11				32.6 m 0.3	3/2 <sup>-</sup> *	07		1942	$\beta^+$ =100	
<sup>47</sup> V <sup>i</sup>	-37856.72	0.16	4150.35	0.11			5/2(-) T=3/2	07			IT=100	
<sup>47</sup> Cr	-34563	5				461.6 ms 1.5	3/2 <sup>-</sup>	07		1972	$\beta^+$ =100	*
<sup>47</sup> Cr <sup>j</sup>	-29803#	21#	4760#	20#			5/2 <sup>-</sup> # T=5/2					
<sup>47</sup> Mn	-22570	30				88.0 ms 1.3	5/2 <sup>-</sup> #	07 07Do17	TD	1987	$\beta^+$ =100; $\beta^+$ p<1.7	
<sup>47</sup> Mn <sup>i</sup>	-15191	17	7380	40	RQ		7/2 <sup>-</sup> # T=5/2	07		2001	p=100	
<sup>47</sup> Fe	-7130#	500#				21.9 ms 0.2	7/2 <sup>-</sup> #	07 07Do17	TD	1992	$\beta^+$ =100; $\beta^+$ p=88.4 9	
<sup>47</sup> Fe <sup>m</sup>	-6360#	510#	770#	100#			3/2 <sup>+</sup> #	Mirror	I		IT ?	
<sup>47</sup> Co	10620#	600#					7/2 <sup>-</sup> #	07 Mirror	I		p ?	
<sup>47</sup> K	J : 19Ko19,14Kr04,14Pa45,13Pa11=1/2											**
<sup>47</sup> Ca	J : also 15Ru02=7/2											**
<sup>47</sup> Cr	T: average 77Ed01=460.0(1.5) 77Ho25=452(18) 85Bu07=508(10)											**
<sup>47</sup> Cr	T: 88HaZB=472.0(6.3) 85HoZS=520(40) 17Ku12=460(80)											**
<sup>48</sup> S	12390#	500#				10# ms >200ns	0 <sup>+</sup>	06		1990	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?	
<sup>48</sup> Cl	-4280#	500#				30# ms >200ns		06 89Gu03	I	1989	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?	
<sup>48</sup> Ar	-22355	17				415 ms 15	0 <sup>+</sup>	10 12We08	TD	2004	$\beta^-$ =100; $\beta^-$ n=38 6	*
<sup>48</sup> K	-32284.5	0.8				6.83 s 0.14	1 <sup>-</sup> *	06		1972	$\beta^-$ =100; $\beta^-$ n=1.14 15	*
<sup>48</sup> Ca	-44224.868	0.018				56 Ey 10	0 <sup>+</sup>	06 20Ba.A	T	1938	IS=0.187 21;2 $\beta^-$ =?; $\beta^-$ ?	*
<sup>48</sup> Sc	-44504	5				43.67 h 0.09	6 <sup>+</sup> *	06		1937	$\beta^-$ =100	
<sup>48</sup> Ti	-48492.95	0.07				STABLE	0 <sup>+</sup>	06		1923	IS=73.72 3	
<sup>48</sup> Ti <sup>i</sup>	-37767	6	10726	6			(6 <sup>+</sup> ) T=3	06				
<sup>48</sup> V	-44478.0	1.0				15.9735 d 0.0025	4 <sup>+</sup> *	06		1937	$\beta^+$ =100	
<sup>48</sup> V <sup>i</sup>	-41459.16	0.23	3018.8	0.9	RQ		(0) <sup>+</sup> T=2	06			IT=100	
<sup>48</sup> Cr	-42821	7				21.56 h 0.03	0 <sup>+</sup>	06		1952	$\beta^+$ =100	
<sup>48</sup> Cr <sup>i</sup>	-37028	7	5792.77	0.24			4 <sup>+</sup> T=1	06		1987	IT=100	
<sup>48</sup> Cr <sup>j</sup>	-34061	15	8760	17	RQ		0 <sup>+</sup> frg. T=2	06				*
<sup>48</sup> Mn	-29297	7				158.1 ms 2.2	4 <sup>+</sup>	06		1987	$\beta^+$ =100; $\beta^+$ p=0.28 4; $\beta^+$ $\alpha$ =6e-4	
<sup>48</sup> Mn <sup>i</sup>	-26260	7	3036.7	0.9	IT		0 <sup>+</sup> T=2	06 MMC12	J		p=100	
<sup>48</sup> Fe	-18010	90				45.3 ms 0.6	0 <sup>+</sup>	06 07Do17	TD	1987	$\beta^+$ =100; $\beta^+$ p=15.3 5	
<sup>48</sup> Co	1730#	500#					6 <sup>+</sup> #	06			p ?	
<sup>48</sup> Ni	18180#	420#				2.8 ms 0.8	0 <sup>+</sup>	06 11Po09	TD	2000	2p=70 20; $\beta^+$ =30 20; $\beta^+$ p ?	*
<sup>48</sup> Ar	T : average 12We08=381(35) 412(19) 04Gr20=475(40)											**
<sup>48</sup> K	J : 14Kr04,14Pa45=1											**
<sup>48</sup> K	T : average 75Mu08=6.8(0.2) 81HuZT=6.9(0.2) 78De17=6(1)											**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>48</sup> Ca	T : 2 $\beta^-$ symmetrized from 20Ba.A=53(+12-8); other 15Ba11=44(+6-5)									**
* <sup>48</sup> Cr <sup>j</sup>	E : strongly fragmented state; other: 10(15) keV lower									**
* <sup>48</sup> Fe	D : % $\beta^+$ p average 07Do17=15.9(6) 16Or03=14.4(7); other 96Fa09>3.6(1.1)									**
* <sup>48</sup> Fe	T : other 16Or03=51(3) 96Fa09=44(7)									**
* <sup>48</sup> Ni	T : average 05Do20=2.1(+2.1-0.7) 14Po05=11Po09=2.1(+1.4-0.4)									**
<sup>49</sup> S	20390#	580#				4# ms >400ns	1/2 <sup>-</sup> #	08 18Ta17 I	2018	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?
<sup>49</sup> Cl	740#	400#				35# ms >200ns	3/2 <sup>+</sup> #	08 89Gu03 I	1989	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?
<sup>49</sup> Ar	-17060#	400#				236 ms 8	3/2 <sup>-</sup> #	08 12We08 TD	1989	$\beta^-$ =100; $\beta^-$ n=29 6; $\beta^-$ 2n ?
<sup>49</sup> K	-29611.5	0.8				1.26 s 0.05	1/2 <sup>+</sup> *	11	1972	$\beta^-$ =100; $\beta^-$ n=86 9
<sup>49</sup> Ca	-41300.00	0.18				8.718 m 0.006	3/2 <sup>-</sup> *	08	1950	$\beta^-$ =100
<sup>49</sup> Sc	-46562.4	2.3				57.18 m 0.13	7/2 <sup>-</sup>	08	1940	$\beta^-$ =100
<sup>49</sup> Ti	-48564.01	0.08				STABLE	7/2 <sup>-</sup> *	08	1934	IS=5.41 2
<sup>49</sup> V	-47962.2	0.8				330 d 15	7/2 <sup>-</sup> *	08	1940	$\epsilon$ =100
<sup>49</sup> Vi	-41531	4	6432	4	RQ		7/2 <sup>-</sup> T=5/2			
<sup>49</sup> Cr	-45332.4	2.2				42.3 m 0.1	5/2 <sup>-</sup> *	08	1942	$\beta^+$ =100
<sup>49</sup> Cr <sup>i</sup>	-40568	5	4764	5			(7/2 <sup>-</sup> ) T=3/2	08 85Fu03 E	1969	IT=100
<sup>49</sup> Mn	-37619.9	2.2				382 ms 7	5/2 <sup>-</sup>	08	1970	$\beta^+$ =100
<sup>49</sup> Mn <sup>i</sup>	-32803	18	4817	18	p		(7/2 <sup>-</sup> ) T=3/2	08		p=100
<sup>49</sup> Fe	-24751	24				64.7 ms 0.3	(7/2 <sup>-</sup> )	08 96Fa09 J	1970	$\beta^+$ =100; $\beta^+$ p=56.7 4
<sup>49</sup> Co	-9780#	500#				<35ns	7/2 <sup>-</sup> #	08 94B110 I		p ?
<sup>49</sup> Ni	8530#	600#				7.5 ms 1.0	7/2 <sup>-</sup> #	08	1996	$\beta^+$ =100; $\beta^+$ p=83.4 13.2
* <sup>49</sup> K	J : 14Kr04, 14Pa45, 13Pa11=1/2									**
* <sup>49</sup> Ca	J : 15Ru02, 16Ga34=3/2									**
* <sup>49</sup> Cr	T : other 18Tu03=44.0(2.7) for q=24+ (bare ion)									**
* <sup>49</sup> Cr <sup>i</sup>	E : strongest component surrounded by several weak l=3 lines.									**
* <sup>49</sup> Cr <sup>i</sup>	E : 85Fu03 cannot confirm IAS identity and fragmentation									**
* <sup>49</sup> Mn	T: average 80Ha12=384(17) 87Ha.A=381.7(7.4) 17Ku12=380(30)									**
<sup>50</sup> Cl	7700#	400#				10# ms >620ns		19 09Ta24 I	2009	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?
<sup>50</sup> Ar	-13230#	500#				106 ms 6	0 <sup>+</sup>	19	1989	$\beta^-$ =100; $\beta^-$ n=37 7; $\beta^-$ 2n ?
<sup>50</sup> K	-25728	8				472 ms 4	0 <sup>-</sup> *	19	1972	$\beta^-$ =100; $\beta^-$ n=28.6 24; $\beta^-$ 2n ?
<sup>50</sup> K <sup>m</sup>	-25556	8	172.0	0.4		125 ns 40	(2 <sup>-</sup> )	19 10Da06 T	1999	IT=100
<sup>50</sup> Ca	-39589.2	1.6				13.45 s 0.05	0 <sup>+</sup>	19	1964	$\beta^-$ =100
<sup>50</sup> Sc	-44537.1	2.5				102.5 s 0.5	5 <sup>+</sup>	19	1959	$\beta^-$ =100
<sup>50</sup> Sc <sup>m</sup>	-44280.2	2.5	256.895	0.010		350 ms 40	2 <sup>+</sup>	19	1963	IT>99; $\beta^-$ <1
<sup>50</sup> Ti	-51431.87	0.08				STABLE	0 <sup>+</sup>	19	1934	IS=5.18 2
<sup>50</sup> V	-49223.24	0.09				271 Py 13	6 <sup>+</sup> *	19 20Da12 T	1949	IS=0.250 10; $\beta^+$ ≈100; $\beta^-$ ?
<sup>50</sup> Vi	-44410.71	0.28	4812.53	0.27	RQ		0 <sup>+</sup> T=3	10		
<sup>50</sup> Cr	-50261.36	0.09				STABLE >1.3Ey	0 <sup>+</sup>	19	1930	IS=4.345 13; 2 $\beta^+$ ?
<sup>50</sup> Cr <sup>i</sup>	-41836	7	8425	7	RQ		6 <sup>+</sup> T=2	19		
<sup>50</sup> Cr <sup>j</sup>	-37039	6	13222	6	RQ		0 <sup>+</sup> T=3	19		
<sup>50</sup> Mn	-42626.89	0.12				283.21 ms 0.07	0 <sup>+</sup> * T=1	19 06Ba33 T	1952	$\beta^+$ =100
<sup>50</sup> Mn <sup>m</sup>	-42401.57	0.11	225.31	0.07	MD	1.75 m 0.03	5 <sup>+</sup> * T=0	19	1962	$\beta^+$ =100
<sup>50</sup> Fe	-34476	8				152.0 ms 0.6	0 <sup>+</sup>	19 15Mo01 T	1977	$\beta^+$ =100; $\beta^+$ p≈0
<sup>50</sup> Fe <sup>i</sup>	-25999	10	8478	13	RQ		(6 <sup>+</sup> ) T=2	19		
<sup>50</sup> Co	-17590	130				38.8 ms 0.2	(6 <sup>+</sup> )	19	1987	$\beta^+$ =100; $\beta^+$ p=70.5 7; $\beta^+$ 2p ?
<sup>50</sup> Co <sup>j</sup>	-12747	15	4840	130	2p		(0 <sup>+</sup> ) T=3	19		p=100
<sup>50</sup> Ni	-3460#	500#				18.5 ms 1.2	0 <sup>+</sup>	19 07Do17 TD	1994	$\beta^+$ =100; $\beta^+$ p=73 6; $\beta^+$ 2p=14 5
* <sup>50</sup> K	J : 14Kr04, 14Pa45=0									**
* <sup>50</sup> K	D : % $\beta^-$ n average 83La23=28(4) 82Ca04=29(3)									**
* <sup>50</sup> V	T : $\beta^+$ average 20Da12=277(+20-19) 19La09=267(+16-18); $\beta^-$ 20Da12>8900 Py									**
* <sup>50</sup> Cr	T : 03Bi05>1.3Ey 85No03>0.18Ey									**
* <sup>50</sup> Mn	T : average 06Ba33=283.10(0.14) 97Ko65=283.29(0.08) 76Wi08=282.72(0.26)									**
* <sup>50</sup> Mn	T : 75Fr02=282.8(0.3) 74Ha59=284.0(0.4)									**
* <sup>50</sup> Fe	T : average 15Mo01=152.1(0.6, beta), 150.1(2.9, gamma); others (outweighed)									**
* <sup>50</sup> Fe	T : 17Ku12=145(13) 97Ko46=155(11)									**
* <sup>50</sup> Ni	T : other 03Ma34=12(+3-2)									**
* <sup>50</sup> Ni	D : % $\beta^+$ p + % $\beta^+$ 2p 07Do17=86.7(3.9), other 03Ma34=70(20); % $\beta^+$ 2p 07Do17=14(5)									**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
<sup>51</sup> Cl	14290#	700#	5# ms >200ns	3/2 <sup>+</sup> #	16	1990	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
<sup>51</sup> Ar	-6490#	400#	30# ms >200ns	1/2 <sup>-</sup> #	16 89Gu03 I	1989	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
<sup>51</sup> K	-22515	13	365 ms 5	3/2 <sup>+</sup> *	16	1983	$\beta^-$ =100; $\beta^-n$ =65 6; $\beta^-2n$ ? *
<sup>51</sup> Ca	-36332.3	0.5	10.0 s 0.8	3/2 <sup>-</sup> *	16	1980	$\beta^-$ =100; $\beta^-n$ ? *
<sup>51</sup> Sc	-43250.4	2.5	12.4 s 0.1	(7/2) <sup>-</sup>	16	1966	$\beta^-$ =100; $\beta^-n$ ?
<sup>51</sup> Ti	-49733.0	0.5	5.76 m 0.01	3/2 <sup>-</sup>	16	1947	$\beta^-$ =100
<sup>51</sup> V	-52203.11	0.10	STABLE	7/2 <sup>-</sup> *	16 76Fu06 J	1924	IS=99.750 10
<sup>51</sup> Cr	-51450.71	0.17	27.7015 d 0.0011	7/2 <sup>-</sup> *	16 FGK204 T	1940	$\epsilon$ =100
<sup>51</sup> Cr <sup>i</sup>	-44838	5	6613 5 RQ	7/2 <sup>-</sup> T=5/2	16		
<sup>51</sup> Mn	-48243.2	0.3	45.81 m 0.21	5/2 <sup>-</sup> *	16	1938	$\beta^+$ =100 *
<sup>51</sup> Mn <sup>i</sup>	-43792.6	1.5	4450.6 1.5 RQ	7/2 <sup>-</sup> T=3/2	16		IT=100
<sup>51</sup> Fe	-40189.2	1.4	305.4 ms 2.3	5/2 <sup>-</sup>	16 15Sh16 T	1972	$\beta^+$ =100 *
<sup>51</sup> Co	-27340	50	68.8 ms 1.9	7/2 <sup>-</sup>	16	1987	$\beta^+$ =100; $\beta^+p$ <3.8
<sup>51</sup> Co <sup>i</sup>	-20674	18	6670 50 p	7/2 <sup>-</sup> # T=5/2	07Do17 D		p=100
<sup>51</sup> Ni	-11650#	500#	23.8 ms 0.2	7/2 <sup>-</sup> #	16 07Do17 TD	1987	$\beta^+$ =100; $\beta^+p$ =87.2 8; $\beta^+2p$ =0.5 2 *
* <sup>51</sup> K	D : % $\beta^-n$ average 06Pe16=63(8) 83La23=68(10); other 82Ca04=47(5) **						
* <sup>51</sup> K	J : 14Pa45,14Kr04,06Pe16,13Pa11=3/2 **						
* <sup>51</sup> Ca	J : 06Pe16,15Ru0,16Ga34=3/2 **						
* <sup>51</sup> Mn	J : 15Ba49,65Sa22,68Jo18=5/2 **						
* <sup>51</sup> Mn	T : average 17Gr12=45.59(0.07) 70Er01=46.2(0.1) 66G102=46.5(0.2); **						
* <sup>51</sup> Fe	T : average 15Sh16=308(5) 13Su07=301(4) 87Ha.B=305(5) 84Ay01=310(5); **						
* <sup>51</sup> Fe	T : other 17Ku12=288(6) is ~7 sigma away from the average value **						
* <sup>51</sup> Ni	D : % $\beta^+2p$ from 12Au08 **						
<sup>52</sup> Cl	22360#	700#	2# ms >400ns		18Ta17 I	2018	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
<sup>52</sup> Ar	-1380#	600#	40# ms >620ns	0 <sup>+</sup>	15 09Ta24 I	2009	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
<sup>52</sup> K	-17140	30	110 ms 4	2 <sup>-</sup> #	15 06Pe16 TD	1983	$\beta^-$ =100; $\beta^-n$ =72.2 9.3; $\beta^-2n$ =2.3 3 *
<sup>52</sup> Ca	-34266.3	0.7	4.6 s 0.3	0 <sup>+</sup>	15 83La23 D	1985	$\beta^-$ =100; $\beta^-n$ <2
<sup>52</sup> Sc	-40524	3	8.2 s 0.2	3( <sup>+</sup> )	15	1980	$\beta^-$ =100; $\beta^-n$ ?
<sup>52</sup> Ti	-49477.7	2.7	1.7 m 0.1	0 <sup>+</sup>	15	1966	$\beta^-$ =100
<sup>52</sup> V	-51443.03	0.16	3.743 m 0.005	3 <sup>+</sup>	15	1934	$\beta^-$ =100
<sup>52</sup> Cr	-55419.51	0.11	STABLE	0 <sup>+</sup>	15	1923	IS=83.789 18
<sup>52</sup> Cr <sup>i</sup>	-44154.6	0.4	11264.9 0.4	3 <sup>+</sup> T=3	15		IT=100
<sup>52</sup> Mn	-50711.39	0.13	5.591 d 0.003	6 <sup>+</sup> *	15	1938	$\beta^+$ =100
<sup>52</sup> Mn <sup>m</sup>	-50333.64	0.13	377.749 0.005	2 <sup>+</sup> *	15	1937	$\beta^+$ =98.22 5; IT=1.78 5 *
<sup>52</sup> Mn <sup>i</sup>	-47785	5	2926 5 RQ	0 <sup>+</sup> T=2	15		IT=100
<sup>52</sup> Fe	-48332.10	0.18	8.275 h 0.008	0 <sup>+</sup>	15	1948	$\beta^+$ =100 *
<sup>52</sup> Fe <sup>m</sup>	-41371.43	0.29	6960.7 0.3 MD	12 <sup>+</sup>	15	1979	$\beta^+$ =99.979 5; IT=0.021 5 *
<sup>52</sup> Fe <sup>i</sup>	-42677.6	0.4	5654.5 0.4	6 <sup>+</sup> T=1	15		IT=100
<sup>52</sup> Fe <sup>j</sup>	-39776	6	8556 6 RQ	0 <sup>+</sup> frg. T=2	15		*
<sup>52</sup> Co	-34344	5	111.7 ms 2.1	6 <sup>+</sup>	15 16Or08 TJ	1987	$\beta^+$ =100; $\beta^+p$ ? *
<sup>52</sup> Co <sup>m</sup>	-33968	8	376 9 MD	2 <sup>+</sup>	16Or08 TJ	2016	$\beta^+$ ≈100; IT ?; $\beta^+p$ ? *
<sup>52</sup> Co <sup>i</sup>	-31420	8	2924 9 IT	0 <sup>+</sup> T=2	16Or03 D	2016	IT=75 23; p=? *
<sup>52</sup> Ni	-22560	80	41.8 ms 1.0	0 <sup>+</sup>	15 16Or03 TD	1987	$\beta^+$ =100; $\beta^+p$ =31.1 5 *
<sup>52</sup> Cu	-1880#	600#		3 <sup>+</sup> #	Mirror I		p ?
* <sup>52</sup> K	T : average 06Pe16=118(6) 85Hu03=110(30) 83La23=105(5) **						
* <sup>52</sup> Mn <sup>m</sup>	T : other: 95Ir01=22.7(3.0) for q=25+ (bare ion) **						
* <sup>52</sup> Fe	T : other: 95Ir01=12.5(+1.5-1.2) for q=26+ (bare ion) 67Pa22=8.23(0.04) **						
* <sup>52</sup> Fe <sup>m</sup>	E : other 6958.0(0.4) keV from a least-squares fit to Eg in Ensdf2015 **						
* <sup>52</sup> Fe <sup>m</sup>	D : %IT from 05Ga20; other 79Ga02<0.4 **						
* <sup>52</sup> Fe <sup>j</sup>	E : probably fragmented, unresolved doublet separated by 4 keV **						
* <sup>52</sup> Co	T : average 17Ku12=111(4) 16Or08=112(3) 15Sh16=112(4) **						
* <sup>52</sup> Co <sup>m</sup>	T : average 16Or08=102(6) 13Su07=103(7) **						
* <sup>52</sup> Ni	T : average 16Or03=42.8(3) 07Do17=40.8(2); other 94Fa06=38(5) **						
* <sup>52</sup> Ni	D : % $\beta^+p$ other 07Do17=31.4(15) 94Fa06=17.0(14) **						
<sup>53</sup> Ar	6790#	700#	20# ms >620ns	5/2 <sup>-</sup> #	11 09Ta24 I	2009	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{53}\text{K}$	-12300	110				30 ms 5	$3/2^+$	09 06Pe16 JD	1983	$\beta^- = 100; \beta^- n \approx 64$	11; *
$^{53}\text{Ca}$	-29390	40				461 ms 90	$1/2^- \#$	14	1983	$\beta^- = 100; \beta^- n \approx 40$	10 *
$^{53}\text{Sc}$	-38770	18				2.4 s 0.6	$(7/2^-)$	14 10Cr02 TJ	1980	$\beta^- = 100; \beta^- n ?$	
$^{53}\text{Ti}$	-46881.4	2.9				32.7 s 0.9	$(3/2^-)$	09	1977	$\beta^- = 100$	
$^{53}\text{V}$	-51852	3				1.543 m 0.014	$7/2^-$	09	1960	$\beta^- = 100$	
$^{53}\text{Cr}$	-55287.62	0.12				STABLE	$3/2^- *$	09	1930	IS=9.501 17	
$^{53}\text{Mn}$	-54690.3	0.3				3.7 My 0.4	$7/2^- *$	09	1955	$\epsilon = 100$	*
$^{53}\text{Mn}^i$	-47717	4	6973	4	RQ		$3/2^- T=5/2$	09	1976		
$^{53}\text{Fe}$	-50947.5	1.7				8.51 m 0.02	$7/2^-$	09	1938	$\beta^+ = 100$	*
$^{53}\text{Fe}^m$	-47907.1	1.7	3040.4	0.3		2.54 m 0.02	$19/2^-$	09	1967	IT=100	
$^{53}\text{Fe}^i$	-46698	3	4250	3			$7/2^- T=3/2$	09			
$^{53}\text{Co}$	-42659.4	1.7				244.6 ms 2.8	$7/2^- \#$	09 17Ku12 T	1970	$\beta^+ = 100$	*
$^{53}\text{Co}^m$	-39485.1	1.9	3174.3	0.9	MD	250 ms 10	$(19/2^-)$	09 72Ce01 D	1970	$\beta^+ = ?; p \approx 1.5$	*
$^{53}\text{Co}^i$	-38334.4	2.6	4325.0	2.0	IT		$(7/2^-) T=3/2$	09 16Su10 ED	1976	IT $\approx 100; p < 0.9$	3
$^{53}\text{Ni}$	-29631	25				55.2 ms 0.7	$(7/2^-)$	13 16Su10 D	1976	$\beta^+ = 100; \beta^+ p = 22.7$	7 *
$^{53}\text{Cu}$	-13140#	500#				<130ns	$3/2^- \#$	13		p ?	
* $^{53}\text{K}$	J : from 20Su06										
* $^{53}\text{Ca}$	D : % $\beta^- n$ 83La23=40(10)% is a lower limit										
* $^{53}\text{Mn}$	J : 15Ba49,56Do45=7/2										
* $^{53}\text{Fe}$	T : other 18Tu03=8.47(0.19) 95Ir01=8.5(0.3) for q=26+ (bare ion)										
* $^{53}\text{Co}$	T : average 17Ku12=245(3) 02Lo13=240(9) 89Ho13=240(20) 73Ko10=262(25);										
* $^{53}\text{Co}$	T : values may contain small contribution from $^{53m}\text{Co}$ decay										
* $^{53}\text{Co}^m$	D : %p from 72Ce01 $\sim 1.5$ %										
* $^{53}\text{Co}^m$	T : average 15Sh16=237(48) 76Vi02=260(20) 72Ce01=247(12)										
* $^{53}\text{Ni}$	D : % $\beta^+ p$ average 16Su10=22(1) 07Do17=23.4(1.0); other: 76Vi02 45										
$^{54}\text{Ar}$	12560#	800#				5# ms >400ns	$0^+$	18Ta17 I	2018	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$	
$^{54}\text{K}$	-5150#	400#				10 ms 5	$2^- \#$	14	1983	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$	
$^{54}\text{Ca}$	-25160	50				90 ms 6	$0^+$	14 08Ma01 TD	1997	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$	*
$^{54}\text{Sc}$	-34438	14				526 ms 15	$(3)^+$	14	1990	$\beta^- = 100; \beta^- n = 16$	9
$^{54}\text{Sc}^m$	-34328	14	110.5	0.3		2.77 $\mu$ s 0.02	$(5^+, 4^+)$	14 10Cr02 J	1998	IT=100	
$^{54}\text{Ti}$	-45744	16				2.1 s 1.0	$0^+$	14 10Cr02 J	1980	$\beta^- = 100$	
$^{54}\text{V}$	-49898	11				49.8 s 0.5	$3^+$	14	1970	$\beta^- = 100$	
$^{54}\text{V}^m$	-49790	11	108.0	1.0		900 ns 500	$(5)^+$	14	1998	IT=100	
$^{54}\text{Cr}$	-56935.38	0.13				STABLE	$0^+$	14	1930	IS=2.365 7	
$^{54}\text{Mn}$	-55558.2	1.0				312.081 d 0.032	$3^+ *$	14 FGK204 T	1938	$\epsilon = 100; \beta^- = 0.93e-4;$	* $e^+ = 1.28e-7$ 25
$^{54}\text{Mn}^i$	-49411.9	2.8	6146.4	3.0	RQ		$0^+ T=3$				
$^{54}\text{Fe}$	-56254.6	0.3				STABLE	$0^+$	14	1923	IS=5.845 105; $2\beta^+ ?$	
$^{54}\text{Fe}^m$	-49727.5	1.1	6527.1	1.1		364 ns 7	$10^+$	14	1983	IT=100	
$^{54}\text{Fe}^i$	-41387	20	14868	20	RQ		$0^+ T=3$	14			
$^{54}\text{Co}$	-48010.1	0.4				193.27 ms 0.06	$0^+ T=1$	14 97Ko65 T	1952	$\beta^+ = 100$	*
$^{54}\text{Co}^m$	-47812.5	0.4	197.57	0.10	MD	1.48 m 0.02	$7^+ T=0$	14	1962	$\beta^+ = 100$	
$^{54}\text{Ni}$	-39278	5				114.1 ms 0.3	$0^+$	14 17Ku12 T	1977	$\beta^+ = 100; \beta^+ p ?$	*
$^{54}\text{Ni}^m$	-32821	5	6457.4	0.9		152 ns 4	$10^+$	14 08Ru09 JD	2008	IT=64 2; p=36 2	
$^{54}\text{Cu}$	-21240#	400#				<75ns	$3^+ \#$	14 94B110 I		p ?	
$^{54}\text{Zn}$	-5700#	220#				1.8 ms 0.5	$0^+$	14 11As08 TD	2005	$2p = 87$	*
* $^{54}\text{Ca}$	T : average 10Cr02=107(14) 08Ma01=86(7)										
* $^{54}\text{Mn}$	D : % $e^+$ average 98Wu01=1.20(0.26)e-7 97Za07=2.2(0.9)e-7										
* $^{54}\text{Co}$	T : average 97Ko65=193.28(0.07) 74Ha59=193.4(0.4) 74Ho21=193.0(0.3)										
* $^{54}\text{Co}$	T : 77Al11=193.28(0.18); other (outweighed) 02Lo13=172(23)										
* $^{54}\text{Ni}$	T : average 17Ku12=110(2) 15Mo01=114.2(0.3), 114.3(1.8) 13Su07=113(9)										
* $^{54}\text{Ni}$	T : 08Fu04=114(5) 02Lo13=103(9) 99Re06=106(12)										
* $^{54}\text{Zn}$	T : symmetrized from 11As08=1.59(+0.60-0.35); other 05B115=3.2(+1.8-0.8)										
* $^{54}\text{Zn}$	D : % $2p$ average 11As08=92(+6-13)% 05B115=87(+10-17)%										
$^{55}\text{K}$	470#	500#				10# ms >620ns	$3/2^+ \#$	09 09Ta24 I	2009	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$	
$^{55}\text{Ca}$	-18650	160				22 ms 2	$5/2^- \#$	09	1997	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$	
$^{55}\text{Sc}$	-30840	60				96 ms 2	$(7/2^-)$	08 10Cr02 TJD	1990	$\beta^- = 100; \beta^- n = 17$	7; $\beta^- 2n ?$ *
$^{55}\text{Ti}$	-41832	29				1.3 s 0.1	$(1/2^-)$	10	1980	$\beta^- = 100; \beta^- n ?$	

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
<sup>55</sup> V	-49125	27				6.54 s 0.15	7/2 <sup>-</sup> #	08	1977	$\beta^-$ =100
<sup>55</sup> Cr	-55110.32	0.23				3.497 m 0.003	3/2 <sup>-</sup>	08	1952	$\beta^-$ =100
<sup>55</sup> Mn	-57712.54	0.26				STABLE	5/2 <sup>-</sup> *	08	1923	IS=100
<sup>55</sup> Fe	-57481.4	0.3				2.7562 y 0.0004	3/2 <sup>-</sup>	09 FGK204	T 1939	$\epsilon$ =100
<sup>55</sup> Fe <sup>i</sup>	-49848	6	7633	6	RQ		5/2 <sup>-</sup> T=5/2	09		
<sup>55</sup> Co	-54030.0	0.4				17.53 h 0.03	7/2 <sup>-</sup> *	09	1938	$\beta^+$ =100
<sup>55</sup> Co <sup>i</sup>	-49308.6	0.4	4721.44	0.10			3/2 <sup>-</sup> frg. T=3/209		1981	IT=100
<sup>55</sup> Ni	-45336.0	0.7				203.9 ms 1.3	7/2 <sup>-</sup>	08 17Ku12	T 1972	$\beta^+$ =100
<sup>55</sup> Ni <sup>i</sup>	-40737.0	1.2	4599	1			7/2 <sup>-</sup> frg. T=3/2	13Tr09	E	
<sup>55</sup> Cu	-31640	160				55.9 ms 1.5	3/2 <sup>-</sup> #	08 13Tr09	T 1987	$\beta^+$ =100; $\beta^+$ p=?
<sup>55</sup> Zn	-14270#	400#				19.8 ms 1.3	5/2 <sup>-</sup> #	08 07Do17	TD 2001	$\beta^+$ =100; $\beta^+$ p=91.0 51
* <sup>55</sup> Sc	T : others 04Li75=115(15) 02Sh43=103(7) 98So03=120(40)									**
* <sup>55</sup> Mn	J : 15Ba49,15He28,79De19=5/2									**
* <sup>55</sup> Co <sup>i</sup>	E : strongly fragmented state; other 26.69(0.15) keV higher									**
* <sup>55</sup> Ni	T : average 17Ku12=203(2) 02Lo13=196(5) 99Re06=204(3) 87Ha.A=212.1(3.8)									**
* <sup>55</sup> Ni	T : 84Ay01=208(5) 77Ho25=189(5) 76Ed.A=219(6)									**
* <sup>55</sup> Ni	J : l=3 in 14Sa46									**
* <sup>55</sup> Ni <sup>i</sup>	E : strongly fragmented state; other 20 keV lower									**
* <sup>55</sup> Cu	T : average 20Gi02=55.5(1.8) 13Tr09=57(3); other 07Do17=27(8), conflicting									**
* <sup>55</sup> Cu	D : % $\beta^+$ p in 07Do17=15.0(4.3), but it is probably a contaminant given the									**
* <sup>55</sup> Cu	D : short and conflicting half-life; not confirmed in 13Tr09									**
<sup>56</sup> K	7980#	600#				5# ms >620ns	2 <sup>-</sup> #	11 09Ta24	I 2009	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?
<sup>56</sup> Ca	-13510	250				11 ms 2	0 <sup>+</sup>	11	1997	$\beta^-$ =100; $\beta^-$ n ?; $\beta^-$ 2n ?
<sup>56</sup> Sc	-25520	260			*	26 ms 6	(1 <sup>+</sup> )	11 10Cr02	JT 1997	$\beta^-$ =100; $\beta^-$ n ?; $\beta^-$ 2n ?
<sup>56</sup> Sc <sup>m</sup>	-25520#	280#	0#	100#	*	75 ms 6	(6 <sup>+</sup> , 5 <sup>+</sup> )	11 10Cr02	JTD 2004	$\beta^-$ =100; $\beta^-$ n >12; $\beta^-$ 2n ?
<sup>56</sup> Sc <sup>n</sup>	-24750	260	775.0	0.1		290 ns 17	(4 <sup>+</sup> )	11 20Mi13	TE 2004	IT=100
<sup>56</sup> Ti	-39420	100				200 ms 5	0 <sup>+</sup>	11 98Am04	D 1980	$\beta^-$ =100; $\beta^-$ n ?
<sup>56</sup> V	-46180	180				216 ms 4	(1 <sup>+</sup> )	11 98Am04	D 1980	$\beta^-$ =100; $\beta^-$ n ?
<sup>56</sup> Cr	-55285.1	0.6				5.94 m 0.10	0 <sup>+</sup>	11 60Dr03	D 1960	$\beta^-$ =100
<sup>56</sup> Mn	-56911.67	0.29				2.5789 h 0.0001	3 <sup>+</sup> *	11	1934	$\beta^-$ =100
<sup>56</sup> Fe	-60607.16	0.27				STABLE	0 <sup>+</sup>	11	1923	IS=91.754 106
<sup>56</sup> Fe <sup>i</sup>	-49103.5	0.4	11503.7	0.3			3 <sup>+</sup> T=3	11		
<sup>56</sup> Co	-56040.5	0.5				77.236 d 0.026	4 <sup>+</sup> *	11	1941	$\beta^+$ =100
<sup>56</sup> Co <sup>i</sup>	-52448	9	3593	9	RQ		(0 <sup>+</sup> ) frg. T=2	11		*
<sup>56</sup> Ni	-53907.6	0.4				6.075 d 0.010	0 <sup>+</sup>	11	1952	$\beta^+$ =100
<sup>56</sup> Ni <sup>p</sup>	-44172.1	1.9	9735.5	1.9			7	11	2008	p≈100
<sup>56</sup> Ni <sup>i</sup>	-47475.7	0.8	6431.9	0.7			4 <sup>+</sup> T=1	11		
<sup>56</sup> Ni <sup>j</sup>	-43964	4	9944	4	RQ		0 <sup>+</sup> frg. T=2			*
<sup>56</sup> Cu	-38630	6				80.8 ms 0.6	(4 <sup>+</sup> )	11 01Bo54	TJD 1987	$\beta^+$ =100; $\beta^+$ p=0.40 12
<sup>56</sup> Cu <sup>i</sup>	-35099	10	3531	12	p		T=2	16Or03	D 2007	IT=51 6;p=49 6
<sup>56</sup> Zn	-25390#	400#				32.4 ms 0.7	0 <sup>+</sup>	11 14Or04	TD 2001	$\beta^+$ =100; $\beta^+$ p=88.0 23
<sup>56</sup> Zn <sup>i</sup>	-21530#	650#	3860#	510#			3 <sup>+</sup> # T=3			p ?
<sup>56</sup> Ga	-3840#	500#					3 <sup>+</sup> #			p ?
* <sup>56</sup> Sc <sup>n</sup>	T : average 20Mi13=290(20) 10Cr02=290(30); other 12Ka36=350(+260-120)									**
* <sup>56</sup> Co <sup>i</sup>	E : strongly fragmented state; other 70(9) keV lower									**
* <sup>56</sup> Ni <sup>j</sup>	E : strongly fragmented state; others 68(6) and 98(6) keV higher									**
* <sup>56</sup> Cu	T : average 20Gi02=80.2(7) 17Ku12=80(2) 02Lo13=82(9) 01Bo54=93(3)									**
* <sup>56</sup> Cu	T : 98Ra15=78(15)									**
* <sup>56</sup> Zn	T : average 14Or04=32.9(0.8) 07Do17=30.0(1.7)									**
* <sup>56</sup> Zn	D : % $\beta^+$ p average 14Or04=88.5(26) 07Do17=86.0(49)									**
<sup>57</sup> K	14130#	600#				2# ms >400ns	3/2 <sup>+</sup> #	18Ta17	I 2018	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?
<sup>57</sup> Ca	-6560#	400#				8# ms >620ns	5/2 <sup>-</sup> #	10 09Ta24	I 2009	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?
<sup>57</sup> Sc	-21380	180				22 ms 2	7/2 <sup>-</sup> #	10 10Cr02	T 1997	$\beta^-$ =100; $\beta^-$ n ?; $\beta^-$ 2n ?
<sup>57</sup> Ti	-34400	210				95 ms 8	5/2 <sup>-</sup> #	10 99So20	T 1985	$\beta^-$ =100; $\beta^-$ n ?
<sup>57</sup> V	-44440	80				350 ms 10	(7/2 <sup>-</sup> )	10 03Ma02	T 1980	$\beta^-$ =100; $\beta^-$ n ?
<sup>57</sup> Cr	-52525.0	1.9				21.1 s 1.0	(3/2 <sup>-</sup> )	10	1978	$\beta^-$ =100
<sup>57</sup> Mn	-57486.3	1.5				85.4 s 1.8	5/2 <sup>-</sup> *	98 15Ba49	J 1954	$\beta^-$ =100
<sup>57</sup> Fe	-60182.02	0.27				STABLE	1/2 <sup>-</sup> *	98	1935	IS=2.119 29
<sup>57</sup> Co	-59345.7	0.5				271.811 d 0.032	7/2 <sup>-</sup> *	98 FGK204	T 1941	$\epsilon$ =100



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{57}\text{Co}^i$	-52092.3	0.4	7253.3	0.6	RQ	$1/2^-$ T=5/2	MMC120J		
$^{57}\text{Ni}$	-56084.0	0.6			35.60 h 0.06	$3/2^-$	98	1938	$\beta^+=100$
$^{57}\text{Ni}^i$	-50845.2	0.9	5238.8	0.7		$7/2^-$ frg. T=3/298			*
$^{57}\text{Cu}$	-47309.0	0.5			196.4 ms 0.7	$3/2^-*$	98 17Ku12 T	1976	$\beta^+=100$
$^{57}\text{Cu}^i$	-42010	25	5299	25	p	$7/2^-$ T=3/2			*
$^{57}\text{Zn}$	-32550#	200#			45.7 ms 0.6	$7/2^-$ #	98 20Gi02 T	1976	$\beta^+=100;\beta^+p=87.9$
$^{57}\text{Ga}$	-15410#	400#				$1/2^-$ #			p ?
$^{57}\text{Sc}$	T : other 03So21=13(4)								
$^{57}\text{Ti}$	T : average 05Li53=98(5) 99So20=67(25) 96Do23=56(20); other								
$^{57}\text{Ti}$	T : 98Am04=180(30) conflicting, not used								
$^{57}\text{V}$	J : 98So03 proposed $3/2^-$ , supported in 03Ma02; same group 05Li53 favors $7/2^-$								
$^{57}\text{Ni}^i$	E : strongly fragmented state; others 98(7) keV lower 128(7) keV higher								
$^{57}\text{Ni}^i$	E : in 79Ik04 and 104(5) keV lower, 129(5) keV higher in 78Na11								
$^{57}\text{Cu}$	T : average 17Ku12=195(4) 02Lo13=183(17) 96Se01=196.3(0.7)								
$^{57}\text{Cu}$	T : 87Ha.A=199.4(3.2) 84Sh28=223(16)								
$^{57}\text{Cu}$	J : 10Co01=3/2								
$^{57}\text{Zn}$	T : others (outweighed) 07Bi09=48(3) 02Lo13=37(5) 76Vi02=40(10)								
$^{57}\text{Zn}$	D : % $\beta^+p$ average 20Ci04=90(10) 07Bi09=78(17)								
$^{58}\text{K}$	21930#	700#			2# ms >400ns	$2^-$ #	18Ta17 I	2019	$\beta^- ?; \beta^-n ?; \beta^-2n ?$
$^{58}\text{Ca}$	-1530#	500#			4# ms >620ns	$0^+$	10 09Ta24 I	2009	$\beta^- ?; \beta^-n ?; \beta^-2n ?$
$^{58}\text{Sc}$	-15480	190			12 ms 5	$3^+ \#$	10	1997	$\beta^- =100; \beta^-n ?; \beta^-2n ?$
$^{58}\text{Sc}^m$	-14060	190	1420.7	2.2	0.60 $\mu\text{s}$ 0.13		20Mi13 ET	2020	IT=100
$^{58}\text{Ti}$	-30920	180			55 ms 6	$0^+$	14 11Da08 T	1992	$\beta^- =100; \beta^-n ?$
$^{58}\text{V}$	-40430	100			191 ms 10	$(1^+)$	10	1980	$\beta^- =100; \beta^-n ?$
$^{58}\text{Cr}$	-51991.8	3.0			7.0 s 0.3	$0^+$	10	1980	$\beta^- =100$
$^{58}\text{Mn}$	-55827.6	2.7			3.0 s 0.1	$1^+*$	10 15He28 J	1961	$\beta^- =100$
$^{58}\text{Mn}^m$	-55755.8	2.7	71.77	0.05	65.4 s 0.5	$4^+*$	10 15He28 J	1961	$\beta^- \approx 100; IT ?$
$^{58}\text{Fe}$	-62155.3	0.3			STABLE	$0^+$	10	1935	IS=0.282 12
$^{58}\text{Co}$	-59847.3	1.2			70.844 d 0.020	$2^+*$	10 FGK204 T	1941	$\beta^+ =100; e^+ =14.79 24; \epsilon =85.21 24$
$^{58}\text{Co}^m$	-59822.4	1.2	24.95	0.06	8.853 h 0.023	$5^+$	10 19Mo11 TD	1950	IT=99.99880 5; $\epsilon =0.00120 5$
$^{58}\text{Co}^n$	-59794.2	1.2	53.15	0.07	10.5 $\mu\text{s}$ 0.3	$4^+$	10	1964	IT=100
$^{58}\text{Co}^i$	-54095	8	5752	8	RQ	$0^+$ frg. T=3	10		*
$^{58}\text{Ni}$	-60228.9	0.3			STABLE >700Ey	$0^+$	10 93Va19 T	1921	IS=68.0769 190; $2\beta^+ ?$
$^{58}\text{Ni}^i$	-51400	40	8830	40	RQ	$2^+ T=2$	10		*
$^{58}\text{Ni}^j$	-45690	7	14539	7	RQ	$0^+ T=3$	10 MMC12 J		
$^{58}\text{Cu}$	-51667.9	0.6			3.204 s 0.007	$1^+* T=0$	10 11Vi03 J	1952	$\beta^+ =100$
$^{58}\text{Cu}^i$	-51464.9	0.6	202.99	0.24		$0^+ T=1$	10		*
$^{58}\text{Zn}$	-42300	50			86.0 ms 1.9	$0^+$	14 20Ci04 D	1986	$\beta^+ =100; \beta^+p=0.7 1$
$^{58}\text{Ga}$	-23540#	300#			*	$2^+ \#$	Mirror I		p ?
$^{58}\text{Ga}^m$	-23510#	320#	30#	100#	*	$5^+ \#$	Mirror I		p ?
$^{58}\text{Ge}$	-7580#	500#				$0^+$	Mirror I		2p ?
$^{58}\text{Sc}^m$	T : average 20Mi13=0.6(0.2) 1.3(0.8) 0.9(0.5) 0.5(0.2) from $\gamma(t)$								
$^{58}\text{Sc}^m$	E : 20Mi13=180.5(0.6), 247(2), 412.3(0.6), 580.9(0.4) gammas in a cascade								
$^{58}\text{Ti}$	T : average 11Da08=57(10) 03So21=59(9) 99So20=47(10)								
$^{58}\text{Co}$	D : from 71GoYM								
$^{58}\text{Co}^i$	E : strongly fragmented state; other 20(8) keV lower								
$^{58}\text{Cu}$	J : also 10Co01=1								
$^{58}\text{Zn}$	T : average 17Ku12=86(2) 09Fu15=90(8) 05Ka46=83(10) 02Lo13=83(10)								
$^{58}\text{Zn}$	T : 98Jo18=86(18)								
$^{59}\text{K}$	28750#	800#			1# ms >400ns	$3/2^+ \#$	18Ta17 I	2018	$\beta^- ?; \beta^-n ?; \beta^-2n ?$
$^{59}\text{Ca}$	5810#	600#			5# ms >400ns	$5/2^- \#$	18Ta17 I	2018	$\beta^- ?; \beta^-n ?; \beta^-2n ?$
$^{59}\text{Sc}$	-10830	250			12# ms >620ns	$7/2^- \#$	18 09Ta24 I	2009	$\beta^- ?; \beta^-n ?; \beta^-2n ?$
$^{59}\text{Ti}$	-25880#	300#			28.5 ms 1.9	$5/2^- \#$	18 11Da08 T	1997	$\beta^- =100; \beta^-n ?; \beta^-2n ?$
$^{59}\text{Ti}^m$	-25770#	300#	108.5	0.5	615 ns 11	$1/2^- \#$	18 19Wi04 TJE	2012	IT=100
$^{59}\text{V}$	-37610	140			95 ms 6	$(5/2^-)$	18 05Li53 TJD	1985	$\beta^- =100; \beta^-n >3$
$^{59}\text{Cr}$	-48115.9	0.7			1050 ms 90	$(1/2^-)$	18 05Li53 TJ	1980	$\beta^- =100$
$^{59}\text{Cr}^m$	-47613.2	1.3	502.7	1.1	96 $\mu\text{s}$ 20	$(9/2^+)$	18	1998	IT=100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{59}\text{Mn}$	-55525.3	2.3	4.59 s 0.05	$5/2^-*$	18 15Ba49 J	1976	$\beta^- = 100$
$^{59}\text{Fe}$	-60665.0	0.3	44.500 d 0.012	$3/2^-*$	18 FGK204 T	1938	$\beta^- = 100$
$^{59}\text{Co}$	-62229.8	0.4	STABLE	$7/2^-*$	18	1923	IS=100
$^{59}\text{Ni}$	-61156.8	0.4	81 ky 5	$3/2^-$	18 94Ru19 T	1951	$\beta^+ = 100$
$^{59}\text{Ni}^i$	-53814.9	2.1	7341.9 2.1 RQ	$7/2^-$ frg. T=5/2			*
$^{59}\text{Cu}$	-56358.5	0.5	81.5 s 0.5	$3/2^-*$	18	1947	$\beta^+ = 100$
$^{59}\text{Cu}^i$	-52473.0	2.2	3885.5 2.1	$3/2^-$ frg. T=3/218			IT=100
$^{59}\text{Zn}$	-47215.7	0.8	178.7 ms 1.3	$3/2^-$	18	1981	$\beta^+ = 100; \beta^+ p = 0.10$
$^{59}\text{Ga}$	-33760#	170#	<43ns	$3/2^- \#$	18		p ?
$^{59}\text{Ge}$	-16370#	400#	13.3 ms 1.7	$7/2^- \#$	18 20Gi02 TD	2015	$\beta^+ \approx 100; \beta^+ p = 93.7; 2p < 0.2$
$^{59}\text{Ti}$	T : average 11Da08=27.5(2.5) 03So21=30(3); other 99So20=58(17)						
$^{59}\text{Ti}^m$	T : average 20Mi13=610(20) 19Wi04=618(13) 12Ka36=587(+57-51); other						
$^{59}\text{Ti}^m$	T : 05Ga01=590(130)						
$^{59}\text{Ti}^m$	E : other 20Mi13=108.9(0.4)						
$^{59}\text{V}$	T : average 05Li53=97(2) 99So20=75(7) (supersedes 98So03=70(40)); other						
$^{59}\text{V}$	T : 98Am04=130(20) conflicting, not used						
$^{59}\text{Cr}$	T : others 96Do23=460(50), 88Bo06=600(300), 85Bo49=1000(400)						
$^{59}\text{Ni}$	T : average 94Ru19=108(13) 94Ru19(meteorite)=120(22) 81Ni08=76(5)						
$^{59}\text{Ni}^i$	E : strongest fragmented state; others 40.1(0.3) keV higher, 17.7(0.3) keV						
$^{59}\text{Ni}^i$	E : higher and 36.3(0.2)keV lower						
$^{59}\text{Cu}$	J : 11Vi03, 11Ko36, 10Co01=3/2						
$^{59}\text{Cu}^i$	E : strongest fragmented state; other 21(6) keV higher						
$^{59}\text{Zn}$	T : average 17Ku12=174(2) 14Ro14=210(34) 02Lo13=173(14)						
$^{59}\text{Zn}$	T : 84Ar12=182.2(1.8) 81Ho19=210(20)						
$^{59}\text{Ge}$	T : other 16Go26 (same as 20Gi02)						
$^{59}\text{Ge}$	D : 2p not observed in 20Gi02 and 16Go26; limit from 16Go26 based on the						
$^{59}\text{Ge}$	D : assumption that one event is not $\beta^+ p$						
$^{60}\text{Ca}$	11000#	700#	2# ms >400ns	$0^+$	18Ta17 I	2018	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$
$^{60}\text{Sc}$	-4550#	500#	10# ms >620ns	$3^+ \#$	09Ta24 I	2009	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$
$^{60}\text{Ti}$	-22100	240	22.2 ms 1.6	$0^+$	14 11Da08 T	1997	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$
$^{60}\text{V}$	-33090	180	122 ms 18	$3^+ \#$	13	1985	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$
$^{60}\text{V}^m$	-33090#	230#	0# 150# *	$1^+ \#$	13	1999	$\beta^- = ?; IT ?; \beta^- n ?; \beta^- 2n ?$
$^{60}\text{V}^n$	-32890	180	203.7 0.7	$(4^+)$	13 12Ka36 ET	1999	IT=100
$^{60}\text{Cr}$	-46908.5	1.1	490 ms 10	$0^+$	13	1980	$\beta^- = 100; \beta^- n ?$
$^{60}\text{Mn}$	-52967.9	2.3	280 ms 20	$1^+*$	13 15He28 J	1978	$\beta^- = 100$
$^{60}\text{Mn}^m$	-52696.0	2.3	271.90 0.10	$4^+*$	13 15He28 J	1978	$\beta^- = 88.5; 8; IT = 11.5$
$^{60}\text{Fe}$	-61413	3	2.62 My 0.04	$0^+$	13 09Ru08 T	1957	$\beta^- = 100$
$^{60}\text{Co}$	-61650.4	0.4	5.2714 y 0.0006	$5^+*$	13 FGK204 T	1941	$\beta^- = 100$
$^{60}\text{Co}^m$	-61591.8	0.4	58.59 0.01	$2^+*$	13	1963	IT $\approx$ 100; $\beta^- = 0.25$
$^{60}\text{Ni}$	-64473.2	0.4	STABLE	$0^+$	13	1921	IS=26.2231 150
$^{60}\text{Ni}^i$	-53347	4	11126 4 RQ	$5^+ T=3$			
$^{60}\text{Cu}$	-58345.3	1.6	23.7 m 0.4	$2^+*$	13 11Vi03 J	1947	$\beta^+ = 100$
$^{60}\text{Cu}^i$	-55804	5	2541 5 RQ	$(0^+) T=2$	13		IT=100
$^{60}\text{Zn}$	-54174.5	0.5	2.38 m 0.05	$0^+$	13	1955	$\beta^+ = 100$
$^{60}\text{Zn}^i$	-49322.3	0.9	4852.2 0.7	$(2^+) T=1$	13		IT=100
$^{60}\text{Zn}^j$	-46807	24	7367 24 RQ	$0^+ T=2$	13		
$^{60}\text{Ga}$	-39590#	200#	72.4 ms 1.7	$(2^+)$	13 20Gi02 T	1995	$\beta^+ = 100; \beta^+ p = 1.6$ ; $7; \beta^+ \alpha < 0.023$
$^{60}\text{Ga}^i$	-37050#	210#	2540# 50#				
$^{60}\text{Ge}$	-27530#	300#	21 ms 6	$0^+$	13 16Ci01 TD	2005	$\beta^+ = 100; \beta^+ p \approx 100; \beta^+ 2p < 14$
$^{60}\text{As}$	-5640#	400#		$5^+ \#$	Mirror I		p ?
$^{60}\text{As}^m$	-5580#	400#	60# 20#	$2^+ \#$	Mirror I		p ?
$^{60}\text{Ti}$	T : average 11Da08=22.4(2.5) 03So21=22(2)						
$^{60}\text{V}^m$	E : 12Ka36=99.7(0.5) and 104.0(0.5) gamma rays in a cascade to gs						
$^{60}\text{V}^n$	T : symmetrized from 12Ka36=229(+25-23); others 10Da06=320(90) 99Da.A=320(90)						
$^{60}\text{Mn}^m$	I : other isomer T=1.0(+0.3-0.2) us decays by 114 keV g-ray (not placed)						
$^{60}\text{Fe}$	T : adopted from 09Ru08; others: 17Os02=2.72(0.16) 15Wa06=2.50(0.12)						
$^{60}\text{Ga}$	T : average 20Gi02=70.8(2.0) 17Ku12=76(3); others 02Lo13=70(13)						
$^{60}\text{Ga}$	T : 01Ma96=70(15), outweighed						
$^{60}\text{Ge}$	T : symmetrized from 16Ci01=20(+7-5)						

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{61}\text{Ca}$	19010#	800#			1# ms	$1/2^-$ #			$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
$^{61}\text{Sc}$	500#	600#			7# ms >620ns	$7/2^-$ #	15 09Ta24	I 2009	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
$^{61}\text{Ti}$	-16370#	300#			15 ms 4	$1/2^-$ #	15	1997	$\beta^-$ =100; $\beta^-n$ ?; $\beta^-2n$ ?
$^{61}\text{Ti}^m$	-16250#	300#	125.0	0.5	200 ns 28	$5/2^-$ #	15 19Wi04	TEJ 2019	IT=100
$^{61}\text{Ti}^n$	-15670#	300#	700.1	0.7	354 ns 69	$9/2^+$ #	15 19Wi04	TEJ 2019	IT=100
$^{61}\text{V}$	-30180	230			48.2 ms 0.6	$(3/2^-)$	15 20On01	TDJ 1992	$\beta^-$ =100; $\beta^-n$ =14.5 20; $\beta^-2n$ ?
$^{61}\text{Cr}$	-42496.5	1.9			243 ms 9	$(5/2^-)$	15 09Cr02	T 1985	$\beta^-$ =100; $\beta^-n$ ?
$^{61}\text{Mn}$	-51742.1	2.3			709 ms 8	$5/2^-*$	15 15Ba49	J 1980	$\beta^-$ =100; $\beta^-n$ ?
$^{61}\text{Fe}$	-58920.5	2.6			5.98 m 0.06	$(3/2^-)$	15	1957	$\beta^-$ =100
$^{61}\text{Fe}^m$	-58058.8	2.6	861.67	0.11	238 ns 5	$9/2^+$	15	1998	IT=100
$^{61}\text{Co}$	-62898.2	0.8			1.649 h 0.005	$7/2^-$	15	1947	$\beta^-$ =100
$^{61}\text{Ni}$	-64222.0	0.4			STABLE	$3/2^-*$	15	1934	IS=1.1399 13
$^{61}\text{Cu}$	-61984.1	1.0			3.343 h 0.016	$3/2^-*$	15	1937	$\beta^+$ =100
$^{61}\text{Cu}^i$	-55611	7	6373	7	RQ	$3/2^-$ frg. T=5/2			
$^{61}\text{Zn}$	-56349	16			89.1 s 0.2	$3/2^-$	15	1955	$\beta^+$ =100
$^{61}\text{Zn}^i$	-53190#	100#	3160#	100#		$3/2^-$ # T=3/2			
$^{61}\text{Zn}^j$	-46360	70	9990	70		$3/2^-$ T=5/2	15		
$^{61}\text{Ga}$	-47130	40			165.9 ms 2.5	$3/2^-$	15 17Ku12	T 1987	$\beta^+$ =100; $\beta^+p$ <0.25
$^{61}\text{Ga}^m$	-47040#	110#	90#	100#		$1/2^-$ #	Mirror	I	
$^{61}\text{Ga}^i$	-43780	30	3360	50	p	$(3/2^-)$ T=3/2	15	1987	p=100
$^{61}\text{Ge}$	-33790#	300#			40.7 ms 0.4	$3/2^-$ #	15 20Gi02	TD 1987	$\beta^+$ =100; $\beta^+p$ =87 3
$^{61}\text{As}$	-17200#	300#				$3/2^-$ #	Mirror	I	p ?
$^{*61}\text{Ti}^m$	E : other 20Mi13=125.2(0.6)								
$^{*61}\text{Ti}^n$	T : other 20Mi13=300(100)								
$^{*61}\text{Ti}^n$	E : other 20Mi13=701.3(0.7)								
$^{*61}\text{Ti}^n$	T : other 20Mi13=200(100)								
$^{*61}\text{V}$	T : average 20On01=48(1) 14Su07=49(1) 11Da08=52.6(4.2) 03So02=47.0(1.2),								
$^{*61}\text{V}$	T : supersedes 99So20=43(7)								
$^{*61}\text{Mn}$	D : $\beta^-n$ observed by 99Ha05; 13Ra17 quotes % $\beta^-n$ =0.6(0.1) (unpublished)								
$^{*61}\text{Cu}$	T : average 15Cv01=3.323(0.010) 82Gr10=3.333 (0.005) 72Cr02=3.34(0.01)								
$^{*61}\text{Cu}$	T : 69Ri04=3.408(0.010); Birge ratio 4.1								
$^{*61}\text{Cu}^i$	E : strongly fragmented state; other 18(7) keV higher								
$^{*61}\text{Ga}$	T : average 17Ku12=163(5) 14Ro14=162(10) 02We07=168(3) 02Lo13=148(19)								
$^{*61}\text{Ga}$	T : 99Oi01=140(70) 93Wi18=150(30)								
$^{62}\text{Sc}$	7310#	600#			2# ms >400ns		18Ta17	I 2018	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
$^{62}\text{Ti}$	-12200#	400#			9# ms >620ns	$0^+$	12 09Ta24	I 2009	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
$^{62}\text{V}$	-25210	260			33.6 ms 2.3	$3^+$ #	12	1997	$\beta^-$ =100; $\beta^-n$ ?; $\beta^-2n$ ?
$^{62}\text{Cr}$	-40853	3			206 ms 12	$0^+$	12	1985	$\beta^-$ =100; $\beta^-n$ ?
$^{62}\text{Mn}$	-48524	7		*	92 ms 13	$1^+*$	12 15He28	J 1983	$\beta^-$ =100; $\beta^-n$ ?
$^{62}\text{Mn}^m$	-48181.0	2.6	343	6	*	$4^+*$	12 15He28	J 1983	$\beta^-$ =100; $\beta^-n$ ?; IT ?
$^{62}\text{Fe}$	-58878.1	2.8			68 s 2	$0^+$	12	1975	$\beta^-$ =100
$^{62}\text{Co}$	-61424	19			1.54 m 0.10	$(2)^+$	12	1949	$\beta^-$ =100
$^{62}\text{Co}^m$	-61402	20	22	5	13.86 m 0.09	$(5)^+$	12 70Jo12	D 1957	$\beta^- \approx 100$ ; IT<0.5
$^{62}\text{Ni}$	-66746.4	0.4			STABLE	$0^+$	12	1934	IS=3.6345 40
$^{62}\text{Cu}$	-62787.5	0.6			9.672 m 0.008	$1^+*$	12 14Un01	T 1936	$\beta^+$ =100
$^{62}\text{Cu}^i$	-58174	6	4614	6	RQ	$(0)^+$ T=3	12		
$^{62}\text{Zn}$	-61168.1	0.6			9.193 h 0.015	$0^+$	12	1948	$\beta^+$ =100
$^{62}\text{Ga}$	-51987.0	0.6			116.122 ms 0.021	$0^+$ T=1	12 13Da16	T 1978	$\beta^+$ =100
$^{62}\text{Ga}^j$	-51415.8	0.6	571.2	0.1		$1(^+)$ T=2	12 98Vi06	EJ 1998	IT=100
$^{62}\text{Ge}$	-42140#	140#			82.5 ms 1.4	$0^+$	12 17Ku12	T 1991	$\beta^+$ =100; $\beta^+p$ ?
$^{62}\text{As}$	-24420#	300#				$1^+\#$			p ?
$^{*62}\text{Mn}$	D : % $\beta^-n$ 99So20~0 99Ha05>0								
$^{*62}\text{Mn}^m$	E : symmetrized from 15Ga38=346(+3-8) keV								
$^{*62}\text{Cu}^i$	E : Ensdf2012=4628(10) keV								
$^{*62}\text{Ga}$	T : average 13Da16=116.15(0.13) 08Gr03=116.100(0.025) 05Hy04=116.01(0.19)								
$^{*62}\text{Ga}$	T : 05Ca06=116.09(0.17) 04Bl03=116.19(0.04) 03Hy02=115.84(0.25)								
$^{*62}\text{Ga}$	T : 79Da04=116.34(0.35) 78Al23=115.95(0.30); others (outweighed)								
$^{*62}\text{Ga}$	T : 02Bl17,02Lo13=114(2) 78Ch11=116.4(1.5) 93Wi03,93Wi18=113(+6-5)								
$^{*62}\text{Ga}$	T : 93Wi03,93Wi18=113(+6-5)								

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
* <sup>62</sup> Ge	T : average 14Gr10=82.9(1.4) 17Ku12=76(6)						**	
* <sup>62</sup> As	D : most likely p-unstable from estimated Sp=-1980#(420#) keV						**	
<sup>63</sup> Sc	13070#	700#	1# ms	7/2 <sup>-</sup> #			$\beta^- ?; \beta^- n ?; \beta^- 2n ?$	
<sup>63</sup> Ti	-5860#	500#	10# ms >620ms	1/2 <sup>-</sup> #	09 09Ta24 I	2009	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$	
<sup>63</sup> V	-21740	340	19.6 ms 0.9	(3/2 <sup>-</sup> , 5/2 <sup>-</sup> )	09 14Su07 TJ	1997	$\beta^- =100; \beta^- n >35; \beta^- 2n ?$	
<sup>63</sup> Cr	-36180	70	129 ms 2	1/2 <sup>-</sup> #	09	1992	$\beta^- =100; \beta^- n ?$	
<sup>63</sup> Mn	-46887	4	275 ms 4	5/2 <sup>-</sup> *	09 15Ba49 J	1985	$\beta^- =100; \beta^- n = ?$	
<sup>63</sup> Fe	-55636	4	6.1 s 0.6	(5/2 <sup>-</sup> )	09	1980	$\beta^- =100$	
<sup>63</sup> Co	-61852	19	26.9 s 0.4	7/2 <sup>-</sup>	09 94It.A T	1960	$\beta^- =100$	
<sup>63</sup> Ni	-65512.9	0.4	101.2 y 1.5	1/2 <sup>-</sup> *	09 17Dy01 J	1951	$\beta^- =100$	
<sup>63</sup> Ni <sup>m</sup>	-65425.8	0.4	87.15 0.11	1.67 $\mu$ s 0.03	5/2 <sup>-</sup>	09	IT=100	
<sup>63</sup> Cu	-65579.9	0.4	STABLE	3/2 <sup>-</sup> *	09	1923	IS=69.15 15	
<sup>63</sup> Zn	-62213.4	1.6	38.47 m 0.05	3/2 <sup>-</sup> *	09	1937	$\beta^+ =100$	
<sup>63</sup> Zn <sup>i</sup>	-56723	6	5490 6 RQ	3/2 <sup>-</sup> T=5/2	09			
<sup>63</sup> Ga	-56547.1	1.3	32.4 s 0.5	3/2 <sup>-</sup> *	09 12Pr11 J	1965	$\beta^+ =100$	
<sup>63</sup> Ge	-46920	40	153.6 ms 1.1	3/2 <sup>-</sup> #	09 20Gi02 T	1991	$\beta^+ =100; \beta^+ p ?$	
<sup>63</sup> As	-33500#	200#	<43ns	3/2 <sup>-</sup> #	09 05St29 I		p ?	
<sup>63</sup> Se	-16850#	500#	13.2 ms 3.9	3/2 <sup>-</sup> #	20Gi02 TD	2016	$\beta^+ =100; \beta^+ p =89 11; 2p <0.5$	
* <sup>63</sup> V	T : average 14Su07=20(1) 11Da08=19.2(2.4) 03So02=17(3)						**	
* <sup>63</sup> Cr	T : other 11Da08=128(8)						**	
* <sup>63</sup> Mn	D : $\beta^- n$ observed by 99Ha05, but not quantified						**	
* <sup>63</sup> Co	T : average 94It.A=26.41(0.27) 72Jo08=27.5(0.3) 69Wa15=26(1)						**	
* <sup>63</sup> Ni	J : 17Dy01=1/2						**	
* <sup>63</sup> Cu	J : also 20De21,10Vi07,11Ko36,10Co01=3/2						**	
* <sup>63</sup> Zn	J : also 17Wr01=3/2						**	
* <sup>63</sup> Ge	T : from 20Gi02, supersedes 19Ru.A=153.3(0.6) (same collaboration);						**	
* <sup>63</sup> Ge	T : others: 17Ku12=156(11) 14Ro14=149(4) 02Lo13=150(9) 93Wi03=95(+23-20)						**	
* <sup>63</sup> As	D : most likely p-unstable from estimated Sp=-950#(240#) keV						**	
* <sup>63</sup> Se	T : other 16Go26 (same as 20Gi02)						**	
* <sup>63</sup> Se	D : 2p not observed in 20Gi02 and 16Go26; limit from 16Go26 based on the						**	
* <sup>63</sup> Se	D : assumption that one event is not $\beta^+ p$						**	
<sup>64</sup> Ti	-1480#	600#	5# ms >620ns	0 <sup>+</sup>	13 13Ta14 I	2013	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$	
<sup>64</sup> V	-16320#	400#	15 ms 2	(1,2)	14	1997	$\beta^- =100; \beta^- n ?; \beta^- 2n ?$	
<sup>64</sup> V <sup>m</sup>	-16240#	400#	81.0 0.7	< 1 $\mu$ s	14	2014	IT $\approx$ 100	
<sup>64</sup> Cr	-33640	300	43 ms 1	0 <sup>+</sup>	14	1992	$\beta^- =100; \beta^- n ?$	
<sup>64</sup> Mn	-42989	4	88.8 ms 2.4	1 <sup>+</sup> *	07 12Pa39 D	1985	$\beta^- =100; \beta^- n =2.7 6$	
<sup>64</sup> Mn <sup>m</sup>	-42815	4	174.1 0.5	439 $\mu$ s 31	(4 <sup>+</sup> )*	07 10Da06 E	1998	IT=100
<sup>64</sup> Fe	-54970	5	2.0 s 0.2	0 <sup>+</sup>	07	1980	$\beta^- =100$	
<sup>64</sup> Co	-59792	20	300 ms 30	1 <sup>+</sup>	07	1969	$\beta^- =100$	
<sup>64</sup> Co <sup>m</sup>	-59686	4	107 20 MD	300# ms	5 <sup>+</sup> #	08B105 E	2008	$\beta^- ?; IT ?$
<sup>64</sup> Ni	-67099.0	0.5	STABLE	0 <sup>+</sup>	07	1935	IS=0.9256 19	
<sup>64</sup> Cu	-65424.4	0.4	12.7004 h 0.0013	1 <sup>+</sup> *	07 FGK204 T	1936	$\beta^+ =61.52 26; \beta^- =38.48 26$	
<sup>64</sup> Cu <sup>i</sup>	-58598	6	6826 6	0 <sup>+</sup> frg. T=4	07 71Be29 E			
<sup>64</sup> Zn	-66004.0	0.6	STABLE	>60Py	0 <sup>+</sup>	07 03Ki08 T	1922	IS=49.17 75; 2 $\beta^+ ?$
<sup>64</sup> Ga	-58832.8	1.4	2.627 m 0.012	0(+ #)	07	1953	$\beta^+ =100$	
<sup>64</sup> Ga <sup>m</sup>	-58790.0	1.4	42.85 0.08	21.9 $\mu$ s 0.7	(2 <sup>+</sup> )	07	1999	IT=100
<sup>64</sup> Ga <sup>i</sup>	-56925.8	2.5	1907.0 2.2 RQ	(0 <sup>+</sup> ) T=2	07			
<sup>64</sup> Ge	-54316	4	63.7 s 2.5	0 <sup>+</sup>	07	1972	$\beta^+ =100$	
<sup>64</sup> As	-39530#	200#	69.0 ms 1.4	0 <sup>+</sup> #	07 20Gi02 T	1995	$\beta^+ =100; \beta^+ p ?$	
<sup>64</sup> Se	-26860#	500#	22.6 ms 0.2	0 <sup>+</sup>	07 19Ru.A T	2005	$\beta^+ ?; \beta^+ p ?$	
* <sup>64</sup> Mn	T : average 11Da08=90(9) 02So.A=91(4) 99So20=85(5) 99Ha05=89(4)						**	
* <sup>64</sup> Mn	J : 15He28=1						**	
* <sup>64</sup> Mn	D : % $\beta^- n$ other 00HaZL=33(2)						**	
* <sup>64</sup> Mn <sup>m</sup>	J : 15He28=(4)						**	
* <sup>64</sup> Mn <sup>m</sup>	T : average 11Li50=400(40) 05Ga.B=500(50)						**	
* <sup>64</sup> Cu	J : 20De21,10Vi07=1						**	
* <sup>64</sup> Cu	D : from 12Be24						**	
* <sup>64</sup> Cu <sup>i</sup>	E : strongest fragment (xs=100); other 16 keV lower (xs=37)						**	
* <sup>64</sup> Zn	T : for 2nu- $\epsilon\epsilon$						**	

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>64</sup> As	T : from 20Gi02, supersedes 19Ru.A=63.4(1.2) (same collaboration);								**
* <sup>64</sup> As	T : others 14Ro14=72(6) 02Lo13=18(+43-7)								**
<sup>65</sup> Ti	5210#	700#			1# ms	1/2 <sup>-</sup> #			$\beta^- ?; \beta^- n ?; \beta^- 2n ?$
<sup>65</sup> V	-12110#	500#			14# ms >620ns	5/2 <sup>-</sup> #	10 09Ta24 I	2009	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$
<sup>65</sup> Cr	-28310#	200#			27.5 ms 2.1	1/2 <sup>-</sup> #	10 11Da08 T	1997	$\beta^- =100; \beta^- n ?; \beta^- 2n ?$
<sup>65</sup> Mn	-40967	4			91.9 ms 0.7	(5/2 <sup>-</sup> )	10 13OI06 TJD	1985	$\beta^- =100; \beta^- n =7.9 12$
<sup>65</sup> Fe	-51218	5			805 ms 10	(1/2 <sup>-</sup> )	10 13OI06 TD	1980	$\beta^- =100; \beta^- n ?$
<sup>65</sup> Fe <sup>m</sup>	-50824	5	393.7	0.2	1.12 s 0.15	(9/2 <sup>+</sup> )	10 13OI06 E	2008	$\beta^- ?$
<sup>65</sup> Fe <sup>n</sup>	-50820	5	397.6	0.2	418 ns 12	(5/2 <sup>+</sup> )	10 13OI06 EJ	1998	IT=100
<sup>65</sup> Co	-59185.2	2.1			1.16 s 0.03	(7/2 <sup>-</sup> )	10	1978	$\beta^- =100$
<sup>65</sup> Ni	-65125.8	0.5			2.5175 h 0.0005	5/2 <sup>-</sup>	10	1946	$\beta^- =100$
<sup>65</sup> Ni <sup>m</sup>	-65062.4	0.5	63.37	0.05	69 $\mu$ s 3	1/2 <sup>-</sup>	10	1978	IT=100
<sup>65</sup> Cu	-67263.7	0.6			STABLE	3/2 <sup>-</sup> *	10 10Vi07 J	1923	IS=30.85 15
<sup>65</sup> Zn	-65912.0	0.6			243.94 d 0.04	5/2 <sup>-</sup> *	10 FGK204 T	1939	$\beta^+ =100$
<sup>65</sup> Zn <sup>m</sup>	-65858.1	0.6	53.928	0.010	1.6 $\mu$ s 0.6	1/2 <sup>-</sup>	10 FGK149 J	1960	IT=100
<sup>65</sup> Ga	-62657.5	0.8			15.133 m 0.028	3/2 <sup>-</sup> *	10 19Gy04 T	1938	$\beta^+ =100$
<sup>65</sup> Ge	-56478.2	2.2			30.9 s 0.5	3/2 <sup>-</sup>	10	1972	$\beta^+ =100; \beta^+ p =0.011 3$
<sup>65</sup> As	-46940	80			130.3 ms 0.6	3/2 <sup>-</sup> #	10 20Gi02 T	1991	$\beta^+ =100; \beta^+ p ?$
<sup>65</sup> As <sup>i</sup>	-43452	11	3490	90	p	(3/2 <sup>-</sup> ) T=3/2	10 11Ro47 J	1993	p=100
<sup>65</sup> Se	-33020#	300#			34.2 ms 0.7	3/2 <sup>-</sup> #	10 20Gi02 T	1993	$\beta^+ =100; \beta^+ p =87 13$
<sup>65</sup> Br	-16490#	500#			<410ns	5/2 <sup>-</sup> #	16B105 I		p ?
* <sup>65</sup> Cr	T : average 11Da08=28(3) 03So21=27(3)								**
* <sup>65</sup> Mn	T : average 13OI06=91.9(0.9) 03So21=92(1); other (recent) 11Da08=84(8),								**
* <sup>65</sup> Mn	T : outweighed (not used)								**
* <sup>65</sup> Mn	D : other $\beta^- n$ observed by 99Ha05, but not quantified								**
* <sup>65</sup> Fe	T : 19OI02=805(10). others 09Pa16=810(50) 99So20=1320(280)								**
* <sup>65</sup> Fe	T : 95Am.A=760(50) supersedes 94Cz02=450(150)								**
* <sup>65</sup> Fe <sup>n</sup>	E : other 10Da06=396.8, uncertainty not given, 98Gr14=364(3)								**
* <sup>65</sup> Fe <sup>n</sup>	T : average 18St18=409(+29-27) 10Da06=420(13)								**
* <sup>65</sup> Cu	J : 20De21, 10Vi07, 10Co01=3/2								**
* <sup>65</sup> Zn	J : also 17Wr01=5/2								**
* <sup>65</sup> Zn <sup>m</sup>	J : E2 to 5/2-								**
* <sup>65</sup> Ga	T : from 19Gy04=15.133 (0.016 stat) (0.023 syst); other 57Da07=15.2(0.2)								**
* <sup>65</sup> Ga	J : 17Fa09=3/2								**
* <sup>65</sup> As	T : others (outweighed) 14Ro14=126(7) 02Lo13=126(16) 95Mo26=190(11)								**
* <sup>65</sup> As <sup>i</sup>	J : IAS studied in 93Ba12 and 11Ro47								**
* <sup>65</sup> Se	T : other 11Ro47=33(4)								**
* <sup>65</sup> Se	D : % $\beta^+ p$ symmetrized from 11Ro47=88(+12-13)								**
<sup>66</sup> V	-6300#	500#			10# ms >620ns		10 09Ta24 I	2009	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$
<sup>66</sup> Cr	-25140#	300#			23.8 ms 1.8	0 <sup>+</sup>	15 11Li50 T	1997	$\beta^- =100; \beta^- n ?; \beta^- 2n ?$
<sup>66</sup> Mn	-36750	11			63.8 ms 0.9	(1 <sup>+</sup> )	10 18St18 TDJ	1992	$\beta^- =100; \beta^- n =7.4 14; \beta^- 2n ?$
<sup>66</sup> Mn <sup>m</sup>	-36286	11	464.5	0.4	780 $\mu$ s 40	(5 <sup>-</sup> )	11Li50 ETJ	2005	IT $\approx$ 100; $\beta^- ?$
<sup>66</sup> Fe	-50068	4			467 ms 29	0 <sup>+</sup>	10 18St18 T	1985	$\beta^- =100; \beta^- n ?$
<sup>66</sup> Co	-56409	14			194 ms 17	(1 <sup>+</sup> )	10 12Li02 J	1985	$\beta^- =100; \beta^- n ?$
<sup>66</sup> Co <sup>m</sup>	-56234	14	175.1	0.3	824 ns 22	(3 <sup>+</sup> )	10 12Li02 EJ	1998	IT=100
<sup>66</sup> Co <sup>n</sup>	-55767	15	642	5	> 100 $\mu$ s	(8 <sup>-</sup> )	10 98Gr14 E	1998	IT=100
<sup>66</sup> Ni	-66006.3	1.4			54.6 h 0.3	0 <sup>+</sup>	10	1948	$\beta^- =100$
<sup>66</sup> Cu	-66258.3	0.6			5.120 m 0.014	1 <sup>+</sup> *	10 20De21 J	1937	$\beta^- =100$
<sup>66</sup> Cu <sup>m</sup>	-65104.1	1.5	1154.2	1.4	600 ns 17	(6 <sup>-</sup> )	10 11Lo01 T	1972	IT=100
<sup>66</sup> Zn	-68899.2	0.7			STABLE	0 <sup>+</sup>	10	1922	IS=27.73 98
<sup>66</sup> Ga	-63723.7	1.1			9.304 h 0.008	0 <sup>+</sup> *	10 10Se16 T	1937	$\beta^+ =100$
<sup>66</sup> Ga <sup>i</sup>	-59874	6	3850	6	RQ	0 <sup>+</sup> T=3			*
<sup>66</sup> Ge	-61607.0	2.4			2.26 h 0.05	0 <sup>+</sup>	10	1950	$\beta^+ =100$
<sup>66</sup> As	-52025	6			95.77 ms 0.23	0 <sup>+</sup> T=1	10 MMC156J	1978	$\beta^+ =100$
<sup>66</sup> As <sup>m</sup>	-50668	6	1356.63	0.17	1.14 $\mu$ s 0.04	5 <sup>+</sup>	10 13Ru10 TJ	1995	IT=100
<sup>66</sup> As <sup>n</sup>	-49001	6	3023.8	0.3	7.98 $\mu$ s 0.26	9 <sup>+</sup>	10 13Ru10 TJ	1998	IT=100
<sup>66</sup> Se	-41660#	200#			54 ms 4	0 <sup>+</sup>	10 14Ro14 T	1993	$\beta^+ =100; \beta^+ p ?$
<sup>66</sup> Br	-23570#	400#			<410ns	0 <sup>+</sup> #	16B105 I		p ?
* <sup>66</sup> Cr	T : average 11Li50=24(2) 11Da08=23(4); other 05Ga01=10(6), outweighed								**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>66</sup> Mn	J : 11Li50=(1+) due to large ground-state feeding from <sup>66</sup> Cr (J=0+);						**
* <sup>66</sup> Mn	J : large direct $\beta^-$ feeding to <sup>66</sup> Fe gs (J=0+) in 18St18						**
* <sup>66</sup> Mn	T : average 18St18=64.1(1.1) 17Ol08=70(15) 13Li04=60(3) 03So21=64(2)						**
* <sup>66</sup> Mn	T : 99Ha05=66(4); other 11Pa.A=64.2(0.8) is superseded by 18St18						**
* <sup>66</sup> Mn	D : % $\beta^-$ -n symmetrized from 18St18=7.3(+1.4-1.3)						**
* <sup>66</sup> Mn <sup>m</sup>	T : other 05Ga.B=750(250)						**
* <sup>66</sup> Fe	T : average 18St18=485(+39-34) 99Le67=440(60) 98Am04=440(60)						**
* <sup>66</sup> Co	J : also large direct $\beta^-$ feeding to <sup>66</sup> Ni gs (J=0+) in 18St18						**
* <sup>66</sup> Co <sup>m</sup>	T : symmetrized 18St18=823(+22-21)						**
* <sup>66</sup> Cu	J : 20De21,10Vi07=1						**
* <sup>66</sup> Cu <sup>m</sup>	T : average 11Lo01=601(30) 72Bl16=600(20)						**
* <sup>66</sup> Ga	T : other 12Gy01=9.312(0.032) not used; Ensdf2010=9.49(0.03)						**
* <sup>66</sup> As	T : average 88Bu12=95.77(0.28) 78Al23=95.78(0.39); other (recent)						**
* <sup>66</sup> As	T : 14Ro14=93(4) 02Lo13=97(2) (outweighed)						**
* <sup>66</sup> As	J : super-allowed $\beta^+$ -decay emitter; see also 98Gr12						**
* <sup>66</sup> As <sup>m</sup>	T : average 13Ru10=1.15(0.04) 01Gr07=1.1(0.1)						**
* <sup>66</sup> As <sup>n</sup>	T : average 13Ru10=7.9(0.3) 01Gr07=8.2(0.5)						**
<sup>67</sup> V	-1740#	600#		8# ms >620ns	5/2 <sup>-</sup> #	13 13Ta14 I 2013	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$
<sup>67</sup> Cr	-19270#	400#		11# ms >300ns	1/2 <sup>-</sup> #	05 97Be70 I 1997	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$
<sup>67</sup> Mn	-33580#	200#		46.7 ms 2.3	5/2 <sup>-</sup> #	05 11Da08 TD 1997	$\beta^- =100; \beta^- n =10.5; \beta^- 2n ?$
<sup>67</sup> Fe	-45708	4		394 $\mu$ s 9	(1/2 <sup>-</sup> )	05 02So.A TD 1985	$\beta^- =100; \beta^- n ?$
<sup>67</sup> Fe <sup>m</sup>	-45305	10	403 9	64 $\mu$ s 17	(5/2 <sup>+</sup> , 7/2 <sup>+</sup> )	05 11Da08 EJ 1998	IT=100
<sup>67</sup> Fe <sup>n</sup>	-45260#	100#	450# 100#	75 $\mu$ s 21	(9/2 <sup>+</sup> )	08B105 TJ 2008	IT=100
<sup>67</sup> Co	-55322	6		329 ms 28	(7/2 <sup>-</sup> )	05 08Pa33 TJ 1985	$\beta^- =100; \beta^- n ?$
<sup>67</sup> Co <sup>m</sup>	-54830	6	491.55 0.11	496 ms 33	(1/2 <sup>-</sup> )	09Pa16 E 2008	IT>80; $\beta^- n ?$
<sup>67</sup> Ni	-63742.7	2.9		21 s 1	1/2 <sup>-</sup>	05 00Ri14 J 1978	$\beta^- =100$
<sup>67</sup> Ni <sup>m</sup>	-62736.1	2.9	1006.6 0.2	13.34 $\mu$ s 0.19	9/2 <sup>+</sup>	05 14Di08 ETJ 1998	IT=100
<sup>67</sup> Cu	-67319.6	0.9		61.83 h 0.12	3/2 <sup>-</sup> *	05 20De21 J 1948	$\beta^- =100$
<sup>67</sup> Zn	-67880.4	0.8		STABLE	5/2 <sup>-</sup> *	05 1928	IS=4.04 16
<sup>67</sup> Zn <sup>m</sup>	-67787.1	0.8	93.312 0.005	9.15 $\mu$ s 0.07	1/2 <sup>-</sup>	05 15Ch57 T 1953	IT=100
<sup>67</sup> Zn <sup>n</sup>	-67275.9	0.8	604.48 0.05	333 ns 14	9/2 <sup>+</sup>	05 1973	IT=100
<sup>67</sup> Ga	-66879.2	1.2		3.2617 d 0.0004	3/2 <sup>-</sup> *	05 FGK204 T 1938	$\epsilon =100$
<sup>67</sup> Ge	-62674	4		18.9 m 0.3	1/2 <sup>-</sup>	05 1950	$\beta^+ =100$
<sup>67</sup> Ge <sup>m</sup>	-62656	4	18.20 0.05	13.7 $\mu$ s 0.9	5/2 <sup>-</sup>	05 1978	IT=100
<sup>67</sup> Ge <sup>n</sup>	-61922	4	751.70 0.06	109.1 ns 3.8	9/2 <sup>+</sup>	05 00Ch07 T 1973	IT=100
<sup>67</sup> As	-56587.2	0.4		42.5 s 1.2	(5/2 <sup>-</sup> )	05 1980	$\beta^+ =100$
<sup>67</sup> Se	-46580	70		133 ms 4	5/2 <sup>-</sup> #	05 14Ro14 T 1991	$\beta^+ =100; \beta^+ p =0.5 1$
<sup>67</sup> Br	-32530#	300#			1/2 <sup>-</sup> #		p ?
<sup>67</sup> Kr	-15550#	420#		7.4 ms 2.9	3/2 <sup>-</sup> #	20Gi02 TD 2016	2p=37 14; $\beta^+ ?$
* <sup>67</sup> Mn	T : average 11Da08=51(4) 03So21=47(4) 99Ha05=42(4)						**
* <sup>67</sup> Fe	T : others (recent) 11Da08=304(81) 08Pa33=416(29), outweighed (not used)						**
* <sup>67</sup> Fe <sup>m</sup>	T : average 03Sa02=75(21) 98Gr14=43(30), same authors, different experiment						**
* <sup>67</sup> Fe <sup>n</sup>	E : less than 30 keV above 387.7-keV level						**
* <sup>67</sup> Co <sup>m</sup>	D : %IT from 08Pa33						**
* <sup>67</sup> Ni <sup>m</sup>	T : average 14Di08=13.7(0.6) 98Gr14=13.3(0.2); other 02Ge16=13(1)						**
* <sup>67</sup> Cu	J : 20De21,10Vi07=3/2						**
* <sup>67</sup> Zn	J : also 17Wr01,16Ya02=5/2						**
* <sup>67</sup> Zn <sup>m</sup>	T : average 15Ch57=9.37(0.04) 98At04=9.34(0.20) 96Hw03=9.01(0.03)						**
* <sup>67</sup> Zn <sup>n</sup>	T : 75Ro25=9.1(0.4) 73Le18=9.20(0.07) 72Le37=9.15(0.05);						**
* <sup>67</sup> Zn <sup>m</sup>	T : Birge ratio=3.27						**
* <sup>67</sup> Ga	J : other 17Fa09=3/2						**
* <sup>67</sup> Ge <sup>n</sup>	T : average 00Ch07=101(3) 79Al04=110.9(1.4); Birge ratio=2.99						**
* <sup>67</sup> Se	D : % $\beta^+ p$ from 95Bl23						**
* <sup>67</sup> Kr	T : other 16Go26=7.4(3.0) (same as 20Gi02)						**
<sup>68</sup> Cr	-15690#	500#		10# ms >620ns	0 <sup>+</sup>	12 09Ta24 I 2009	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$
<sup>68</sup> Mn	-28920#	300#		33.7 ms 1.5	(3)	12 15Be32 TD 1995	$\beta^- =100; \beta^- n =18 10; \beta^- 2n ?$
<sup>68</sup> Fe	-43900#	190#		188 ms 4	0 <sup>+</sup>	12 1985	$\beta^- =100; \beta^- n >0$
<sup>68</sup> Co	-51643	4	*	200 ms 20	(7 <sup>-</sup> )	12 1985	$\beta^- =100; \beta^- n ?$
<sup>68</sup> Co <sup>m</sup>	-51490#	150#	150# 150#	1.6 s 0.3	(2 <sup>-</sup> )	12 15Fl01 JD 1998	$\beta^- =100; \beta^- n >2.6$

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{68}\text{Co}^n$	-51450# 150#	195# 150#	101 ns 10	(1)	12 10Da06 T	2010	IT=100 *
$^{68}\text{Ni}$	-63463.8 3.0		29 s 2	$0^+$	12	1977	$\beta^-$ =100
$^{68}\text{Ni}^m$	-61860 3	1603.51 0.28	270 ns 5	$0^+$	12 15Fi01 E	1984	IT=100 *
$^{68}\text{Ni}^n$	-60615 3	2849.1 0.3	850 $\mu\text{s}$ 30	$5^-$	12 15Wi02 T	1995	IT=100 *
$^{68}\text{Cu}$	-65567.0 1.6		30.9 s 0.6	$1^+*$	12 20De21 J	1953	$\beta^-$ =100 *
$^{68}\text{Cu}^m$	-64845.7 1.6	721.26 0.08	3.75 m 0.05	$6^-*$	12 20De21 J	1969	IT=86 2; $\beta^-$ =14 2 *
$^{68}\text{Zn}$	-70007.2 0.8		STABLE	$0^+$	12	1922	IS=18.45 63
$^{68}\text{Ga}$	-67086.1 1.4		67.842 m 0.016	$1^+*$	12 FGK204 T	1937	$\beta^+$ =100
$^{68}\text{Ge}$	-66978.8 1.9		271.05 d 0.08	$0^+$	12 18Be03 T	1948	$\epsilon$ =100 *
$^{68}\text{As}$	-58894.5 1.8		151.6 s 0.8	$3^+$	12	1971	$\beta^+$ =100
$^{68}\text{As}^m$	-58469.4 1.8	425.1 0.2	111 ns 0.7	$1^+$	12	1994	IT=100 *
$^{68}\text{Se}$	-54189.4 0.5		35.5 s 0.7	$0^+$	12	1990	$\beta^+$ =100
$^{68}\text{Br}$	-38790# 260#		$\sim$ 40ns	$3^+*$	12 19Wi08 T	1995	p ?
$^{68}\text{Kr}$	-25630# 500#		21.6 ms 3.3	$0^+$	20Gi02 TD	2016	$\beta^+$ = ?; $\beta^+$ p=90 11;p ? *
$^{*68}\text{Mn}$	T : average 15Be32=38.3(3.6) and 35.2(2.0) 11Da08=29(4) 03So21=28(8)						
$^{*68}\text{Mn}$	T : 99Ha05=28(4).						
$^{*68}\text{Mn}$	D : $\beta^-$ n observed by 99Ha05, but not quantified						
$^{*68}\text{Mn}$	J : direct $\beta^-$ feeding to 2+ and 4+ in 15Be32 (incomplete decay scheme)						
$^{*68}\text{Co}^n$	J : strong feeding in 68Fe (J=0+) $\beta^-$ decay and possible gamma-ray decay						
$^{*68}\text{Co}^n$	J : to 2- in 12Li02						
$^{*68}\text{Ni}^m$	E : average 15Fi01=1603.6(0.8) 13Re18=1603.5(0.3) from g-ray differences						
$^{*68}\text{Ni}^n$	T : average 15Wi02=840(40) 95Br10=860(50)						
$^{*68}\text{Cu}$	J : 20De21,10Vi07=1						
$^{*68}\text{Cu}^m$	J : 20De21,10Vi07=6						
$^{*68}\text{Ge}$	T : average 18Be03=271.14(0.15,NaI), 271.07(0.12,IC) 94Sc44=270.99(0.19)						
$^{*68}\text{Ge}$	T : 81Wa26=270.82(0.27)						
$^{*68}\text{As}^m$	T : symmetrized from 94Ba50=107(+23-16)						
$^{*68}\text{Kr}$	D : % $\beta^+$ p symmetrized from 20Gi02=89(+11-10)						
$^{69}\text{Cr}$	-9630# 500#		6# ms >620ns	$7/2^+*$	14 13Ta14 I	2013	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?
$^{69}\text{Mn}$	-25360# 400#		22.1 ms 1.6	$5/2^-*$	14 15Be32 TD	1995	$\beta^-$ =100; $\beta^-$ n=40 20; $\beta^-$ 2n ? *
$^{69}\text{Fe}$	-39200# 200#		162 ms 7	$1/2^-*$	14 15Li33 T	1992	$\beta^-$ =100; $\beta^-$ n ?; $\beta^-$ 2n ?
$^{69}\text{Co}$	-50390 90		180 ms 20	$(7/2^-)$	14 15Li33 T	1985	$\beta^-$ =100; $\beta^-$ n ?
$^{69}\text{Co}^m$	-50213 13	170 90 MD*	750 ms 250	$1/2^-*$	15Li33 TD	2015	$\beta^-$ =100
$^{69}\text{Ni}$	-59979 4		11.4 s 0.3	$(9/2^+)$	14	1984	$\beta^-$ =100
$^{69}\text{Ni}^m$	-59658 4	321 2	3.5 s 0.4	$(1/2^-)$	14 98Gr14 E	1998	$\beta^- \approx 100$ ;IT<0.01 *
$^{69}\text{Ni}^n$	-57279 4	2700.0 1.0	439 ns 3	$(17/2^-)$	14	1998	IT=100
$^{69}\text{Cu}$	-65736.2 1.4		2.85 m 0.15	$3/2^-*$	14 20De21 J	1966	$\beta^-$ =100 *
$^{69}\text{Cu}^m$	-62994.2 1.6	2742.0 0.7	357 ns 2	$(13/2^+)$	14 16Ku11 T	1997	IT=100 *
$^{69}\text{Zn}$	-68417.9 0.8		56.4 m 0.9	$1/2^-*$	14 17Wr01 J	1937	$\beta^-$ =100
$^{69}\text{Zn}^m$	-67979.3 0.8	438.636 0.018	13.747 h 0.011	$9/2^+*$	14 17Wr01 J	1970	IT=99.967 3; $\beta^-$ =0.033 3 *
$^{69}\text{Ga}$	-69327.8 1.2		STABLE	$3/2^-*$	14	1923	IS=60.108 50 *
$^{69}\text{Ge}$	-67100.7 1.3		39.05 h 0.10	$5/2^-*$	14	1938	$\beta^+$ =100
$^{69}\text{Ge}^m$	-67013.9 1.3	86.76 0.02	5.1 $\mu\text{s}$ 0.2	$1/2^-*$	14	1978	IT=100
$^{69}\text{Ge}^n$	-66702.8 1.3	397.94 0.02	2.81 $\mu\text{s}$ 0.05	$9/2^+$	14	1978	IT=100
$^{69}\text{As}$	-63110 30		15.2 m 0.2	$5/2^-*$	14	1955	$\beta^+$ =100
$^{69}\text{Se}$	-56434.7 1.5		27.4 s 0.2	$1/2^-*$	14	1974	$\beta^+$ =100; $\beta^+$ p=0.052 8
$^{69}\text{Se}^m$	-56395.9 1.5	38.85 0.22	2.0 $\mu\text{s}$ 0.2	$5/2^-*$	14	1988	IT=100
$^{69}\text{Se}^n$	-55860.7 1.6	574.0 0.4	955 ns 16	$9/2^+$	14 00Ch07 T	1988	IT=100 *
$^{69}\text{Br}$	-46260 40		< 24 ns	$(5/2^-)$	15	1988	p=100
$^{69}\text{Br}^m$	-46220# 110#	40# 100#		$5/2^-*$	Mirror I		
$^{69}\text{Br}^n$	-45690# 110#	570# 100#		$9/2^+*$	Mirror I		
$^{69}\text{Br}^i$	-42771 19	3490 50 p		$(5/2^-)$ T=3/2	14 11Ro47 I	2011	p=100
$^{69}\text{Kr}$	-32140# 300#		27.9 ms 0.8	$(5/2^-)$	15 02Gi02 DT	1995	$\beta^+$ =100; $\beta^+$ p=94 5 *
$^{*69}\text{Mn}$	T : average 15Be32=24.1(2.6) 25.8(2.8) 11Da08=18(4) 99Ha05=14(4)						
$^{*69}\text{Ni}^m$	E : from 98Gr14; E(9/2+) in $^{73}\text{Ge}$ =67 keV and $^{71}\text{Zn}$ =156 keV						
$^{*69}\text{Ni}^m$	E : isotones exhibits a large variation						
$^{*69}\text{Cu}$	J : 20De21,10Vi07=3/2						
$^{*69}\text{Cu}^m$	T : average 16Ku11=351(14) 12Di03=360(20) 02Ge16=357(2)						
$^{*69}\text{Cu}^m$	T : 98Gr14=360(50) 97Is13=360(30)						
$^{*69}\text{Zn}^m$	T : average 17Kr01=13.742(0.014) 77He20=13.756(0.018)						
$^{*69}\text{Ga}$	J : other 17Fa09=3/2						

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>69</sup> Se <sup>n</sup>	T : average 00Ch07=950(21) 95Po01=960(23)						**
* <sup>69</sup> Kr	T : average 20Gi02=27.8(1.6) 14De41=28(1) 11Ro47=27(3); other 97Xu01=32(10)						**
* <sup>69</sup> Kr	D : %β <sup>+</sup> p average 02Gi02=93(+7-6) 11Ro47=99(+1-11); other						**
* <sup>69</sup> Kr	D : 14De41=52.5(6.5) + 2.4(0.5), conflicting						**
<sup>70</sup> Cr	-5640# 600#		6# ms >620ns	0 <sup>+</sup>	16 13Ta14 I	2013	β <sup>-</sup> ?; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>70</sup> Mn	-20450# 500#		19.9 ms 1.7	(4,5)	16 15Be32 TD	2009	β <sup>-</sup> =100; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>70</sup> Fe	-36890# 300#		61.4 ms 0.7	0 <sup>+</sup>	16 17Mo02 T	1997	β <sup>-</sup> =100; β <sup>-</sup> n ?
<sup>70</sup> Co	-46525 11		* & 508 ms 7	(1 <sup>+</sup> )	16 17Mo02 JT	1998	β <sup>-</sup> =100; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>70</sup> Co <sup>m</sup>	-46330# 200#	200# 200#	* & 112 ms 7	(7 <sup>-</sup> )	16 FGK205 J	1985	β <sup>-</sup> =100; IT ?; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>70</sup> Ni	-59213.9 2.1		6.0 s 0.3	0 <sup>+</sup>	16	1987	β <sup>-</sup> =100
<sup>70</sup> Ni <sup>m</sup>	-56353.0 2.1	2860.91 0.08	232 ns 1	8 <sup>+</sup>	16	1997	IT=100
<sup>70</sup> Cu	-62976.4 1.1		44.5 s 0.2	6 <sup>-*</sup>	16 20De21 J	1971	β <sup>-</sup> =100
<sup>70</sup> Cu <sup>m</sup>	-62875.3 1.1	101.1 0.3	33 s 2	3 <sup>-*</sup>	16 20De21 J	2002	β <sup>-</sup> =52.9; IT=48.9
<sup>70</sup> Cu <sup>n</sup>	-62733.8 1.2	242.6 0.5	6.6 s 0.2	1 <sup>+</sup> *	16 20De21 J	1971	β <sup>-</sup> =93.2.9; IT=6.8.9
<sup>70</sup> Zn	-69564.7 1.9		STABLE >3.8Ey	0 <sup>+</sup>	16	1922	IS=0.61 10; 2β <sup>-</sup> ?
<sup>70</sup> Ga	-68910.2 1.2		21.14 m 0.05	1 <sup>+</sup> *	16	1937	β <sup>-</sup> =99.59.5; ε=0.41.5
<sup>70</sup> Ge	-70562.0 0.8		STABLE	0 <sup>+</sup>	16	1923	IS=20.52.19
<sup>70</sup> As	-64334.0 1.4		52.6 m 0.3	4 <sup>+</sup> *	16 76He24 J	1950	β <sup>+</sup> =100
<sup>70</sup> As <sup>m</sup>	-64302.0 1.4	32.046 0.023	96 μs 3	2 <sup>+</sup>	16	1979	IT=100
<sup>70</sup> Se	-61929.9 1.6		41.1 m 0.3	0 <sup>+</sup>	16	1950	β <sup>+</sup> =100
<sup>70</sup> Br	-51426 15		78.8 ms 0.3	0 <sup>+</sup> T=1	16 17Mo18 T	1978	β <sup>+</sup> =100; β <sup>+</sup> p ?
<sup>70</sup> Br <sup>m</sup>	-49134 15	2292.3 0.8	2.16 s 0.05	9 <sup>+</sup> T=0	16 17Mo18 T	1981	β <sup>+</sup> =100; β <sup>+</sup> p ?
<sup>70</sup> Kr	-41100# 200#		45.00 ms 0.14	0 <sup>+</sup>	16 20Vi02 T	1995	β <sup>+</sup> =100; β <sup>+</sup> p <1.3
* <sup>70</sup> Fe	T : others (not used): 14XuZZ=66(7) 13Ma87=61(5) 11Da08=71(10) 03So21=94(17)						**
* <sup>70</sup> Co	T : others (not used) 15Pr10=470(50) 00Mu10=500(180)						**
* <sup>70</sup> Cu	J : 20De21, 10Vi07, 16Bi08=6						**
* <sup>70</sup> Zn	T : 2ν-ββ >3.8 Ey Onu-BB >32 Ey in 11Be39; Onu-BB >6.8 Ey in 16Eb03						**
* <sup>70</sup> Ga	J : also 12Pr11=1						**
* <sup>70</sup> Br	T : average 17Mo18=78.42(0.51) 88Bu12=78.54(0.59) 78Al23=80.2(0.8);						**
* <sup>70</sup> Br	T : other (recent) 19Si33=79.7(2.4) (outweighed)						**
* <sup>70</sup> Br <sup>m</sup>	T : symmetrized from 17Mo18=2.157(+0.053-0.049); others (not used):						**
* <sup>70</sup> Br <sup>m</sup>	T : 81Vo04=2.2(0.2) 02Ro25=2.2(0.3) (outweighed)						**
* <sup>70</sup> Kr	T : average 20Vi02=44.99(0.14,stat)(0.06,syst) 45.16(0.68,stat)(0.20,syst);						**
* <sup>70</sup> Kr	T : others (outweighed) 16De29=31(+13-7) 14Ro14=40(6) 02B117=42(31)						**
* <sup>70</sup> Kr	T : 00Oi02=57(21)						**
<sup>71</sup> Mn	-16620# 500#		16# ms >400ns	5/2 <sup>-</sup> #	10 10Oh02 I	2010	β <sup>-</sup> ?; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>71</sup> Fe	-31930# 400#		34.3 ms 2.6	7/2 <sup>+</sup> #	10 13Ma87 T	1997	β <sup>-</sup> =100; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>71</sup> Co	-44370 470		80 ms 3	(7/2 <sup>-</sup> )	10 12Ra10 TJD	1992	β <sup>-</sup> =100; β <sup>-</sup> n=3.1
<sup>71</sup> Ni	-55406.2 2.2		2.56 s 0.03	(9/2 <sup>+</sup> )	10	1987	β <sup>-</sup> =100
<sup>71</sup> Ni <sup>m</sup>	-55406.0 2.3	499 5	2.3 s 0.3	(1/2 <sup>-</sup> )	10	2009	β <sup>-</sup> =100
<sup>71</sup> Cu	-62711.1 1.5		19.4 s 1.4	3/2 <sup>-*</sup>	10	1983	β <sup>-</sup> =100
<sup>71</sup> Cu <sup>m</sup>	-59955.4 1.6	2755.7 0.6	271 ns 13	(19/2 <sup>-</sup> )	10 98Gr14 TJ	1998	IT=100
<sup>71</sup> Zn	-67328.8 2.7		2.40 m 0.05	1/2 <sup>-*</sup>	10 17Wr01 J	1955	β <sup>-</sup> =100
<sup>71</sup> Zn <sup>m</sup>	-67171.1 2.4	157.7 1.3 MD	4.148 h 0.012	9/2 <sup>+</sup> *	10	1958	β <sup>-</sup> ≈ 100; IT ?
<sup>71</sup> Ga	-70139.1 0.8		STABLE	3/2 <sup>-*</sup>	10	1923	IS=39.892.50
<sup>71</sup> Ge	-69906.7 0.8		11.43 d 0.03	1/2 <sup>-*</sup>	10	1941	ε=100
<sup>71</sup> Ge <sup>m</sup>	-69708.3 0.8	198.354 0.014	20.41 ms 0.18	9/2 <sup>+</sup>	10	1959	IT=100
<sup>71</sup> As	-67893 4		65.30 h 0.07	5/2 <sup>-*</sup>	10	1939	β <sup>+</sup> =100
<sup>71</sup> Se	-63146.5 2.8		4.74 m 0.05	(5/2 <sup>-</sup> )	10	1957	β <sup>+</sup> =100
<sup>71</sup> Se <sup>m</sup>	-63097.7 2.8	48.79 0.05	5.6 μs 0.7	(1/2 <sup>-</sup> )	10	1982	IT=100
<sup>71</sup> Se <sup>n</sup>	-62886.0 2.8	260.48 0.10	19.0 μs 0.5	(9/2 <sup>+</sup> )	10	1982	IT=100
<sup>71</sup> Br	-56502 5		21.4 s 0.6	(5/2 <sup>-</sup> )	10	1981	β <sup>+</sup> =100
<sup>71</sup> Kr	-46330 130		98.8 ms 0.3	(5/2 <sup>-</sup> )	10 19Si33 T	1981	β <sup>+</sup> =100; β <sup>+</sup> p=2.1.7 p ?
<sup>71</sup> Rb	-32290# 400#		*	5/2 <sup>-</sup> #			
<sup>71</sup> Rb <sup>m</sup>	-32240# 410#	50# 100#	*	1/2 <sup>-</sup> #	Mirror I		
<sup>71</sup> Rb <sup>n</sup>	-32030# 410#	260# 100#		9/2 <sup>+</sup> #	Mirror I		
* <sup>71</sup> Fe	T : average 14XuZZ=34.7(3.6) 13Ma87=42(6) 11Da08=28(5)						**
* <sup>71</sup> Co	D : %β <sup>-</sup> n from 12Ra10 <2.7(0.9) and 05Ma95 >3(1) of the same group						**
* <sup>71</sup> Co	T : others 19Ly02=86(10) 12Ra10=10RaZY=80(3) 04Sa59=79(5) 03So21=97(2)						**



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>71</sup> Co	T : 98Am04=210(40) 95Am.A=200(50)						**
* <sup>71</sup> Cu	T : average 99Pr10=19(3) 83Ru06=19.5(1.6)						**
* <sup>71</sup> Cu	J : 20De21,10Vi07=3/2						**
* <sup>71</sup> Cu <sup>m</sup>	T : average 98Is11=250(30) 98Gr14=275(14)						**
* <sup>71</sup> Zn	T : average 17Kr01=2.36(0.08) 61Th04=2.45(0.10)						**
* <sup>71</sup> Zn <sup>m</sup>	J : 17Wr01=9/2						**
* <sup>71</sup> Zn <sup>m</sup>	T : average 17Kr01=4.155(0.004) 12Re05=4.127(0.007); Birge ratio=3.47						**
* <sup>71</sup> Zn <sup>m</sup>	D : 156 keV depopulating transition not observed experimentally and						**
* <sup>71</sup> Zn <sup>m</sup>	D : only a limit of %IT<0.05 given in 70Zo01						**
* <sup>71</sup> Ga	J : other 17Fa09=3/2						**
* <sup>71</sup> Kr	T : others 14Ro14=92(9) 97Oi01=100(3) 81Ew01=97(9) 95Bi23=64(+8-5).						**
* <sup>71</sup> Kr	T : Values from 95Bi23 for <sup>67</sup> Se and <sup>71</sup> Kr questioned in 97Oi01						**
* <sup>71</sup> Kr	D : %β <sup>+</sup> p from 97Oi01=2.1 7; other 95Bi23=5.2(0.6) conflicting not trusted						**
<sup>72</sup> Mn	-11170#	600#	12# ms >620ns		13 13Ta14	I 2013	β <sup>-</sup> ?; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>72</sup> Fe	-29250#	500#	17.0 ms 1.0	0 <sup>+</sup>	10 13Ma87	TD 1997	β <sup>-</sup> =100; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>72</sup> Co	-40300#	300#	52.5 ms 0.3	(6 <sup>-</sup> , 7 <sup>-</sup> )	10 16Mo07	T 1992	β <sup>-</sup> =100; β <sup>-</sup> n >4; β <sup>-</sup> 2n ?
<sup>72</sup> Co <sup>m</sup>	-40100#	360#	47.8 ms 0.5	(0 <sup>+</sup> , 1 <sup>+</sup> )	16Mo07	TJ 2016	β <sup>-</sup> =100
<sup>72</sup> Ni	-54226.1	2.2	1.57 s 0.05	0 <sup>+</sup>	10	1987	β <sup>-</sup> =100; β <sup>-</sup> n ?
<sup>72</sup> Cu	-59783.0	1.4	6.63 s 0.03	2 <sup>-</sup> *	10 20De21	J 1983	β <sup>-</sup> =100
<sup>72</sup> Cu <sup>m</sup>	-59512.7	1.7	1.76 μs 0.03	(6 <sup>-</sup> )	10	1998	IT=100
<sup>72</sup> Zn	-68145.5	2.1	46.5 h 0.1	0 <sup>+</sup>	10	1951	β <sup>-</sup> =100
<sup>72</sup> Ga	-68588.3	0.8	14.025 h 0.010	3 <sup>-</sup> *	10 12Kr07	T 1939	β <sup>-</sup> =100
<sup>72</sup> Ga <sup>m</sup>	-68468.6	0.8	39.68 ms 0.13	(0 <sup>+</sup> )	10	1968	IT=100
<sup>72</sup> Ge	-72585.91	0.08	STABLE	0 <sup>+</sup>	10	1923	IS=27.45 15
<sup>72</sup> Ge <sup>m</sup>	-71894.48	0.09	444.2 ns 0.8	0 <sup>+</sup>	10	1984	IT=100
<sup>72</sup> As	-68230	4	26.0 h 0.1	2 <sup>-</sup> *	10	1939	β <sup>+</sup> =100
<sup>72</sup> Se	-67868.2	2.0	8.40 d 0.08	0 <sup>+</sup>	10	1948	ε=100
<sup>72</sup> Br	-59061.8	1.0	78.6 s 2.4	1 <sup>+</sup>	10	1970	β <sup>+</sup> =100
<sup>72</sup> Br <sup>m</sup>	-58961.0	1.0	100.76 0.15	(3 <sup>-</sup> )	10	1980	IT≈100; β <sup>+</sup> ?
<sup>72</sup> Kr	-53941	8	17.16 s 0.18	0 <sup>+</sup>	10 03Pi03	T 1973	β <sup>+</sup> =100
<sup>72</sup> Rb	-38330#	500#	103 ns 22	1 <sup>+</sup> #	17Su31	T 2017	p ?
<sup>72</sup> Rb <sup>m</sup>	-38230#	510#	100# 100#	3 <sup>-</sup> #			p ?
* <sup>72</sup> Fe	T : average 14XuZZ=16.9(1.0) 13Ma87=19(4)						**
* <sup>72</sup> Co	T : others 14Xu07=52.8(1.6) 14Ra20=55(4) 05Ma59=59(2) 03Sa40=62(3)						**
* <sup>72</sup> Co	J : β <sup>-</sup> feeding of the 6+ level in <sup>72</sup> Ni and shell model						**
* <sup>72</sup> Co	D : from %β <sup>-</sup> n >6(2) in 05Ma95						**
* <sup>72</sup> Cu	J : 20De21,10Vi07=2						**
* <sup>72</sup> Cu <sup>m</sup>	D : no β <sup>-</sup> decay observed in 05Th.A						**
* <sup>72</sup> Kr	T : average 03Pi03=17.1(0.2) 73Da22=17.4(0.4)						**
* <sup>72</sup> Rb	J : 19Si33=p3/2[321] n1/2[321], K=1+; similarity with the mirror <sup>72</sup> Br						**
<sup>73</sup> Mn	-6700#	600#	12# ms >410ns	5/2 <sup>-</sup> #	19 17Su15	I 2017	β <sup>-</sup> ?
<sup>73</sup> Fe	-23990#	500#	12.9 ms 1.6	7/2 <sup>+</sup> #	19 14XuZZ	T 2010	β <sup>-</sup> =100; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>73</sup> Co	-37970#	300#	42.0 ms 0.8	(7/2 <sup>-</sup> )	19 20Go10	JTD 1995	β <sup>-</sup> =100; β <sup>-</sup> n=6 3; β <sup>-</sup> 2n ?
<sup>73</sup> Ni	-50108.2	2.4	840 ms 30	(9/2 <sup>+</sup> )	19	1987	β <sup>-</sup> =100; β <sup>-</sup> n ?
<sup>73</sup> Cu	-58987.4	1.9	4.20 s 0.12	3/2 <sup>-</sup> *	19 00KoZH	TD 1983	β <sup>-</sup> =100; β <sup>-</sup> n=0.029 6
<sup>73</sup> Zn	-65593.4	1.9	24.5 s 0.2	1/2 <sup>-</sup> *	19 17Ve05	T 1972	β <sup>-</sup> =100
<sup>73</sup> Zn <sup>m</sup>	-65397.9	1.9	13.0 ms 0.2	5/2 <sup>+</sup> *	19 18Ya11	J 1985	IT=100
<sup>73</sup> Ga	-69699.3	1.7	4.86 h 0.03	1/2 <sup>-</sup> *	19 10Ch16	J 1949	β <sup>-</sup> =100
<sup>73</sup> Ga <sup>m</sup>	-69699.2	1.7	< 200 ms	3/2 <sup>-</sup>	19 17Ve05	E 1949	β <sup>-</sup> ?; IT ?
<sup>73</sup> Ge	-71297.53	0.06	STABLE	9/2 <sup>+</sup> *	19 49To09	J 1933	IS=7.76 8
<sup>73</sup> Ge <sup>m</sup>	-71284.25	0.06	13.2845 0.0015	5/2 <sup>+</sup>	19	1975	IT=100
<sup>73</sup> Ge <sup>n</sup>	-71230.80	0.06	66.725 0.009	1/2 <sup>-</sup>	19	1957	IT=100
<sup>73</sup> As	-70953	4	80.30 d 0.06	3/2 <sup>-</sup>	19	1948	ε=100
<sup>73</sup> As <sup>m</sup>	-70525	4	427.902 0.021	9/2 <sup>+</sup>	19	1956	IT=100
<sup>73</sup> Se	-68227	7	7.15 h 0.09	9/2 <sup>+</sup> *	19 88Be39	J 1948	β <sup>+</sup> =100
<sup>73</sup> Se <sup>m</sup>	-68201	7	25.71 0.04	3/2 <sup>-</sup>	19	1960	IT=72.6 3; β <sup>+</sup> =27.4 3
<sup>73</sup> Br	-63646	7	3.4 m 0.2	1/2 <sup>-</sup>	19	1970	β <sup>+</sup> =100
<sup>73</sup> Kr	-56552	7	27.3 s 1.0	(3/2 <sup>-</sup> )	19	1972	β <sup>+</sup> =100; β <sup>+</sup> p=0.25 3
<sup>73</sup> Kr <sup>m</sup>	-56118	7	433.55 0.13	(9/2 <sup>+</sup> )	19	1993	IT=100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{73}\text{Kr}^i$	-53350# 120#	3204# 118#		T=3/2	20Ho17 E		
$^{73}\text{Rb}$	-46010 40		< 81 ns	3/2 <sup>-</sup> #	19 17Su31 T	1996	$\beta^+$ ?; p $\approx$ 100
$^{73}\text{Rb}^m$	-45580# 110#	430# 100#		9/2 <sup>+</sup> #	Mirror I		
$^{73}\text{Rb}^i$	-42809 20	3200 40 p		(5/2 <sup>-</sup> ) T=5/2	19 20Ho06 JD	1993	p=100
$^{73}\text{Sr}$	-31950# 400#		25.3 ms 1.4	(5/2 <sup>-</sup> )	19 20Ho06 TDJ	1993	$\beta^+$ =100; $\beta^+$ p=63 3
$^{*73}\text{Co}$	D : % $\beta^-$ n from 20Go10=6(3), supersedes 12Ra10<22(8) 05Ma95>9(4); other						
$^{*73}\text{Co}$	D : 10Ho12<7.9						
$^{*73}\text{Co}$	T : average 20Go10=43(1), supersedes 12Ra10=42(3), 14Xu07=40.4(1.3),						
$^{*73}\text{Co}$	T : supersedes 14Xu.A=40.5(3.3), 11Da08,04Sa59=41(4) 10Ho12=41(6)						
$^{*73}\text{Cu}$	T : average 00KhZH=4.22(0.15) 98Hu20=4.4(0.3) 83Ru06=3.9(0.3)						
$^{*73}\text{Cu}$	J : 20De21,17De30,10Vi07,09Fi03=3/2						
$^{*73}\text{Zn}$	J : 17Wr01,18Ya11=1/2						
$^{*73}\text{Ga}^m$	E : from <0.3 keV in 17Ve05						
$^{*73}\text{Rb}^i$	J : other 93Ba61=1/2-, T=3/2						
$^{*73}\text{Rb}^i$	E : from 20Ho17						
$^{*73}\text{Sr}$	T : average 20Ho06=23.1(1.4) 19Si33=28(+5-4); others: 20Ho06=23.5(1.8)						
$^{*73}\text{Sr}$	T : 19Si33=24.3(5.3) using a least-squares fit analysis						
$^{74}\text{Fe}$	-20660# 500#		5 ms 5	0 <sup>+</sup>	17 10Oh02 I	2010	$\beta^-$ =100; $\beta^-$ n ?; $\beta^-$ 2n ?
$^{74}\text{Co}$	-33540# 400#		31.3 ms 1.3	7 <sup>-</sup> #	06 10Ho12 TD	1995	$\beta^-$ =100; $\beta^-$ n=18 15; $\beta^-$ 2n ?
$^{74}\text{Ni}$	-48700# 200#		507.7 ms 4.6	0 <sup>+</sup>	06 14Xu07 T	1987	$\beta^-$ =100; $\beta^-$ n ?
$^{74}\text{Cu}$	-56006 6		1.606 s 0.009	2 <sup>-</sup> *	06 00KoZH TD	1987	$\beta^-$ =100; $\beta^-$ n=0.075 16
$^{74}\text{Zn}$	-65756.7 2.5		95.6 s 1.2	0 <sup>+</sup>	06	1972	$\beta^-$ =100
$^{74}\text{Ga}$	-68049.6 3.0		8.12 m 0.12	(3 <sup>-</sup> )*	06 13Ma15 J	1956	$\beta^-$ =100
$^{74}\text{Ga}^m$	-67990 3	59.571 0.014	9.5 s 1.0	(0)( <sup>+</sup> #)	06	1974	IT=75 25; $\beta^-$ ?
$^{74}\text{Ge}$	-73422.451 0.013		STABLE	0 <sup>+</sup>	06	1923	IS=36.52 12
$^{74}\text{As}$	-70860.1 1.7		17.77 d 0.02	2 <sup>-</sup>	06	1938	$\beta^+$ =66 2; $\beta^+$ =34 2
$^{74}\text{Se}$	-72213.210 0.015		STABLE >2.3Ey	0 <sup>+</sup>	06 20Ba08 T	1922	IS=0.86 3; $2\beta^+$ ?
$^{74}\text{Br}$	-65288 6		25.4 m 0.3	(0 <sup>-</sup> )	06	1952	$\beta^+$ =100
$^{74}\text{Br}^m$	-65274 6	13.58 0.21	46 m 2	4 <sup>+</sup> *	06	1953	$\beta^+$ =100
$^{74}\text{Kr}$	-62331.8 2.0		11.50 m 0.11	0 <sup>+</sup>	06	1960	$\beta^+$ =100
$^{74}\text{Rb}$	-51916 3		64.78 ms 0.03	0 <sup>+</sup> * T=1	06 11Ma66 J	1977	$\beta^+$ =100; $\beta^+$ p ?
$^{74}\text{Sr}$	-40830# 100#		27.6 ms 2.6	0 <sup>+</sup>	15 19Si33 T	1995	$\beta^+$ =100; $\beta^+$ p ?
$^{*74}\text{Fe}$	T : symmetrized from 14XuZZ=8.2(+2.6-7.1)						
$^{*74}\text{Co}$	T : average 14Xu07=31.6(1.5) 05Ma95=30(3)						
$^{*74}\text{Co}$	T : others (recent) 11Da08=19(7) 10Ho12=34(+6-9) outweighed (not used)						
$^{*74}\text{Cu}$	T : average 05Va19=1.75(0.06) 00KoZH=1.68(0.03) 91Kr15=1.594(0.010)						
$^{*74}\text{Cu}$	T : 89Wi11=1.59(0.05); others 90Be13=1.51(0.27)						
$^{*74}\text{Cu}$	J : 20De21,17De30,10Vi07,10Fi02=2						
$^{*74}\text{Rb}$	T : average 01Ba12=64.761(0.031) 02Oi02,01Oi04=64.90(0.09); other (recent)						
$^{*74}\text{Rb}$	T : 19Si33=65.1(0.5) (outweighed)						
$^{*74}\text{Sr}$	T : average 19Si33=27.7(2.8) 14He29=27(8)						
$^{75}\text{Fe}$	-14700# 600#		9# ms >620ns	9/2 <sup>+</sup> #	13 13Ta14 I	2013	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?
$^{75}\text{Co}$	-30560# 400#		26.5 ms 1.2	7/2 <sup>-</sup> #	13 14Xu07 T	1995	$\beta^-$ =100; $\beta^-$ n<16; $\beta^-$ 2n ?
$^{75}\text{Ni}$	-44240# 200#		331.6 ms 3.2	9/2 <sup>+</sup> #	13 14Xu07 T	1992	$\beta^-$ =100; $\beta^-$ n=10.0 28
$^{75}\text{Cu}$	-54470.2 0.7		1.224 s 0.003	5/2 <sup>-</sup> *	13 00KoZH D	1985	$\beta^-$ =100; $\beta^-$ n=2.7 4
$^{75}\text{Cu}^m$	-54408.5 0.8	61.7 0.4	310 ns 8	1/2 <sup>-</sup>	13 16Pe14 ET	2010	IT=100
$^{75}\text{Cu}^n$	-54404.0 0.8	66.2 0.4	149 ns 5	3/2 <sup>-</sup>	13 16Pe14 ET	2010	IT=100
$^{75}\text{Zn}$	-62558.9 2.0		10.2 s 0.2	7/2 <sup>+</sup> *	13 17Wr01 J	1974	$\beta^-$ =100
$^{75}\text{Zn}^m$	-62432.0 2.0	126.94 0.09	5# s	1/2 <sup>-</sup> *	13 17Wr01 J	2011	$\beta^-$ ?;IT ?
$^{75}\text{Ga}$	-68460.6 0.7		126 s 2	3/2 <sup>-</sup> *	13	1960	$\beta^-$ =100
$^{75}\text{Ge}$	-71856.97 0.05		82.78 m 0.04	1/2 <sup>-</sup> *	13	1939	$\beta^-$ =100
$^{75}\text{Ge}^m$	-71717.28 0.06	139.69 0.03	47.7 s 0.5	7/2 <sup>+</sup>	13	1952	IT $\approx$ 100; $\beta^-$ =0.030 6
$^{75}\text{Ge}^n$	-71664.78 0.08	192.19 0.06	216 ns 5	5/2 <sup>+</sup>	13	1982	IT=100
$^{75}\text{As}$	-73034.2 0.9		STABLE	3/2 <sup>-</sup> *	13	1920	IS=100
$^{75}\text{As}^m$	-72730.3 0.9	303.9243 0.0008	17.62 ms 0.23	9/2 <sup>+</sup>	13	1957	IT=100
$^{75}\text{Se}$	-72169.49 0.07		119.78 d 0.03	5/2 <sup>+</sup> *	13 FGK209 T	1947	$\epsilon$ =100
$^{75}\text{Br}$	-69107 4		96.7 m 1.3	3/2 <sup>-</sup>	13	1948	$\beta^+$ =100
$^{75}\text{Kr}$	-64324 8		4.60 m 0.07	5/2 <sup>+</sup> *	13 95Ke04 J	1960	$\beta^+$ =100
$^{75}\text{Rb}$	-57218.7 1.2		19.0 s 1.2	3/2 <sup>-</sup> *	13	1975	$\beta^+$ =100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>75</sup> Sr	-46620	220			85.2 ms 2.3	(3/2 <sup>-</sup> )	13 19Si33 T	1991	$\beta^+=100; \beta^+p=5.2$ 9 *	
<sup>75</sup> Y	-31820#	300#			100# $\mu$ s	5/2 <sup>+</sup> #			$\beta^+ ?; \beta^+p ?$ *	
* <sup>75</sup> Co	D : % $\beta^-$ n from 11Ho21<16								**	
* <sup>75</sup> Co	T : from 14Xu07=26.5(1.2); others 20Go10=27(13) 11Ho21=30(11)								**	
* <sup>75</sup> Cu	T : average 11Ho1=1.222(0.008) 91Kr15=1.224(0.003) 00KoZH=1.225(0.007)								**	
* <sup>75</sup> Cu	D : % $\beta^-$ n average 00KoZH=2.2(0.5) 85Re01=3.5(0.6)								**	
* <sup>75</sup> Cu	J : 20De21,17De30,10Vi07,11Ko36,09Fi03=5/2								**	
* <sup>75</sup> Cu <sup>m</sup>	J : from 19Ic02								**	
* <sup>75</sup> Cu <sup>n</sup>	J : from 19Ic02								**	
* <sup>75</sup> Ga	J : other 17Fa09=3/2								**	
* <sup>75</sup> Sr	T : average 19Si33=81.7(3.4) 03Hu01=88(3).								**	
* <sup>75</sup> Sr	T : other 01Ki13=71(+71-24) and 80(+400-40)								**	
<sup>76</sup> Fe	-10590#	600#			3# ms >410ns	0 <sup>+</sup>	17Su15 I	2017	$\beta^- ?$ *	
<sup>76</sup> Co	-25660#	500#			23 ms 6	(8 <sup>-</sup> )	14 14Xu07 TD	2010	$\beta^-=100; \beta^-n ?; \beta^-2n ?$ *	
<sup>76</sup> Co <sup>m</sup>	-25560#	510#	100#	100#	16 ms 4	(1 <sup>-</sup> )	15So23 TJD	2015	$\beta^-=100$ *	
<sup>76</sup> Co <sup>n</sup>	-24920#	510#	740#	100#	2.99 $\mu$ s 0.27	(3 <sup>+</sup> )	15So23 TJD	2015	IT=100 *	
<sup>76</sup> Ni	-42190#	300#			234.6 ms 2.7	0 <sup>+</sup>	07 14Xu07 T	1995	$\beta^-=100; \beta^-n=14.0$ 36 *	
<sup>76</sup> Ni <sup>m</sup>	-39770#	300#	2418.0	0.5	547.8 ns 3.3	(8 <sup>+</sup> )	07 15So23 TE	2005	IT=100 *	
<sup>76</sup> Cu	-50981.6	0.9			637.7 ms 5.5	3 <sup>-*</sup>	95 09Wi03 D	1987	$\beta^-=100; \beta^-n=7.2$ 5 *	
<sup>76</sup> Cu <sup>m</sup>			<i>non-exist</i>	RN	1.27 s 0.30	(1,3)	95 90Wi12 IJT	1990	$\beta^-=100$ *	
<sup>76</sup> Zn	-62303.0	1.5			5.7 s 0.3	0 <sup>+</sup>	95	1974	$\beta^-=100$ *	
<sup>76</sup> Ga	-66296.6	2.0			30.6 s 0.6	2 <sup>-*</sup>	95	1961	$\beta^-=100$ *	
<sup>76</sup> Ge	-73212.898	0.018			1.88 Zy 0.08	0 <sup>+</sup>	95 20Ba.A T	1933	IS=7.75 12; $\beta^-=100$ *	
<sup>76</sup> As	-72291.4	0.9			1.0933 d 0.0038	2 <sup>-*</sup>	95	1934	$\beta^-=100$ *	
<sup>76</sup> As <sup>m</sup>	-72247.0	0.9	44.425	0.001	1.84 $\mu$ s 0.06	(1) <sup>+</sup>	95	1966	IT=100 *	
<sup>76</sup> Se	-75251.959	0.016			STABLE	0 <sup>+</sup>	95	1922	IS=9.23 7 *	
<sup>76</sup> Br	-70289	9			16.2 h 0.2	1 <sup>-*</sup>	95	1952	$\beta^+=100$ *	
<sup>76</sup> Br <sup>m</sup>	-70186	9	102.58	0.03	1.31 s 0.02	(4) <sup>+</sup>	95	1979	IT $\approx$ 100; $\beta^+ < 0.6$ *	
<sup>76</sup> Kr	-69014	4			14.8 h 0.1	0 <sup>+</sup>	95	1954	$\beta^+=100$ *	
<sup>76</sup> Rb	-60479.1	0.9			36.5 s 0.6	1 <sup>-*</sup>	95	1969	$\beta^+=100; \beta^+ \alpha=3.8e-7$ 10 *	
<sup>76</sup> Rb <sup>m</sup>	-60162.2	0.9	316.93	0.08	3.050 $\mu$ s 0.007	(4) <sup>+</sup>	95 00Ch07 T	1986	IT=100 *	
<sup>76</sup> Sr	-54250	30			7.89 s 0.07	0 <sup>+</sup>	11	1990	$\beta^+=100; \beta^+p=3.4e-3$ 8 *	
<sup>76</sup> Y	-38250#	300#			28 ms 9	1 <sup>-#</sup>	07 19Si33 TJ	2001	$\beta^+ ?; p ?; \beta^+p ?$ *	
* <sup>76</sup> Co	T : symmetrized from 14Xu07=21.7(+6.5-4.9)								**	
* <sup>76</sup> Co	J : from 15So23								**	
* <sup>76</sup> Co <sup>n</sup>	E : 15So23=638.4(0.8) above <sup>76</sup> Co <sup>m</sup>								**	
* <sup>76</sup> Co <sup>n</sup>	T : symmetrized from 15So23=2.96(+0.29-0.25)								**	
* <sup>76</sup> Cu	T : average 10Ho12=599(18) 05Va19=653(24) 91Kr15=641(6)								**	
* <sup>76</sup> Cu	J : 20De21,17De30=3								**	
* <sup>76</sup> Cu <sup>m</sup>	I : reported only in 90Wi12; not confirmed in 05Va19								**	
* <sup>76</sup> Ga	T : average 16Do05=30.6(0.3) 85Ta01=32.6(0.6) 74Gr29=29.8(0.4) Birge B=2.7								**	
* <sup>76</sup> Ge	T : value for 2v- $\beta\beta$ ; other 15Ba11=1.65(+0.14-0.12) (evaluation).								**	
* <sup>76</sup> Ge	T : 0nu-BB 19Al24>27Yy, 18Aa02>19Yy, 18Ag03>80Yy,								**	
* <sup>76</sup> Ge	T : 13Ag11>30Yy combined GERDA+HDM+IGEX results; all at (90% C.L.);								**	
* <sup>76</sup> Ge	T : others 01K113=15 Yy 04K103=11.2 Yy not trusted. See also								**	
* <sup>76</sup> Ge	T : 02Aa.A and 02Zd02								**	
* <sup>76</sup> Rb	J : also 11Ma66=1								**	
* <sup>76</sup> Y	T : symmetrized from 19Si33=24(+12-6)								**	
* <sup>76</sup> Y	J : 19Si33=p5/2[422] n3/2[312], K=1-; similarity with the mirror <sup>76</sup> Br								**	
<sup>77</sup> Co	-21910#	600#			15 ms 6	7/2 <sup>-</sup> #	20	2014	$\beta^-=100; \beta^-n ?; \beta^-2n ?; \beta^-3n ?$ *	
<sup>77</sup> Ni	-37350#	400#			158.9 ms 4.2	9/2 <sup>+</sup> #	20 14Xu07 T	1995	$\beta^-=100; \beta^-n=26$ 13; $\beta^-2n ?$ *	
<sup>77</sup> Cu	-48862.8	1.2			470.3 ms 1.7	5/2 <sup>-*</sup>	20 20De21 J	1987	$\beta^-=100; \beta^-n=30.1$ 13 *	
<sup>77</sup> Zn	-58789.2	2.0			2.08 s 0.05	7/2 <sup>+</sup> *	20 17Wr01 J	1977	$\beta^+=100$ *	
<sup>77</sup> Zn <sup>m</sup>	-58016.8	2.0	772.440	0.015	1.05 s 0.10	1/2 <sup>-*</sup>	20 17Wr01 J	1986	$\beta^-=66$ 7; IT=34 7 *	
<sup>77</sup> Ga	-65992.4	2.4			13.2 s 0.2	3/2 <sup>-*</sup>	20	1968	$\beta^-=100$ [gs=12,m=88] *	
<sup>77</sup> Ge	-71212.87	0.05			11.211 h 0.003	7/2 <sup>+</sup>	20	1939	$\beta^-=100$ *	
<sup>77</sup> Ge <sup>m</sup>	-71053.16	0.08	159.71	0.06	53.7 s 0.6	1/2 <sup>-</sup>	20	1947	$\beta^-=81$ 2; IT=19 2 *	
<sup>77</sup> As	-73916.3	1.7			38.79 h 0.05	3/2 <sup>-</sup>	20	1951	$\beta^-=100$ *	

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{77}\text{As}^m$	-73440.8	1.7	475.48	0.04	114.0 $\mu\text{s}$ 2.5	9/2 <sup>+</sup>	20	1957	IT=100	
$^{77}\text{Se}$	-74599.50	0.06			STABLE	1/2 <sup>-</sup> *	20	1922	IS=7.60 7	
$^{77}\text{Se}^m$	-74437.58	0.06	161.9223	0.0010	17.36 s 0.05	7/2 <sup>+</sup>	20	1947	IT=100	
$^{77}\text{Br}$	-73234.8	2.8			57.04 h 0.12	3/2 <sup>-</sup> *	20	1948	$\beta^+$ =100	
$^{77}\text{Br}^m$	-73128.9	2.8	105.86	0.08	4.28 m 0.10	9/2 <sup>+</sup> *	20	1961	IT=100	
$^{77}\text{Kr}$	-70169.5	2.0			72.6 m 0.9	5/2 <sup>+</sup> *	20 19Ze02 T	1948	$\beta^+$ =100	*
$^{77}\text{Kr}^m$	-70103.0	2.0	66.50	0.05	118 ns 12	3/2 <sup>-</sup>	20	1975	IT=100	
$^{77}\text{Rb}$	-64830.5	1.3			3.78 m 0.04	3/2 <sup>-</sup> *	20	1972	$\beta^+$ =100	*
$^{77}\text{Sr}$	-57803	8			9.0 s 0.2	5/2 <sup>+</sup> *	20 13Ma15 J	1976	$\beta^+$ =100; $\beta^+$ p=0.08 3	
$^{77}\text{Y}$	-46440#	200#			63 ms 17	5/2 <sup>+</sup> #	20 00We.A D	1999	$\beta^+$ $\approx$ 100; $\beta^+$ p ?	*
$^{77}\text{Zr}$	-31600#	400#			100# $\mu\text{s}$	3/2 <sup>-</sup> #	20 17Su31 I	2017	$\beta^+$ ?; $\beta^+$ p ?	
* $^{77}\text{Co}$	T : symmetrized from 14Xu07=13.0(+7.2-4.3)									**
* $^{77}\text{Ni}$	D : % $\beta^-$ -n average 10Ho12=30(24) 14XuZZ=24(16)									**
* $^{77}\text{Cu}$	J : 20De21,17De30,11Ko36=5/2									**
* $^{77}\text{Cu}$	D : % $\beta^-$ -n average 18Ra27=29.2(3.0) 10Ho12=31.0(3.8) 09Ho1=30.3(2.0)									**
* $^{77}\text{Cu}$	D : 09Wi03=30.0(2.7); other 02Pf04=15(+10-5)									**
* $^{77}\text{Cu}$	T : average 14XuZZ=476.8(3.4) 09Pa35=467.4(2.1) 09Ti01=480(9) 91Kr15=469(8)									**
* $^{77}\text{Kr}$	T : average 19Ze02=71.25(42) 73Ba22=75(3) 71Bo30=74.7(0.4) 60Bu22=71.1(0.5)									**
* $^{77}\text{Kr}$	T : 57Be46=69(6); Birge ratio=3.46									**
* $^{77}\text{Kr}$	J : 95Ke04=5/2									**
* $^{77}\text{Rb}$	J : also 81Th04=3/2									**
* $^{77}\text{Y}$	T : symmetrized from 01Ki13=57(+22-12)									**
$^{78}\text{Co}$	-15320#	700#			11# ms >410ns		17Su15 I	2017	$\beta^-$ ?	
$^{78}\text{Ni}$	-34880#	400#			122.2 ms 5.1	0 <sup>+</sup>	09 14Xu07 T	1995	$\beta^-$ =100; $\beta^-$ -n ?; $\beta^-$ 2n ?	
$^{78}\text{Cu}$	-44789	13			330.7 ms 2.0	(6 <sup>-</sup> )*	09 14Xu07 T	1991	$\beta^-$ =100; $\beta^-$ -n=50.6 45;	*
									$\beta^-$ 2n ?	
$^{78}\text{Zn}$	-57483.2	1.9			1.47 s 0.15	0 <sup>+</sup>	09	1977	$\beta^-$ =100; $\beta^-$ -n ?	
$^{78}\text{Zn}^m$	-54809.5	2.0	2673.7	0.6	320 ns 6	(8 <sup>+</sup> )	09 12Ka36 ET	1998	IT=100	*
$^{78}\text{Ga}$	-63704.1	1.1			5.09 s 0.05	2 <sup>-</sup> *	09	1972	$\beta^-$ =100	
$^{78}\text{Ga}^m$	-63205.2	1.2	498.9	0.5	110 ns 3		09 10Da06 ET	2010	IT=100	*
$^{78}\text{Ge}$	-71862	4			88.0 m 1.0	0 <sup>+</sup>	09	1953	$\beta^-$ =100	
$^{78}\text{As}$	-72817	10			90.7 m 0.2	2 <sup>-</sup>	09	1937	$\beta^-$ =100	
$^{78}\text{Se}$	-77025.95	0.18			STABLE	0 <sup>+</sup>	09	1922	IS=23.69 22	
$^{78}\text{Br}$	-73452	4			6.45 m 0.04	1 <sup>+</sup> *	09 73Hi01 D	1937	$\beta^+$ $\approx$ 100; $\beta^-$ <0.01	
$^{78}\text{Br}^m$	-73271	4	180.89	0.13	119.4 $\mu\text{s}$ 1.0	(4 <sup>+</sup> )	09	1958	IT=100	
$^{78}\text{Kr}$	-74178.3	0.3			STABLE	0 <sup>+</sup>	09 94Sa31 T	1920	IS=0.355 3;2 $\beta^+$ ?	*
$^{78}\text{Rb}$	-66935	3			17.66 m 0.03	0 <sup>+</sup> *	09	1968	$\beta^+$ =100	*
$^{78}\text{Rb}^m$	-66888	3	46.84	0.14	910 ns 40	(1 <sup>-</sup> )	09	1996	IT=100	
$^{78}\text{Rb}^n$	-66824	3	111.19	0.22	5.74 m 0.03	4 <sup>-</sup> *	09	1968	$\beta^+$ =91 2;IT=9 2	*
$^{78}\text{Rb}^x$	-66861	12	74	12	R = 2.0 0.5	<i>spmix</i>				
$^{78}\text{Sr}$	-63174	7			156.1 s 2.7	0 <sup>+</sup>	09 11Pe29 T	1982	$\beta^+$ =100	*
$^{78}\text{Y}$	-52170#	300#		*	54 ms 5	(0 <sup>+</sup> )	09 01Ga24 TJ	1992	$\beta^+$ =100; $\beta^+$ p ?	*
$^{78}\text{Y}^m$	-52170#	580#	0#	500#	5.8 s 0.6	(5 <sup>+</sup> )	09	1998	$\beta^+$ =100; $\beta^+$ p ?	
$^{78}\text{Zr}$	-40850#	400#			50# ms >200ns	0 <sup>+</sup>	09 01Ki13 I	2001	$\beta^+$ ?; $\beta^+$ p ?	*
* $^{78}\text{Cu}$	D : % $\beta^-$ -n average 10Ho12=44.0(5.4) 09Wi03=65(8)									**
* $^{78}\text{Cu}$	J : 20De21,17De30,11Ko36=(6); other 12Ko29=(5)									**
* $^{78}\text{Zn}^m$	E : from 12Ko29; other 12Ka36=2675.3(1.0)									**
* $^{78}\text{Zn}^m$	T : average 12Ka36=320(+9-8) 00Da07=319(9)									**
* $^{78}\text{Ga}^m$	ET : other E=559.6(0.7) keV, T1/2<500 ns in Ensdf2009									**
* $^{78}\text{Kr}$	T : limit given here is for the K-e <sup>+</sup> decay (theoretically faster)									**
* $^{78}\text{Rb}$	J : other 11Ma66,81Th04=0									**
* $^{78}\text{Rb}^n$	J : other 11Ma66,81Th04=4									**
* $^{78}\text{Sr}$	T : average 11Pe29=155(3) 97Mu02=168(12) 92Gr09=159(8)									**
* $^{78}\text{Y}$	T : average 01Ga24=50(8) 01Ki13=55(+9-6)									**
* $^{78}\text{Zr}$	I : other 00We.A>170 ns, same group as 01Ki13									**
$^{79}\text{Ni}$	-28160#	500#			44 ms 8	5/2 <sup>+</sup> #	16	2010	$\beta^-$ =100; $\beta^-$ -n ?; $\beta^-$ 2n ?	
$^{79}\text{Cu}$	-42410	100			241.3 ms 2.1	(5/2 <sup>-</sup> )*	16 14Xu07 T	1991	$\beta^-$ =100; $\beta^-$ -n=66 10; $\beta^-$ 2n ?	*
$^{79}\text{Zn}$	-53432.3	2.2			746 ms 42	9/2 <sup>+</sup> *	16	1981	$\beta^-$ =100; $\beta^-$ -n=1.7 5	*
$^{79}\text{Zn}^m$	-52330	150	1100	150	> 200 ms	1/2 <sup>+</sup> *	16 17Wr01 J	2015	IT ?; $\beta^-$ ?	*

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>79</sup> Ga	-62548.4	1.2			2.848 s 0.003	3/2 <sup>-</sup> *	16	1974	$\beta^-$ =100; $\beta^-$ -n=0.089 19	*
<sup>79</sup> Ge	-69530	40			18.98 s 0.03	(1/2) <sup>-</sup>	16	1970	$\beta^-$ =100	
<sup>79</sup> Ge <sup>m</sup>	-69340	40	185.95	0.04	39.0 s 1.0	7/2 <sup>+</sup> #	16	1970	$\beta^-$ =96 1;IT=4 1	
<sup>79</sup> As	-73636	5			9.01 m 0.15	3/2 <sup>-</sup>	16	1950	$\beta^-$ =100	
<sup>79</sup> As <sup>m</sup>	-72863	5	772.81	0.06	1.21 $\mu$ s 0.01	(9/2) <sup>+</sup>	16 98Gr14 T	1998	IT=100	
<sup>79</sup> Se	-75917.47	0.22			327 ky 28	7/2 <sup>+</sup> *	16	1950	$\beta^-$ =100	
<sup>79</sup> Se <sup>m</sup>	-75821.70	0.22	95.77	0.03	3.900 m 0.018	1/2 <sup>-</sup>	16 88KI03 D	1950	IT $\approx$ 100; $\beta^-$ =0.056 11	*
<sup>79</sup> Br	-76068.1	1.0			STABLE	3/2 <sup>-</sup> *	16	1920	IS=50.65 9	
<sup>79</sup> Br <sup>m</sup>	-75860.5	1.0	207.61	0.09	4.85 s 0.04	9/2 <sup>+</sup>	16	1954	IT=100	
<sup>79</sup> Kr	-74442	3			35.04 h 0.10	1/2 <sup>-</sup> *	16 95Ke04 J	1948	$\beta^+$ =100	
<sup>79</sup> Kr <sup>m</sup>	-74312	3	129.77	0.05	50 s 3	7/2 <sup>+</sup> *	16 95Ke04 J	1940	IT=100	
<sup>79</sup> Rb	-70802.8	1.9			22.9 m 0.5	5/2 <sup>+</sup> *	16	1957	$\beta^+$ =100	*
<sup>79</sup> Sr	-65480	7			2.25 m 0.10	3/2 <sup>-</sup> *	16	1972	$\beta^+$ =100	*
<sup>79</sup> Y	-57800	80			14.8 s 0.6	5/2 <sup>+</sup> #	16	1992	$\beta^+$ =100	
<sup>79</sup> Zr	-46770#	300#			56 ms 30	5/2 <sup>+</sup> #	16	1999	$\beta^+$ =100; $\beta^+$ p ?	
<sup>79</sup> Nb	-31650#	500#				9/2 <sup>+</sup> #			p ?; $\beta^+$ ?; $\beta^+$ p ?	
* <sup>79</sup> Cu	J : 17De30=(5/2)									**
* <sup>79</sup> Cu	T : others 10Ho12=257(+29-26) 91Kr15=188(25)									**
* <sup>79</sup> Cu	D : % $\beta^-$ -n average 10Ho12=72(12) 91Kr15=55(17)									**
* <sup>79</sup> Zn	J : 17Wr01,16Ya02=9/2									**
* <sup>79</sup> Zn <sup>m</sup>	J : 17Wr01,16Ya02=1/2									**
* <sup>79</sup> Ga	J : also 17Fa09=3/2									**
* <sup>79</sup> Se <sup>m</sup>	T : average 19De24=3.884(0.009) (quoted in the text in hours is a typo)									**
* <sup>79</sup> Se <sup>m</sup>	T : 90Ab02=3.92(0.01); Birge ratio=2.68									**
* <sup>79</sup> Rb	J : also 81Th04=5/2									**
* <sup>79</sup> Sr	J : 90Li28=3/2									**
<sup>80</sup> Ni	-23240#	600#			30 ms 22	0 <sup>+</sup>	14	2014	$\beta^-$ =100; $\beta^-$ -n ?; $\beta^-$ 2n ?	*
<sup>80</sup> Cu	-36680#	300#			113.3 ms 6.4		14 14Xu07 T	1995	$\beta^-$ =100; $\beta^-$ -n=58 9; $\beta^-$ 2n ?	*
<sup>80</sup> Zn	-51648.6	2.6			562.2 ms 3.0	0 <sup>+</sup>	14 14Xu07 T	1981	$\beta^-$ =100; $\beta^-$ -n=1.36 12	*
<sup>80</sup> Ga	-59223.7	2.9			1.9 s 0.1	6 <sup>-</sup> *	14 13Ve03 TJ	1974	$\beta^-$ =100; $\beta^-$ -n=0.86 7	*
<sup>80</sup> Ga <sup>m</sup>	-59201.3	2.9	22.45	0.10	1.3 s 0.2	3 <sup>-</sup> *	14 13Ve03 TJ	2011	$\beta^-$ $\approx$ 100; $\beta^-$ -n ?;IT ?	
<sup>80</sup> Ge	-69535.3	2.1			29.5 s 0.4	0 <sup>+</sup>	05	1972	$\beta^-$ =100	
<sup>80</sup> As	-72215	3			15.2 s 0.2	1 <sup>+</sup>	05	1954	$\beta^-$ =100	
<sup>80</sup> Se	-77759.5	0.9			STABLE	0 <sup>+</sup>	05	1922	IS=49.80 36;2 $\beta^-$ ?	
<sup>80</sup> Br	-75889.0	1.0			17.68 m 0.02	1 <sup>+</sup> *	05	1937	$\beta^-$ =91.7 2; $\beta^+$ =8.3 2	
<sup>80</sup> Br <sup>m</sup>	-75803.2	1.0	85.843	0.004	4.4205 h 0.0008	5 <sup>-</sup> *	05	1937	IT=100	
<sup>80</sup> Kr	-77893.5	0.7			STABLE	0 <sup>+</sup>	05	1920	IS=2.286 10	
<sup>80</sup> Rb	-72175.5	1.9			33.4 s 0.7	1 <sup>+</sup> *	05 93Al03 T	1961	$\beta^+$ =100	*
<sup>80</sup> Rb <sup>m</sup>	-71681.6	2.0	493.9	0.5	1.63 $\mu$ s 0.04	(6 <sup>+</sup> )	05 92Do10 E	1980	IT=100	
<sup>80</sup> Sr	-70311	3			106.3 m 1.5	0 <sup>+</sup>	05	1961	$\beta^+$ =100	
<sup>80</sup> Y	-61148	6			30.1 s 0.5	4 <sup>-</sup>	05	1981	$\beta^+$ =100	
<sup>80</sup> Y <sup>m</sup>	-60920	6	228.5	0.1	4.8 s 0.3	1 <sup>-</sup>	05 01No07 J	1998	IT=81 2; $\beta^+$ =19 2	*
<sup>80</sup> Y <sup>n</sup>	-60835	6	312.6	0.9	4.7 $\mu$ s 0.3	(2 <sup>+</sup> )	05	1997	IT=100	
<sup>80</sup> Zr	-54760#	300#			4.6 s 0.6	0 <sup>+</sup>	05 01Ki13 T	1987	$\beta^+$ =100	*
<sup>80</sup> Nb	-38420#	400#				4 <sup>-</sup> #			p ?; $\beta^+$ ?; $\beta^+$ p ?	
* <sup>80</sup> Ni	T : symmetrized from 14Xu07=23.9(+26.0-17.2)									**
* <sup>80</sup> Cu	T : other 10Ho12=170(+110-50)									**
* <sup>80</sup> Cu	D : % $\beta^-$ -n from 14XuZZ									**
* <sup>80</sup> Zn	D : % $\beta^-$ -n from 19To09; others 91Kr15=1.0(0.5) 10Ho12<1.8%									**
* <sup>80</sup> Ga	D : % $\beta^-$ -n is probably a mixture of values for <sup>80</sup> Ga and <sup>80</sup> Ga <sup>m</sup>									**
* <sup>80</sup> Rb	J : also 81Th04=1									**
* <sup>80</sup> Y <sup>m</sup>	J : 228.5 keV M3 to 4-									**
* <sup>80</sup> Zr	T : average 01Ki13=5.3(+1.1-0.9) 00Re03=4.1(+0.8-0.6)									**
<sup>81</sup> Ni	-16090#	700#			30# ms >410ns	3/2 <sup>+</sup> #	17Su15 I	2017	$\beta^-$ ?	
<sup>81</sup> Cu	-31910#	300#			73.2 ms 6.8	5/2 <sup>-</sup> #	10 14Xu07 TD	2010	$\beta^-$ =100; $\beta^-$ -n=81 20; $\beta^-$ 2n ?	*
<sup>81</sup> Zn	-46200	5			299.4 ms 2.1	(1/2 <sup>+</sup> , 5/2 <sup>+</sup> )	08 20Pa26 TDJ	1991	$\beta^-$ =100; $\beta^-$ -n=23 4; $\beta^-$ 2n ?	*
<sup>81</sup> Ga	-57628	3			1.217 s 0.005	5/2 <sup>-</sup> *	08 20Pa26 T	1976	$\beta^-$ =100; $\beta^-$ -n=12.5 5	*
<sup>81</sup> Ge	-66291.7	2.1			9 s 2	9/2 <sup>+</sup> #	08	1972	$\beta^-$ =100	*
<sup>81</sup> Ge <sup>m</sup>	-65612.6	2.1	679.14	0.04	6 s 2	(1/2 <sup>+</sup> )	08	1981	$\beta^-$ $\approx$ 100;IT<1	*

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>81</sup> As	-72533.3	2.6	33.3 s 0.8	3/2 <sup>-</sup>	08	1960	$\beta^-$ =100	
<sup>81</sup> Se	-76389.0	1.0	18.45 m 0.12	1/2 <sup>-</sup>	08	1948	$\beta^-$ =100	
<sup>81</sup> Se <sup>m</sup>	-76286.0	1.0	103.00 0.06	57.28 m 0.02	7/2 <sup>+</sup>	08	1971	IT≈100; $\beta^-$ =0.051 14
<sup>81</sup> Br	-77977.1	1.0	STABLE	3/2 <sup>-*</sup>	08	1920	IS=49.35 9	
<sup>81</sup> Br <sup>m</sup>	-77440.9	1.0	536.20 0.09	34.6 $\mu$ s 2.8	9/2 <sup>+</sup>	08	1967	IT=100
<sup>81</sup> Kr	-77696.2	1.1	229 ky 11	7/2 <sup>+</sup> *	08 95Ke04 J	1950	$\epsilon$ =100	
<sup>81</sup> Kr <sup>m</sup>	-77505.6	1.1	190.64 0.04	13.10 s 0.03	1/2 <sup>-*</sup>	08 95Ke04 J	1940	IT≈100; $\epsilon$ =0.0025 4
<sup>81</sup> Rb	-75457	5	4.572 h 0.004	3/2 <sup>-*</sup>	08	1949	$\beta^+$ =100	
<sup>81</sup> Rb <sup>m</sup>	-75371	5	86.31 0.07	30.5 m 0.3	9/2 <sup>+</sup> *	08	1956	IT=97.6 6; $\beta^+$ =2.4 6
<sup>81</sup> Sr	-71528	3	22.3 m 0.4	1/2 <sup>-*</sup>	08	1952	$\beta^+$ =100	
<sup>81</sup> Sr <sup>m</sup>	-71449	3	79.23 0.04	390 ns 50	(5/2) <sup>-</sup>	08	1983	IT=100
<sup>81</sup> Sr <sup>n</sup>	-71439	3	89.05 0.07	6.4 $\mu$ s 0.5	(7/2 <sup>+</sup> )	08	1989	IT ?
<sup>81</sup> Y	-65713	5	70.4 s 1.0	(5/2 <sup>+</sup> )	08	1981	$\beta^+$ =100	
<sup>81</sup> Zr	-57520	90	5.5 s 0.4	(3/2 <sup>-</sup> )	08	1997	$\beta^+$ =100; $\beta^+$ p=0.12 2	
<sup>81</sup> Nb	-46360#	400#	<44ns	9/2 <sup>+</sup> #	08 00We.A I		p ?; $\beta^+$ ?; $\beta^+$ p ?	
<sup>81</sup> Mo	-31460#	500#	1# ms >400ns	5/2 <sup>+</sup> #	15 13Su23 I	2013	$\beta^+$ ?; $\beta^+$ p ?	
* <sup>81</sup> Cu	D : % $\beta^-$ -n from 14XuZZ							**
* <sup>81</sup> Zn	D : % $\beta^-$ -n from 20Pa26; others 12Ma37=12(4) 91Kr15=7.5(3.0) 10Ho12=30(13)							**
* <sup>81</sup> Zn	T : average 20Pa26=290(4) 14Xu07=303.2(2.6) 10Pa33=304(13)							**
* <sup>81</sup> Ga	J : also 17Fa09=3/2							**
* <sup>81</sup> Ga	D : % $\beta^-$ -n average 19To09=11.2(2.6) 93Ru01=12.9(0.8) 81Ho07=12.7(1.2)							**
* <sup>81</sup> Ga	D : 80Lu04=12.0(0.9); others 10Ho12<21 86ReZU=11.7(1.2)							**
* <sup>81</sup> Ge	T : derived from 81Ho24=7.6(0.6) 72De43=10.1(0.8) for a mixture of gs and							**
* <sup>81</sup> Ge	T : isomer that have similar T1/2							**
* <sup>81</sup> Ge <sup>m</sup>	T : derived from 81Ho24=7.6(0.6) 72De43=10.1(0.8) for mixture of gs and							**
* <sup>81</sup> Ge <sup>m</sup>	T : isomer that have similar T1/2							**
* <sup>81</sup> Rb	J : also 81Th04=3/2							**
* <sup>81</sup> Rb <sup>m</sup>	J : also 81Th04=9/2							**
* <sup>81</sup> Nb	I : also 99Ja02<80 ns 01Ki13<200 ns							**
* <sup>81</sup> Nb	T : estimated $\beta^+$ half-life 01Ki13=100# ms							**
<sup>82</sup> Ni	-10720#	800#	16# ms >410ns	0 <sup>+</sup>	19 17Su15 I	2017	$\beta^-$ ?	
<sup>82</sup> Cu	-25730#	400#	34 ms 7		19 10Oh02 I	2010	$\beta^-$ =100; $\beta^-$ -n ?; $\beta^-$ 2n ?	
<sup>82</sup> Zn	-42314	3	177.9 ms 2.5	0 <sup>+</sup>	19 14Xu07 T	1997	$\beta^-$ =100; $\beta^-$ -n=69 7; $\beta^-$ 2n ?	
<sup>82</sup> Ga	-52930.7	2.4	600 ms 2	2 <sup>-*</sup>	19 12Ch51 J	1976	$\beta^-$ =100; $\beta^-$ -n=21.2 10; $\beta^-$ 2n ?	
<sup>82</sup> Ga <sup>m</sup>	-52790.0	2.4	140.7 0.3	93.5 ns 6.7	(4 <sup>-</sup> )	19 16Al10 TJ	2009	IT=100
<sup>82</sup> Ge	-65415.1	2.2	4.31 s 0.19	4.31 s 0.19	0 <sup>+</sup>	19	1972	$\beta^-$ =100
<sup>82</sup> As	-70105	4	19.1 s 0.5	19.1 s 0.5	(2 <sup>-</sup> )	19 04Ga44 J	1968	$\beta^-$ =100
<sup>82</sup> As <sup>m</sup>	-69973	4	131.6 0.5	13.6 s 0.4	(5 <sup>-</sup> )	19	1970	$\beta^-$ =100
<sup>82</sup> Se	-77593.9	0.5	87.6 Ey 1.5	87.6 Ey 1.5	0 <sup>+</sup>	19 20Ba.A T	1922	IS=8.82 15;2 $\beta^-$ =100
<sup>82</sup> Br	-77498.7	1.0	35.282 h 0.007	35.282 h 0.007	5 <sup>-*</sup>	19 81Th04 J	1937	$\beta^-$ =100
<sup>82</sup> Br <sup>m</sup>	-77452.8	1.0	45.9492 0.0010	6.13 m 0.05	2 <sup>-</sup>	19	1965	IT=97.6 3; $\beta^-$ =2.4 3
<sup>82</sup> Kr	-80591.795	0.006	STABLE	STABLE	0 <sup>+</sup>	19	1920	IS=11.593 31
<sup>82</sup> Rb	-76188	3	1.2575 m 0.0002	1.2575 m 0.0002	1 <sup>+</sup> *	19 81Th04 J	1949	$\beta^+$ =100
<sup>82</sup> Rb <sup>m</sup>	-76118.8	2.6	69.0 1.5 IT	6.472 h 0.006	5 <sup>-*</sup>	19 81Th04 J	1957	$\beta^+$ ≈100;IT<0.33
<sup>82</sup> Sr	-76010	6	25.35 d 0.03	25.35 d 0.03	0 <sup>+</sup>	19	1952	$\epsilon$ =100
<sup>82</sup> Y	-68064	5	8.30 s 0.20	8.30 s 0.20	1 <sup>+</sup>	19	1980	$\beta^+$ =100
<sup>82</sup> Y <sup>m</sup>	-67661	5	402.63 0.14	258 ns 22	4 <sup>-</sup>	19 94Mu02 T	1994	IT=100
<sup>82</sup> Y <sup>n</sup>	-67557	5	507.50 0.13	148 ns 6	6 <sup>+</sup>	19	1994	IT=100
<sup>82</sup> Zr	-63614.1	1.6	32 s 5	32 s 5	0 <sup>+</sup>	19	1982	$\beta^+$ =100
<sup>82</sup> Nb	-51810#	300#	51 ms 5	51 ms 5	(0 <sup>+</sup> )	19	1992	$\beta^+$ =100; $\beta^+$ p ?
<sup>82</sup> Nb <sup>m</sup>	-50630#	300#	1180 1	93 ns 20	(5 <sup>+</sup> )	19 09Ga40 ETJ	2008	IT=100
<sup>82</sup> Mo	-40370#	400#	30# ms >400ns	30# ms >400ns	0 <sup>+</sup>	19 13Su23 I	2013	$\beta^+$ ?; $\beta^+$ p ?
* <sup>82</sup> Cu	T : symmetrized from 14Xu07=33(+7-6)							**
* <sup>82</sup> Zn	T : others 16Al10=155(26) 12Ma37=228(10) outweighed (not used)							**
* <sup>82</sup> Ga	D : % $\beta^-$ -n average 17Ve01=22(2) 16Te09=22.2(2.0) 86Wa17=19.8(1.7)							**
* <sup>82</sup> Ga	D : 80Lu04=21.4(2.2); other 93Ru01=31.1(4.4)% at variance (not used)							**
* <sup>82</sup> Ga <sup>m</sup>	T : average 16Al10=89(9) 12Ka36=98(+10-9); other 09Fo05<500 ns							**
* <sup>82</sup> Ge	T : average 15Et01=4.04(0.27) 80Ze.A=4.5(0.4) 72De43=4.60(0.35);							**
* <sup>82</sup> Ge	T : other (not used) 73Kr.A=5(1)							**
* <sup>82</sup> Se	T : 2v- $\beta\beta$ symmetrized from 20Ba.A=87(+2-1) (evaluation); others							**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>82</sup> Se	T : 19Az04=86.0(0.3)(+1.9-1.3) 12Si23=96(10) 15Ba11=92(7) (evaluation);						**
* <sup>82</sup> Se	T : 0nu-BB: 19Az02>3.5 Yy 18Az05>2.4 Yy 99Pi08=83(+9-7) 98Ar10=83(12)						**
* <sup>82</sup> Se	T : 92El07=108(+26-6) 88Li11=120(10)						**
* <sup>82</sup> Y <sup>m</sup>	T : average 94Mu02=220(50) 93Wo04=268(25)						**
* <sup>82</sup> Nb <sup>m</sup>	T : 09Ga40 superseded 08Ga04=92(17); other 07Ca26=80(50)						**
<sup>83</sup> Cu	-20390#	500#					
<sup>83</sup> Zn	-36290#	300#	21# ms >410ns	5/2 <sup>-</sup> #	17Su15 I	2017	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
<sup>83</sup> Ga	-49257.1	2.6	100 ms 3	3/2 <sup>+</sup> #	15 14XuZZ	TD 1997	$\beta^-$ =100; $\beta^-n$ ≈71 29; $\beta^-2n$ ? *
<sup>83</sup> Ga <sup>m</sup>	-49059.8	2.6	310.0 ms 0.7	5/2 <sup>-</sup> #	15 17Ve01	TD 1976	$\beta^-$ =100; $\beta^-n$ =85 4; $\beta^-2n$ ? *
<sup>83</sup> Ge	-60976.4	2.4	120 ns 5		16Al10	ETD2016	IT=100
<sup>83</sup> As	-69669.3	2.8	1.85 s 0.06	(5/2 <sup>+</sup> )	15	1972	$\beta^-$ =100; $\beta^-n$ ?
<sup>83</sup> Se	-75341	3	13.4 s 0.4	5/2 <sup>-</sup> #	15	1968	$\beta^-$ =100
<sup>83</sup> Se <sup>m</sup>	-75112	3	22.25 m 0.04	9/2 <sup>+</sup>	15 15Kr02	T 1937	$\beta^-$ =100
<sup>83</sup> Br	-79014	4	70.1 s 0.4	1/2 <sup>-</sup>	15	1969	$\beta^-$ =100
<sup>83</sup> Br <sup>m</sup>	-75945	4	2.374 h 0.004	3/2 <sup>-</sup>	15	1937	$\beta^-$ =100
<sup>83</sup> Kr	-79990.643	0.009	729 ns 77	(19/2 <sup>-</sup> )	15 11Ru.A	T 1989	IT=100 *
<sup>83</sup> Kr <sup>m</sup>	-79981.238	0.009	STABLE	9/2 <sup>+</sup> *	15	1920	IS=11.500 19 *
<sup>83</sup> Kr <sup>n</sup>	-79949.086	0.009	156.8 ns 0.5	7/2 <sup>+</sup>	15	1963	IT=100
<sup>83</sup> Rb	-79070.6	2.3	1.830 h 0.013	1/2 <sup>-</sup> *	15 10Li13	T 1971	IT=100 *
<sup>83</sup> Rb <sup>m</sup>	-79028.5	2.3	86.2 d 0.1	5/2 <sup>-</sup> *	15	1950	$\epsilon$ =100 *
<sup>83</sup> Sr	-76798	7	7.8 ms 0.7	9/2 <sup>+</sup>	15 68Et01	T 1968	IT=100
<sup>83</sup> Sr <sup>m</sup>	-76539	7	32.41 h 0.03	7/2 <sup>+</sup> *	15	1952	$\beta^+$ =100 *
<sup>83</sup> Y	-72206	19	4.95 s 0.12	1/2 <sup>-</sup> *	15	1972	IT=100 *
<sup>83</sup> Y <sup>m</sup>	-72144	19	7.08 m 0.08	(9/2 <sup>+</sup> )	15 92Bu10	J 1962	$\beta^+$ =100
<sup>83</sup> Zr	-65912	6	2.85 m 0.02	(3/2 <sup>-</sup> )	15	1972	$\beta^+$ =60 5; IT=40 5
<sup>83</sup> Zr <sup>m</sup>	-65859	6	42 s 2	1/2 <sup>-</sup> #	15	1974	$\beta^+$ =100; $\beta^+p$ ?
<sup>83</sup> Zr <sup>n</sup>	-65835	6	530 ns 120	(5/2 <sup>-</sup> )	15	1988	IT=100
<sup>83</sup> Nb	-57610	160	1.8 $\mu$ s 0.1	(7/2 <sup>+</sup> )	15	1988	IT=100
<sup>83</sup> Mo	-46340#	400#	3.9 s 0.2	9/2 <sup>+</sup> #	15	1988	$\beta^+$ =100
<sup>83</sup> Tc	-31320#	500#	23 ms 19	3/2 <sup>-</sup> #	15 01Ki13	TD 1999	$\beta^+$ =100; $\beta^+p$ ? *
* <sup>83</sup> Zn	T : average 16Al10=122(28) 14XuZZ=99.4(3.0) 12Ma37=117(20)			1/2 <sup>-</sup> #			$p$ ?; $\beta^+$ ?; $\beta^+p$ ?
* <sup>83</sup> Ga	T : average 17Ve01=312(1) 16Al10=309(6) 14Xu.A=296.1(6.4) 06Pe20=317(17)						**
* <sup>83</sup> Ga	T : 319(24) 93Ru01=307(7) 91Kr15=308(1) 86Wa17,80Lu04,76Ru01=310(10)						**
* <sup>83</sup> Ga	D : % $\beta^-n$ others 16Ma50=56(7) 09Wi03=62.8(2.5) 80Lu04=43(7) 93Ru01=14.9(1.8)						**
* <sup>83</sup> Br <sup>m</sup>	T : average 11Ru.A=862(148) 97Is13=700(100) 89Wi01=600(200)						**
* <sup>83</sup> Kr	J : also 95Ke04=9/2						**
* <sup>83</sup> Kr <sup>n</sup>	T : average 10Li13=1.82(0.02) 09Ka30=1.85(0.03) 71Ru17=1.83(0.02)						**
* <sup>83</sup> Kr <sup>n</sup>	J : 95Ke04=1/2						**
* <sup>83</sup> Rb	J : also 81Th04=5/2						**
* <sup>83</sup> Sr	J : 90Li28=7/2						**
* <sup>83</sup> Sr <sup>m</sup>	J : 90Li28=1/2						**
* <sup>83</sup> Mo	T : symmetrized from 01Ki13=6(+30-3)						**
<sup>84</sup> Cu	-13720#	500#					
<sup>84</sup> Zn	-31830#	400#	54 ms 8	0 <sup>+</sup>	10 14XuZZ	TD 2010	$\beta^-$ ?; $\beta^-n$ ?
<sup>84</sup> Ga	-44094	30	97.6 ms 1.2	0 <sup>-</sup> #	09 19Yo03	TD 1991	$\beta^-$ =100; $\beta^-n$ =73 26; $\beta^-2n$ ? *
<sup>84</sup> Ga <sup>m</sup>		<i>non-exist</i>					$\beta^-$ =100; $\beta^-n$ =43 4; $\beta^-2n$ =1.6 2 *
<sup>84</sup> Ge	-58148	3	< 85 ms	(3 <sup>-</sup> , 4 <sup>-</sup> )	09 09Le26	TD	$\beta^-$ ?; IT ? *
<sup>84</sup> As	-65854	3	951 ms 9	0 <sup>+</sup>	09 13Ma22	T 1972	$\beta^-$ =100; $\beta^-n$ =10.6 6 *
<sup>84</sup> As <sup>m</sup>		<i>non-exist</i>					$\beta^-$ =100; $\beta^-n$ =0.28 4 *
<sup>84</sup> Se	-75947.7	2.0	3.16 s 0.58	(2 <sup>-</sup> )	09 16Ko24	J 1968	$\beta^-$ =100
<sup>84</sup> Br	-77783	26	650 ms 150		09 74KrZG	IT 1974	$\beta^-$ =100 *
<sup>84</sup> Br <sup>m</sup>	-77470	100	3.26 m 0.10	0 <sup>+</sup>	09	1960	$\beta^-$ =100
<sup>84</sup> Br <sup>n</sup>	-77375	26	31.76 m 0.08	2 <sup>-</sup>	09	1943	$\beta^-$ =100
<sup>84</sup> Kr	-82439.345	0.004	6.0 m 0.2	(6 <sup>-</sup> )	09	1957	$\beta^-$ =100
<sup>84</sup> Kr <sup>m</sup>	-79203.27	0.18	< 140 ns	1 <sup>+</sup>	09	1970	IT=100
<sup>84</sup> Rb	-79759.0	2.2	STABLE	0 <sup>+</sup>	09	1920	IS=56.987 15
<sup>84</sup> Rb <sup>m</sup>	-79295.4	2.2	1.83 $\mu$ s 0.04	8 <sup>+</sup>	09	1982	IT=100
<sup>84</sup> Sr	-80649.6	1.2	32.82 d 0.07	2 <sup>-</sup> *	09	1947	$\beta^+$ =96.1 20; $\beta^-$ =3.9 20 *
			20.26 m 0.04	6 <sup>-</sup> *	09	1940	IT≈100; $\beta^+$ <0.0012 *
			STABLE	0 <sup>+</sup>	09	1936	IS=00.56 2; $\beta^+$ ?

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{84}\text{Y}$	-73894	4	39.5 m 0.8	(6 <sup>+</sup> )	09	1962	$\beta^+=100$
$^{84}\text{Y}^m$	-73827	4	67.0 0.2	1 <sup>+</sup>	09	1976	$\beta^+=100$
$^{84}\text{Y}^n$	-73684	4	210.42 0.16	(4 <sup>-</sup> )	09	2005	IT=100
$^{84}\text{Zr}$	-71422	5	25.8 m 0.5	0 <sup>+</sup>	09	1977	$\beta^+=100$
$^{84}\text{Nb}$	-61193.8	0.4	9.8 s 0.9	(1 <sup>+</sup> )	09 09St04 J	1977	$\beta^+=100$
$^{84}\text{Nb}^m$	-61145.8	1.1	48 1	(3 <sup>+</sup> )	09Ga40 ETJ	2009	IT=100
$^{84}\text{Nb}^n$	-60856.1	0.6	337.7 0.4	(5 <sup>-</sup> )	09 09Ga40 T	2000	IT=100
$^{84}\text{Mo}$	-54170#	300#	2.3 s 0.3	0 <sup>+</sup>	09	1991	$\beta^+=100; \beta^+p ?$
$^{84}\text{Tc}$	-37700#	400#		1 <sup>+</sup> #			$p ?; \beta^+ ?; \beta^+p ?$
$^{84}\text{Ga}$	D : % $\beta^-$ -n average 19Yo03=44(4) and 16Ma50=40(7) (same group as 19Yo03, but						
$^{84}\text{Ga}$	D : different experiment); others 17Ve01=53(20) 10Wi03=74(14)						
$^{84}\text{Ga}$	D : 09Gr06=80(15) 91Kr15=70(15)						
$^{84}\text{Ga}^m$	I : proposed in 09Le26 (and Ensdf2009), but not confirmed in 10Wi03 data						
$^{84}\text{Ga}^m$	I : of much bigger statistics						
$^{84}\text{Ge}$	T : average 13Ma22=942(17) 93Ru01=947(11) 91Kr15=984(23)						
$^{84}\text{Ge}$	D : % $\beta^-$ -n average 93Ru01=10.8(0.6) 91Kr15=9.5(2.0) 91Om01=9(3)						
$^{84}\text{As}$	T : from 13Ma22; others: 96WaZX=3.24(0.26) 93Ru01=4.02(0.03) 91Om01=4.5(0.2)						
$^{84}\text{As}$	T : 81Ho10=4.5(0.2) 75Kr08=5.3(0.4) 68De19=5.8(0.5)						
$^{84}\text{As}$	D : % $\beta^-$ -n from 91Ru01; others 02Pf04=0.18(0.10) 73Kr06=0.13(0.06)						
$^{84}\text{As}^m$	I : from 74KrZG (also 75Kr08), but not confirmed in 81Ho10						
$^{84}\text{Rb}$	J : other 14Ya28=1.9(1), 81Th04=2						
$^{84}\text{Rb}^m$	J : other 14Ya28=6.2(2), 81Th04=6						
$^{85}\text{Zn}$	-25100#	500#	40# ms >400ns	5/2 <sup>+</sup> #	14 10Oh02 I	2010	$\beta^- ?; \beta^-n ?; \beta^-2n ?$
$^{85}\text{Ga}$	-39740	40	95.3 ms 1.0	(5/2 <sup>-</sup> )	14 19Yo03 TD	1997	$\beta^-=100; \beta^-n=77.4;$ $\beta^-2n=1.3.2$
$^{85}\text{Ge}$	-53123	4	495 ms 5	(3/2 <sup>+</sup> , 5/2 <sup>+</sup> )#	14 13Ma22 T	1991	$\beta^-=100; \beta^-n=17.2.18;$ $\beta^-2n ?$
$^{85}\text{As}$	-63189	3	2.022 s 0.007	(5/2 <sup>-</sup> )	14 12Ku06 J	1967	$\beta^-=100; \beta^-n=62.6.9$
$^{85}\text{Se}$	-72413.6	2.6	32.9 s 0.3	(5/2 <sup>+</sup> )	14	1960	$\beta^-=100$
$^{85}\text{Br}$	-78575	3	2.90 m 0.06	3/2 <sup>-</sup>	14	1943	$\beta^-=100$
$^{85}\text{Kr}$	-81480.3	2.0	10.728 y 0.007	9/2 <sup>+</sup> *	14 FGK209 T	1940	$\beta^+=100$
$^{85}\text{Kr}^m$	-81175.4	2.0	304.871 0.020	1/2 <sup>-</sup> *	14 95Ke04 J	1937	$\beta^-=78.8.5; IT=21.2.5$
$^{85}\text{Kr}^n$	-79488.5	2.0	1991.8 0.2	(17/2 <sup>+</sup> )	14 11Ru.A T	1989	IT=100
$^{85}\text{Rb}$	-82167.341	0.005	STABLE	5/2 <sup>-</sup> *	14	1921	IS=72.17.2
$^{85}\text{Rb}^m$	-81653.335	0.005	514.0065 0.0022	9/2 <sup>+</sup> *	14 19Ta19 T	1964	IT=100
$^{85}\text{Sr}$	-81103.3	2.8	64.846 d 0.006	9/2 <sup>+</sup> *	14 FGK204 T	1940	$\epsilon=100$
$^{85}\text{Sr}^m$	-80864.5	2.8	238.79 0.05	1/2 <sup>-</sup> *	14	1940	IT=86.6.4; $\beta^+=13.4.4$
$^{85}\text{Y}$	-77842	19	2.68 h 0.05	(1/2 <sup>-</sup> )	14	1952	$\beta^+=100$
$^{85}\text{Y}^m$	-77822	19	19.68 0.17	(9/2 <sup>+</sup> )	14	1952	$\beta^+\approx 100; IT ?$
$^{85}\text{Y}^n$	-77576	19	266.18 0.10	(5/2 <sup>-</sup> )	14	1977	IT=100
$^{85}\text{Zr}$	-73175	6	7.86 m 0.04	(7/2 <sup>+</sup> )	14	1963	$\beta^+=100$
$^{85}\text{Zr}^m$	-72883	6	292.2 0.3	1/2 <sup>-</sup> #	14	1976	IT=?; $\beta^+=?$
$^{85}\text{Nb}$	-66280	4	20.5 s 0.7	9/2 <sup>+</sup> #	14	1988	$\beta^+=100$
$^{85}\text{Nb}^m$	-66130#	80#	150# 80#	(1/2 <sup>-</sup> )	14 05Ka39 J	1988	IT=?; $\beta^+=?$
$^{85}\text{Nb}^n$		<i>non - exist</i>			14 98Oi02 IT		$\beta^- ?; IT ?$
$^{85}\text{Mo}$	-57510	16	12 s 5				$\beta^+=100; \beta^+p=0.14.2$
$^{85}\text{Tc}$	-45850#	400#	3.2 s 0.2	(1/2 <sup>+</sup> )	14 05Xu04 J	1992	$\beta^+=100; \beta^+p=0.14.2$
$^{85}\text{Ru}$	-30630#	500#	<110ns	1/2 <sup>-</sup> #	14 00We.A I		p ?
$^{85}\text{Ru}$			1# ms >400ns	3/2 <sup>-</sup> #	15 13Su23 I	2013	$\beta^+ ?; \beta^+p ?; p ?$
$^{85}\text{Ga}$	D : % $\beta^-$ -n average 19Yo03=90(7)% 18Mi03=70(5)%						
$^{85}\text{Ge}$	D : % $\beta^-$ -n from 14Ag12; others 18Mi03=15(5) 91Kr15=14(3)						
$^{85}\text{Ge}$	T : average 14XuZZ=495(6) 13Ma22=494(8); others 91Kr15=535(47) 91Om01=580(50)						
$^{85}\text{As}$	D : % $\beta^-$ -n average 14Ag12=63.1(1.0) 93Ru01=59.3(2.5)						
$^{85}\text{As}$	T : average 13Ma22=2.08(0.14) 93Ru01=2.002(0.013) 91Kr15=2.032(0.012)						
$^{85}\text{As}$	T : 68To19=2.028(0.012); others 73Kr06=2.05(0.05), superseded by 91Kr15						
$^{85}\text{As}$	T : 76Ru01=2.08(0.05), superseded by 93Ru01 91Om01=2.0(0.1) 78Cr03=1.9(0.1)						
$^{85}\text{Kr}$	J : also 95Ke04=9/2						
$^{85}\text{Rb}$	J : also 14Ya28=2.5(1), 81Th04=5/2						
$^{85}\text{Rb}^m$	T : average 19Ta19=1.0202(0.0060) 72Mi23=1.015(0.001)						
$^{85}\text{Sr}$	J : 90Li28=9/2						
$^{85}\text{Sr}^m$	J : 90Li28=1/2						
$^{85}\text{Nb}$	T : average 05Ka39=17(2) 88Ku14=20.9(0.7)						



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
* <sup>85</sup> Nb <sup>m</sup>	E : from 05Ka39 > 69 keV						**	
* <sup>85</sup> Nb <sup>n</sup>	I : activity reported in 98Oi02 and adopted in Ensdf2014 as a new isomer;						**	
* <sup>85</sup> Nb <sup>n</sup>	I : not confirmed in 05Ka39 (the same laboratory as 98Oi02)						**	
* <sup>85</sup> Tc	I : also 99Ja02<100 ns; estimated T1/2=100# ms for $\beta^+$ decay						**	
<sup>86</sup> Zn	-20060#	500#		0 <sup>+</sup>			$\beta^-$ ?; $\beta^-n$ ?	
<sup>86</sup> Ga	-33760#	400#	49 ms 2		15 19Yo03	TD 1997	$\beta^-$ =100; $\beta^-n$ =69 6; $\beta^-2n$ =16.2 11	
<sup>86</sup> Ge	-49400	440	221.6 ms 11	0 <sup>+</sup>	15 14XuZZ	T 1994	$\beta^-$ =100; $\beta^-n$ =45 15	
<sup>86</sup> As	-58962	3	945 ms 8	(1 <sup>-</sup> , 2 <sup>-</sup> )	15 15Ma61	J 1973	$\beta^-$ =100; $\beta^-n$ =35.5 6; $\beta^-2n$ ?	
<sup>86</sup> Se	-70503.2	2.5	14.3 s 0.3	0 <sup>+</sup>	16	1973	$\beta^-$ =100; $\beta^-n$ ?	
<sup>86</sup> Br	-75632	3	55.1 s 0.4	(1 <sup>-</sup> )	15	1962	$\beta^-$ =100	
<sup>86</sup> Kr	-83265.676	0.004	STABLE	0 <sup>+</sup>	15	1920	IS=17.279 41; 2 $\beta^-$ ?	
<sup>86</sup> Rb	-82747.00	0.20	18.645 d 0.008	2 <sup>-*</sup>	15	1941	$\beta^-$ ≈100; $\epsilon$ =0.0052 5	
<sup>86</sup> Rb <sup>m</sup>	-82190.95	0.27	556.05 0.18	1.017 m 0.003	6 <sup>-*</sup>	15	1951	IT≈100; $\beta^-$ <0.3
<sup>86</sup> Sr	-84523.100	0.005	STABLE	0 <sup>+</sup>	15	1931	IS=9.86 20	
<sup>86</sup> Sr <sup>m</sup>	-81567.01	0.12	2956.09 0.12	455 ns 7	8 <sup>+</sup>	15	1971	IT=100
<sup>86</sup> Y	-79283	14	14.74 h 0.02	4 <sup>-*</sup>	15	1951	$\beta^+$ =100	
<sup>86</sup> Y <sup>m</sup>	-79065	14	218.21 0.09	47.4 m 0.4	(8 <sup>+</sup> )	15	1962	IT=99.31 4; $\beta^+$ =0.69 4
<sup>86</sup> Y <sup>n</sup>	-78981	14	302.18 0.09	125.3 ns 5.5	6 <sup>+</sup>	15 10Ru07	J 2000	IT=100
<sup>86</sup> Zr	-77969	4	16.5 h 0.1	0 <sup>+</sup>	15	1951	$\beta^+$ =100	
<sup>86</sup> Nb	-69134	5	*	88 s 1	(6 <sup>+</sup> )	15	1974	$\beta^+$ =100
<sup>86</sup> Nb <sup>m</sup>	-68980#	100#	150# 100#	20# s	(0 <sup>-</sup> , 1 <sup>-</sup> , 2 <sup>-</sup> )	15 05Ka39	J 1994	$\beta^+$ =100; IT ?
<sup>86</sup> Nb <sup>n</sup>		non-exist	RN	56.3 s 8.3		15 94Sh07	IT	$\beta^+$ = ?; IT ?
<sup>86</sup> Mo	-64110.9	2.9	19.1 s 0.3	0 <sup>+</sup>	15	1991	$\beta^+$ =100	
<sup>86</sup> Tc	-51570#	300#	55 ms 7	(0 <sup>+</sup> )	15	1992	$\beta^+$ =100; $\beta^+p$ ?	
<sup>86</sup> Tc <sup>m</sup>	-50050#	300#	1524 10	1.10 $\mu$ s 0.12	(6 <sup>+</sup> )	15 08Ga04	T 2000	IT=100
<sup>86</sup> Ru	-39770#	400#	50# ms >400ns	0 <sup>+</sup>	15 13Su23	I 2013	$\beta^+$ ?; $\beta^+p$ ?	
* <sup>86</sup> Ga	D : % $\beta^-n$ average 19Yo03=74(8) 13Mi19=60(10)						**	
* <sup>86</sup> Ge	T : other 13Ma22=226(21), supersedes 12Ma.A=219(40)						**	
* <sup>86</sup> Rb	J : also 14Ya28=1.9(2), 81Th04=2						**	
* <sup>86</sup> Rb	T : average 16Ma49=18.648(0.009) 81Mi10=18.631(0.018)						**	
* <sup>86</sup> Y	J : 07Ch07=4						**	
* <sup>86</sup> Y <sup>n</sup>	T : average 10Ru07=127(14) 00Io02=125(6)						**	
* <sup>86</sup> Nb <sup>m</sup>	I : from 94Sh07 and 05Ka39, populated in $\beta^+$ decay of <sup>86</sup> Mo (0 <sup>+</sup> )						**	
* <sup>86</sup> Nb <sup>n</sup>	I : half-life deduced in 94Sh07 by gating on Zr X-rays, which would be						**	
* <sup>86</sup> Nb <sup>n</sup>	I : consistent with decay of two isomers, one with T1/2=88 s and the						**	
* <sup>86</sup> Nb <sup>n</sup>	I : second with a half-life similar to that of <sup>86</sup> Mo, T1/2=19.1 s;						**	
* <sup>86</sup> Nb <sup>n</sup>	I : not confirmed in 05Ka39, 97Ta10 (sensitive to high-spin structures)						**	
* <sup>86</sup> Tc <sup>m</sup>	T : average 08Ga04=1.10(0.14) 00Ch07=1.11(0.21)						**	
* <sup>86</sup> Tc <sup>m</sup>	E : uncertainty estimated by GAU						**	
<sup>87</sup> Ga	-28870#	500#	29 ms 4	5/2 <sup>-</sup> #	15 19Yo03	TD 2010	$\beta^-$ =100; $\beta^-n$ =81 12; $\beta^-2n$ =10.2 2.8	
<sup>87</sup> Ge	-43590#	300#	103 ms 4	5/2 <sup>+</sup> #	15 14XuZZ	T 1997	$\beta^-$ =100; $\beta^-n$ ?; $\beta^-2n$ ?	
<sup>87</sup> As	-55617.9	3.0	492 ms 25	(5/2 <sup>-</sup> , 3/2 <sup>-</sup> )	15 15Ko19	TJ 1970	$\beta^-$ =100; $\beta^-n$ =15.4 22; $\beta^-2n$ ?	
<sup>87</sup> Se	-66426.1	2.2	5.50 s 0.06	(3/2 <sup>+</sup> )	15 15Ko19	J 1968	$\beta^-$ =100; $\beta^-n$ =0.60 12	
<sup>87</sup> Br	-73892	3	55.68 s 0.12	5/2 <sup>-</sup>	15 19Wi11	J 1943	$\beta^-$ =100; $\beta^-n$ =2.60 4	
<sup>87</sup> Kr	-80709.53	0.25	76.3 m 0.5	5/2 <sup>+</sup> *	15	1940	$\beta^-$ =100	
<sup>87</sup> Rb	-84597.802	0.006	49.7 Gy 0.3	3/2 <sup>-*</sup>	15	1921	IS=27.83 2; $\beta^-$ =100	
<sup>87</sup> Sr	-84880.076	0.005	STABLE	9/2 <sup>+</sup> *	15	1931	IS=7.00 20	
<sup>87</sup> Sr <sup>m</sup>	-84491.547	0.006	388.5287 0.0023	2.805 h 0.009	1/2 <sup>-*</sup>	15 21Kr.A	T 1940	IT=99.70 8; $\epsilon$ =0.30 8
<sup>87</sup> Y	-83018.4	1.1	79.8 h 0.3	1/2 <sup>-*</sup>	15	1940	$\beta^+$ =100	
<sup>87</sup> Y <sup>m</sup>	-82637.6	1.1	380.82 0.07	13.37 h 0.03	9/2 <sup>+</sup> *	15	1940	IT=98.43 11; $\beta^+$ =1.57 11
<sup>87</sup> Zr	-79347	4	1.68 h 0.01	9/2 <sup>+</sup>	15	1948	$\beta^+$ =100	
<sup>87</sup> Zr <sup>m</sup>	-79011	4	335.84 0.19	14.0 s 0.2	1/2 <sup>-</sup>	15	1972	IT=100
<sup>87</sup> Nb	-73874	7	3.7 m 0.1	(1/2 <sup>-</sup> )	15	1971	$\beta^+$ =100	
<sup>87</sup> Nb <sup>m</sup>	-73870	7	3.9 0.1	2.6 m 0.1	(9/2 <sup>+</sup> )	15	1972	$\beta^+$ =100
<sup>87</sup> Mo	-66884.8	2.9	14.1 s 0.3	7/2 <sup>+</sup> #	15	1977	$\beta^+$ =100; $\beta^+p$ =15 5	
<sup>87</sup> Tc	-57690	4	*	2.14 s 0.17	9/2 <sup>+</sup> #	15 19Pa16	TD 1991	$\beta^+$ =100; $\beta^+p$ <0.7

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)				
$^{87}\text{Tc}^m$	-57683	4	7	1	*	2# s	1/2 <sup>-</sup> #	09Ga40 E	$\beta^+$ ?; IT ?	*		
$^{87}\text{Tc}^n$	-57619	4	71	1		647 ns	24	7/2 <sup>+</sup> #	15	2007	IT=100	
$^{87}\text{Ru}$	-45730#	400#				50# ms	>1.5us	1/2 <sup>-</sup> #	15	95Ry03 I	1995	$\beta^+$ ?; $\beta^+$ p ?
* $^{87}\text{As}$	T : average 15Ko19=560(80) 13Ma22=484(35) 93Ru01=485(40); others									**		
* $^{87}\text{As}$	T : 12Qu01=1450(550)(+3900-1100) 78Cr03=730(60)									**		
* $^{87}\text{Se}$	T : average 93Ru01=5.29(11) 70Kr05=5.85(15) 70De08=5.90(20) 71To13=5.41(10)									**		
* $^{87}\text{Se}$	D : % $\beta^-$ n from 93Ru01									**		
* $^{87}\text{Rb}$	J : also 14Ya28=1.53(6)									**		
* $^{87}\text{Sr}$	J : 90Li28=9/2									**		
* $^{87}\text{Sr}^m$	J : 90Li28=1/2									**		
* $^{87}\text{Sr}^m$	T : average 21Kr.A=2.808(0.003) 97We13=2.811(0.027) 82Gr07=2.795(0.013)									**		
* $^{87}\text{Sr}^m$	T : 70Le07=2.793(0.009) 68Go30=3.805(0.001); other 92An19=2.827(0.001),									**		
* $^{87}\text{Sr}^m$	T : discrepant (not used)									**		
* $^{87}\text{Y}$	J : 07Ch07=1/2									**		
* $^{87}\text{Y}^m$	J : 07Ch07=9/2									**		
* $^{87}\text{Tc}$	T : average 19Pa16=2.0(0.3) 01Ki13=2.2(0.2)									**		
* $^{87}\text{Tc}^m$	E : 64 keV gamma ray observed in parallel to the 71 keV one, depopulating									**		
* $^{87}\text{Tc}^m$	E : $^{87}\text{Tc}^n$									**		
$^{88}\text{Ga}$	-22390#	500#										$\beta^-$ ?; $\beta^-$ n ?
$^{88}\text{Ge}$	-39520#	400#				61 ms	6	0 <sup>+</sup>	14	14XuZZ T	1997	$\beta^-$ =100; $\beta^-$ n ?; $\beta^-$ 2n ?
$^{88}\text{As}$	-50450#	200#				270 ms	150		14	12Qu01 T	1994	$\beta^-$ =100; $\beta^-$ n ?
$^{88}\text{Se}$	-63884	3				1.53 s	0.06	0 <sup>+</sup>	14		1970	$\beta^-$ =100; $\beta^-$ n=0.99 10
$^{88}\text{Br}$	-70716	3				16.34 s	0.08	(1 <sup>-</sup> )	14	15Cz01 J	1948	$\beta^-$ =100; $\beta^-$ n=6.58 18
$^{88}\text{Br}^m$	-70446	3	270.17	0.11		5.51 $\mu$ s	0.04	(4 <sup>-</sup> )	14	11Ru.A T	1970	IT=100
$^{88}\text{Kr}$	-79691.3	2.6				2.825 h	0.019	0 <sup>+</sup>	14		1939	$\beta^-$ =100
$^{88}\text{Rb}$	-82609.00	0.16				17.78 m	0.03	2 <sup>-</sup> *	14	20Ch42 T	1939	$\beta^-$ =100
$^{88}\text{Rb}^m$	-81235.2	0.3	1373.8	0.3		123 ns	13	(7 <sup>+</sup> )	14		2000	IT=100
$^{88}\text{Sr}$	-87921.629	0.006				STABLE		0 <sup>+</sup>	14		1923	IS=82.58 35
$^{88}\text{Y}$	-84299.0	1.5				106.629 d	0.024	4 <sup>-</sup> *	14	FGK204 T	1948	$\beta^+$ =100
$^{88}\text{Y}^m$	-83906.1	1.5	392.86	0.09		301 $\mu$ s	3	1 <sup>+</sup>	14		1955	IT=100
$^{88}\text{Y}^n$	-83624.5	1.5	674.55	0.04		13.98 ms	0.17	8 <sup>+</sup> *	14		1962	IT=100
$^{88}\text{Zr}$	-83629	5				83.4 d	0.3	0 <sup>+</sup>	14		1951	$\epsilon$ =100
$^{88}\text{Zr}^m$	-80741	5	2887.79	0.06		1.320 $\mu$ s	0.025	8 <sup>+</sup>	14		1978	IT=100
$^{88}\text{Nb}$	-76170	60			*	14.50 m	0.11	(8 <sup>+</sup> )	14		1964	$\beta^+$ =100
$^{88}\text{Nb}^m$	-76040	100	130	120	BD*	7.7 m	0.1	(4 <sup>-</sup> )	14		1971	$\beta^+$ =100
$^{88}\text{Mo}$	-72687	4				8.0 m	0.2	0 <sup>+</sup>	14		1971	$\beta^+$ =100
$^{88}\text{Tc}$	-61670	4				6.4 s	0.8	(2 <sup>+</sup> )	14	09Ga40 J	1991	$\beta^+$ =100; $\beta^+$ p ?
$^{88}\text{Tc}^m$	-61600	5	70	3	MD	5.8 s	0.2	(6 <sup>+</sup> )	14	19Vi05 J	1993	$\beta^+$ =100; $\beta^+$ p ?
$^{88}\text{Tc}^n$	-61575	4	95	1		146 ns	12	(4 <sup>+</sup> )	14	09Ga40 TJ	2009	IT=100
$^{88}\text{Ru}$	-54340#	300#				1.5 s	0.3	0 <sup>+</sup>	14	19Pa16 TD	1994	$\beta^+$ =100; $\beta^+$ p<3.6
$^{88}\text{Rh}$	-36860#	400#				1# ms						$\beta^+$ ?
* $^{88}\text{As}$	T : symmetrized from 12Qu01=200(5)(+200-90)									**		
* $^{88}\text{Br}^m$	J : 15Cz01=(4-)									**		
* $^{88}\text{Br}^m$	T : also 18Rz01=5.5(0.1)									**		
* $^{88}\text{Rb}$	J : also 81Th04=2									**		
* $^{88}\text{Rb}$	T : average 20Ch42=17.78(0.05) 89Ab22=17.773(0.033), uncertainty increased									**		
* $^{88}\text{Rb}$	T : to 3 $\sigma$ by evaluator, 69Ra05=17.78(0.11); other 69He16=17.7(0.1)									**		
* $^{88}\text{Y}$	J : 07Ch07=4									**		
* $^{88}\text{Y}^n$	J : 07Ch07=8									**		
* $^{88}\text{Zr}^m$	T : other 17Pa35=1.40(0.07)									**		
* $^{88}\text{Ru}$	T : average 19Pa16=1.9(0.5) 01Ki13=1.2(+0.3-0.2)									**		
$^{89}\text{Ge}$	-33040#	400#				60# ms	>300ns	3/2 <sup>+</sup> #	13		1997	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?
$^{89}\text{As}$	-46530#	300#				220# ms	>150ns	5/2 <sup>-</sup> #	13	94Be24 I	1994	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?
$^{89}\text{Se}$	-58992	4				430 ms	50	5/2 <sup>+</sup> #	13		1971	$\beta^-$ =100; $\beta^-$ n=7.8 25
$^{89}\text{Br}$	-68274	3				4.357 s	0.022	(3/2 <sup>-</sup> , 5/2 <sup>-</sup> )	13		1959	$\beta^-$ =100; $\beta^-$ n=13.8 4
$^{89}\text{Kr}$	-76535.8	2.1				3.15 m	0.04	3/2 <sup>+</sup> *	13	95Ke04 J	1940	$\beta^-$ =100
$^{89}\text{Rb}$	-81712	5				15.32 m	0.10	3/2 <sup>-</sup> *	13		1940	$\beta^-$ =100
$^{89}\text{Sr}$	-86209.03	0.09				50.563 d	0.025	5/2 <sup>+</sup> *	13		1937	$\beta^-$ =100
$^{89}\text{Y}$	-87711.2	0.3				STABLE		1/2 <sup>-</sup> *	13		1923	IS=100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{89}\text{Y}^m$	-86802.2	0.3	908.97	0.03	15.663 s 0.005	9/2+*	13 94It.A T	1951	IT=100	*
$^{89}\text{Zr}$	-84878.0	2.8			78.360 h 0.023	9/2+	13 20Fe10 T	1948	$\beta^+=100$	*
$^{89}\text{Zr}^m$	-84290.2	2.8	587.82	0.10	4.161 m 0.010	1/2-	13	1953	IT=93.77 12; $\beta^+=6.23$ 12	
$^{89}\text{Nb}$	-80626	24			2.03 h 0.07	(9/2+)	13	1954	$\beta^+=100$	
$^{89}\text{Nb}^m$	-80630#	40#	0#	30#	1.10 h 0.03	(1/2)-	13	1954	$\beta^+=100$	
$^{89}\text{Mo}$	-75015	4			2.11 m 0.10	(9/2+)	13	1980	$\beta^+=100$	
$^{89}\text{Mo}^m$	-74628	4	387.5	0.2	190 ms 15	(1/2-)	13	1980	IT=100	
$^{89}\text{Tc}$	-67395	4			12.8 s 0.9	(9/2+)	13	1991	$\beta^+=100$	
$^{89}\text{Tc}^m$	-67332	4	62.6	0.5	12.9 s 0.8	(1/2-)	13	1991	$\beta^+\approx 100$ ;IT ?	
$^{89}\text{Ru}$	-58369	24			1.32 s 0.03	(9/2+)	13 19Pa16 TD	1992	$\beta^+=100$ ; $\beta^+p=3.1$ 2	
$^{89}\text{Rh}$	-45650#	360#			<120ns	9/2+#	16 16Ce02 TI		$\beta^+ ?$ ; $\beta^+p ?$ ; $p ?$	
* $^{89}\text{Rb}$	J : also 81Th04=3/2									**
* $^{89}\text{Y}$	J : 07Ch07=1/2									**
* $^{89}\text{Y}^m$	J : 07Ch07=9/2									**
* $^{89}\text{Zr}$	T : average 20Fe10=78.368(0.032) 18Ga04=78.333(0.038) 64Va03=78.43(0.08)									**
$^{90}\text{Ge}$	-28470#	500#			30# ms >400ns	0+	20 100h02 I	2010	$\beta^- ?$ ; $\beta^-n ?$ ; $\beta^-2n ?$	
$^{90}\text{As}$	-40990#	400#			70# ms >300ns		20 97Be70 I	1997	$\beta^- ?$ ; $\beta^-n ?$ ; $\beta^-2n ?$	
$^{90}\text{As}^m$	-40870#	400#	124.5	0.5	220 ns 100		12Ka36 ET	2012	IT=100	*
$^{90}\text{Se}$	-55800	330			210 ms 80	0+	20 12Qu01 T	1994	$\beta^+=100$ ; $\beta^-n ?$	*
$^{90}\text{Br}$	-64000	3			1.910 s 0.010		20	1959	$\beta^-=100$ ; $\beta^-n=25.3$ 15	
$^{90}\text{Kr}$	-74959.3	1.9			32.32 s 0.09	0+	20	1951	$\beta^-=100$	
$^{90}\text{Rb}$	-79366	6			158 s 5	0*	20	1951	$\beta^-=100$	*
$^{90}\text{Rb}^m$	-79259	6	106.90	0.03	258 s 4	3*	20	1967	$\beta^-=97.4$ 4;IT=2.5 4	*
$^{90}\text{Rb}^x$	-79295	14	71	12	R = 2 1	fsmix				
$^{90}\text{Sr}$	-85950.9	1.4			28.91 y 0.03	0+	20	1948	$\beta^-=100$	
$^{90}\text{Y}$	-86496.9	0.4			64.05 h 0.05	2*-	20	1937	$\beta^-=100$	*
$^{90}\text{Y}^m$	-85814.9	0.4	682.01	0.05	3.226 h 0.011	7+*	20 20Kr06 T	1961	IT=99.9982 2; $\beta^-=0.0018$ 2	*
$^{90}\text{Zr}$	-88772.55	0.12			STABLE	0+	20	1924	IS=51.45 4	
$^{90}\text{Zr}^m$	-86453.55	0.12	2319.000	0.009	809.2 ms 2.0	5-	20	1972	IT=100	
$^{90}\text{Zr}^n$	-85183.13	0.12	3589.418	0.015	131 ns 4	8+	20	1977	IT=100	
$^{90}\text{Nb}$	-82662	3			14.60 h 0.05	8+*	20	1951	$\beta^+=100$	*
$^{90}\text{Nb}^m$	-82540	3	122.370	0.022	63 $\mu$ s 2	6+	20	1967	IT=100	
$^{90}\text{Nb}^n$	-82537	3	124.67	0.25	18.81 s 0.06	4*-	20	1969	IT=100	*
$^{90}\text{Nb}^p$	-82491	3	171.10	0.10	< 1 $\mu$ s	7+	20	1981	IT=100	
$^{90}\text{Nb}^q$	-82280	3	382.01	0.25	6.19 ms 0.08	1+	20	1967	IT=100[gs=0,m=100]	
$^{90}\text{Nb}^r$	-80782	3	1880.21	0.20	471 ns 6	(11-)	20 05Ch65 TJ	1978	IT=100	*
$^{90}\text{Mo}$	-80173	3			5.56 h 0.09	0+	20	1953	$\beta^+=100$	
$^{90}\text{Mo}^m$	-77298	3	2874.73	0.15	1.14 $\mu$ s 0.05	8+	20	1971	IT=100	
$^{90}\text{Tc}$	-70724.7	1.0			49.2 s 0.4	(8+)	20	1974	$\beta^+=100$	
$^{90}\text{Tc}^m$	-70580.7	1.3	144.0	1.7 MD	8.7 s 0.2	1+	20	1974	$\beta^+=100$	
$^{90}\text{Ru}$	-64884	4			11.7 s 0.9	0+	20	1991	$\beta^+=100$	
$^{90}\text{Rh}$	-51630#	200#			29 ms 3	(0+)	20 19Pa16 JTD	1994	$\beta^+=100$ ; $\beta^+p<0.7$	
$^{90}\text{Rh}^m$	-51630#	540#	0#	500#	0.56 s 0.02	(7+)	20 19Pa16 JTD	2001	$\beta^+=100$ ; $\beta^+p=9.6$ 10	
$^{90}\text{Pd}$	-39710#	400#			10# ms >400ns	0+	20 16Ce02 I	2016	$\beta^+ ?$ ; $\beta^+p ?$ 2p ?	
* $^{90}\text{As}^m$	T : symmetrized from 12Ka36=200(+120-90)									**
* $^{90}\text{Se}$	T : symmetrized from 12Qu01=195(7,stat)(+95-65,syst)									**
* $^{90}\text{Rb}$	J : also 81Th04=0									**
* $^{90}\text{Rb}^m$	J : also 81Th04=3									**
* $^{90}\text{Y}$	J : also 07Ch07,78Fu06=2									**
* $^{90}\text{Y}^m$	J : 07Ch07=7									**
* $^{90}\text{Y}^m$	T : average 20Kr06=3.178(0.012) 92An19=3.244(0.005) 67Gr02=3.19(0.01)									**
* $^{90}\text{Y}^m$	T : 62Ab03=3.15(0.05) 61Ca12=3.2(0.1) 61He09=3.14(0.10) 61Ha17=3.19(0.06);									**
* $^{90}\text{Y}^m$	T : Birge ratio=2.75									**
* $^{90}\text{Nb}$	J : also 09Ch25=8									**
* $^{90}\text{Nb}^n$	J : 09Ch25=4									**
* $^{90}\text{Nb}^r$	T : average 05Ch65=470(10) 81Fi02=440(20) 78Ha52=477(8); other									**
* $^{90}\text{Nb}^r$	T : 17Pa35=415(67)									**
$^{91}\text{As}$	-36500#	400#			100# ms >300ns	5/2-#	13 97Be70 I	1997	$\beta^- ?$ ; $\beta^-n ?$ ; $\beta^-2n ?$	
$^{91}\text{Se}$	-50580	430			270 ms 50	1/2+#	13	1975	$\beta^-=100$ ; $\beta^-n=21$ 10; $\beta^-2n ?$	

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
<sup>91</sup> Br	-61107	4		543 ms	4	13 14Ag12	D 1974	$\beta^- = 100; \beta^-_n = 29.5$
<sup>91</sup> Kr	-70974.0	2.2		8.57 s	0.04	13	1951	$\beta^- = 100; \beta^-_n ?$
<sup>91</sup> Rb	-77745	8		58.2 s	0.3	13	1951	$\beta^- = 100; \beta^-_n ?$
<sup>91</sup> Sr	-83652	5		9.65 h	0.06	13 19Kr10	T 1943	$\beta^- = 100$
<sup>91</sup> Y	-86351.3	1.8		58.51 d	0.06	13	1943	$\beta^- = 100$
<sup>91</sup> Y <sup>m</sup>	-85795.7	1.8	555.58	49.71 m	0.04	13	1953	IT $\approx$ 100; $\beta^- ?$
<sup>91</sup> Zr	-87895.59	0.09		STABLE		13	1934	IS=11.22
<sup>91</sup> Zr <sup>m</sup>	-84728.3	0.4	3167.3	4.35 $\mu$ s	0.14	13	1985	IT=100
<sup>91</sup> Nb	-86638.0	2.9		680 y	130	13 91Hi.A	D 1951	$\epsilon \approx 100; e^+ = 0.0138$
<sup>91</sup> Nb <sup>m</sup>	-86533.4	2.9	104.60	60.86 d	0.22	13 91Hi.A	D 1950	IT=96.6 5; $\epsilon = 3.4$ 5; $e^+ = 0.0028$ 2
<sup>91</sup> Nb <sup>n</sup>	-84603.6	2.9	2034.42	3.76 $\mu$ s	0.12	13	1974	IT=100
<sup>91</sup> Mo	-82209	6		15.49 m	0.01	13	1948	$\beta^+ = 100$
<sup>91</sup> Mo <sup>m</sup>	-81556	6	653.01	64.6 s	0.6	13	1953	IT=50.0 16; $\beta^+ = 50.0$ 16
<sup>91</sup> Tc	-75986.7	2.4		3.14 m	0.02	13	1974	$\beta^+ = 100$
<sup>91</sup> Tc <sup>m</sup>	-75847.4	2.4	139.3	3.3 m	0.1	13	1975	$\beta^+ \approx 100; IT ?$
<sup>91</sup> Ru	-68239.8	2.2		8.0 s	0.4	13	1983	$\beta^+ = 100; \beta^+ p ?$
<sup>91</sup> Ru <sup>m</sup>	-68580	500	-340	7.6 s	0.8	13	1983	$\beta^+ \approx 100; \beta^+ p = ?; IT ?$
<sup>91</sup> Rh	-58570#	300#		1.47 s	0.22	13 04De40	TJ 1994	$\beta^+ = 100; \beta^+ p = 1.3$ 5
<sup>91</sup> Rh <sup>m</sup>	-58400#	300#	172.9	1.8# s		13	2004	$\beta^+ ?; IT ?; \beta^+ p ?$
<sup>91</sup> Pd	-46170#	420#		32 ms	3	13 18Pa20	TD 1995	$\beta^+ = 100; \beta^+ p = 3.1$ 10
* <sup>91</sup> Rb	J : other 81Th04=3/2							
* <sup>91</sup> Sr	J : 90Li28=5/2							
* <sup>91</sup> Sr	T : other 19Kr10=9.66(0.09)							
* <sup>91</sup> Zr	J : 02Ca37=5/2							
* <sup>91</sup> Nb	J : 09Ch25=9/2							
* <sup>91</sup> Nb <sup>m</sup>	J : 09Ch25=1/2							
* <sup>91</sup> Rh	T : from 04De40=1.47(0.22) using time spectra gated by gamma rays feeding							
* <sup>91</sup> Rh	T : the <sup>91</sup> Ru gs (9/2+); others: 19Pa16=1.60(0.02) 01Ki13=1.7(0.2)							
* <sup>91</sup> Rh	T : 00We.A=1.74(0.14) (same group as 01Ki13) probably include both gs							
* <sup>91</sup> Rh	T : and isomer							
* <sup>91</sup> Rh <sup>m</sup>	T : Ensdf2013 assign T1/2=1.47(0.22) from 04De40, but this value is							
* <sup>91</sup> Rh <sup>m</sup>	T : unambiguously associated in 04De04 with the decay of the 9/2+ gs.							
* <sup>91</sup> Rh <sup>m</sup>	T : 19Pa16=1.60(0.02) 01Ki13=1.7(0.2) 00We.A=1.74(0.14) probably include							
* <sup>91</sup> Rh <sup>m</sup>	T : both gs and isomer							
* <sup>91</sup> Pd	D : % $\beta^+ p$ symmetrized from 18Pa20=3.0(+1.1-0.9)							
<sup>92</sup> As	-30380#	500#		45# ms	>300ns	12 97Be70	I 1997	$\beta^- ?; \beta^-_n ?; \beta^-_n ?$
<sup>92</sup> Se	-46720#	400#		90# ms	>300ns	12 97Be70	I 1997	$\beta^- ?; \beta^-_n ?; \beta^-_n ?$
<sup>92</sup> Se <sup>m</sup>	-44780#	400#	3072	15.7 $\mu$ s	0.7	20Li15	ETJ 2012	IT=100
<sup>92</sup> Br	-56233	7		314 ms	16	12	1974	$\beta^- = 100; \beta^-_n = 33.1$ 25; $\beta^-_n ?$
<sup>92</sup> Br <sup>m</sup>	-55571	7	662	88 ns	8	12Ka36	ET 2012	IT=100
<sup>92</sup> Br <sup>n</sup>	-55095	7	1138	85 ns	10	12Ka36	ET 2012	IT=100
<sup>92</sup> Kr	-68769.3	2.7		1.840 s	0.008	12	1951	$\beta^- = 100; \beta^-_n = 0.0332$ 25
<sup>92</sup> Rb	-74772	6		4.48 s	0.03	12	1960	$\beta^- = 100; \beta^-_n = 0.0107$ 5
<sup>92</sup> Sr	-82867	3		2.611 h	0.017	12	1956	$\beta^- = 100$
<sup>92</sup> Y	-84816	9		3.54 h	0.01	12	1940	$\beta^- = 100$
<sup>92</sup> Y <sup>m</sup>	-84010#	50#	807#	3.7 $\mu$ s	0.5	12 11Ru.A	ET 2009	IT=100
<sup>92</sup> Zr	-88459.02	0.09		STABLE		12	1924	IS=17.15 3
<sup>92</sup> Nb	-86453.3	1.8		34.7 My	2.4	12	1938	$\beta^+ = 100$
<sup>92</sup> Nb <sup>m</sup>	-86317.8	1.8	135.5	10.116 d	0.013	12 19Kr13	T 1959	$\beta^+ = 100$
<sup>92</sup> Nb <sup>n</sup>	-86227.5	1.8	225.8	5.9 $\mu$ s	0.2	12	1958	IT=100
<sup>92</sup> Nb <sup>p</sup>	-84250.0	1.8	2203.3	167 ns	4	12	1989	IT=100
<sup>92</sup> Mo	-86808.59	0.16		STABLE	>190Ey	12 97Ba35	T 1930	IS=14.649 106; $2\beta^+ ?$
<sup>92</sup> Mo <sup>m</sup>	-84048.07	0.21	2760.52	190 ns	3	12	1964	IT=100
<sup>92</sup> Tc	-78926	3		4.25 m	0.15	12	1964	$\beta^+ = 100$
<sup>92</sup> Tc <sup>m</sup>	-78656	3	270.09	1.03 $\mu$ s	0.06	12 17Pa35	T 1976	IT=100
<sup>92</sup> Tc <sup>n</sup>	-78397	3	529.42	< 0.1 $\mu$ s		12	1976	IT=100
<sup>92</sup> Tc <sup>p</sup>	-78215	3	711.33	< 0.1 $\mu$ s		12	1976	IT=100
<sup>92</sup> Ru	-74301.2	2.7		3.65 m	0.05	12	1971	$\beta^+ = 100$
<sup>92</sup> Ru <sup>m</sup>	-71467	3	2833.9	100 ns	8	12 19Ha26	T 1980	IT=100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{92}\text{Rh}$	-62999	4	5.61 s 0.08	(6 <sup>+</sup> )	12 19Pa16	TD 1994	$\beta^+=100; \beta^+p=2.05$ 7 *	
$^{92}\text{Rh}^m$	-62950#	100#	3.18 s 0.22	(2 <sup>+</sup> )	12 19Pa16	TD 2004	$\beta^+=100; \beta^+p=1.7$ 3 *	
$^{92}\text{Rh}^n$	-62890#	100#	232 ns 15	(4 <sup>+</sup> )	17Pa35	ETJ 2017	IT=100 *	
$^{92}\text{Pd}$	-54780	350	1.06 s 0.03	0 <sup>+</sup>	12 19Pa16	TD 1994	$\beta^+=100; \beta^+p=1.6$ 2 *	
$^{92}\text{Ag}$	-37530#	400#	1# ms >400ns		16 16Ce02	I 2016	$\beta^+ ?; p ?$ *	
$^{*92}\text{Se}^m$	T : other 12Ka36=10.3(+5.5-2.8) **							
$^{*92}\text{Se}^m$	E : uncertainty estimated by NuBase **							
$^{*92}\text{Br}^m$	T : symmetrized from 12Ka36=89(+7-8) **							
$^{*92}\text{Br}^m$	E : uncertainty estimated by NuBase **							
$^{*92}\text{Br}^n$	T : symmetrized from 12Ka36=84(+10-9) **							
$^{*92}\text{Br}^n$	E : uncertainty estimated by NuBase **							
$^{*92}\text{Rb}$	J : also 81Th04=0 **							
$^{*92}\text{Sr}$	T : other (recent) 19Kr10=2.66(0.06) **							
$^{*92}\text{Y}$	J : 07Ch07=2 **							
$^{*92}\text{Y}^m$	T : average 11Ru.A=3.3(0.6) 09Fo05=4.2(+0.8-0.6) **							
$^{*92}\text{Y}^m$	E : observed 315-keV and 419-keV gamma rays in a cascade; low energy **							
$^{*92}\text{Y}^m$	E : transition may directly depopulate the isomer **							
$^{*92}\text{Nb}$	J : 09Ch25=7 **							
$^{*92}\text{Nb}^m$	T : average 19Kr13=10.07(0.02) 68Re04=10.14(0.03) 62Bu16=10.16(0.03) **							
$^{*92}\text{Nb}^m$	T : 59We30=10.15(0.03) **							
$^{*92}\text{Mo}^m$	T : other 17Pa35=200(37) **							
$^{*92}\text{Tc}^m$	T : average 17Pa35=1.02(0.17) 71Ho27=1.03(0.07) **							
$^{*92}\text{Ru}^m$	T : average 19Ha26=100(10) 80No06=100(14) **							
$^{*92}\text{Rh}$	D : % $\beta^+p$ average 19Pa16=2.2(0.1) 12Lo08=1.9(0.1) **							
$^{*92}\text{Rh}$	J : from 97Ka07; 01Xu05>4 **							
$^{*92}\text{Rh}^m$	T : also 04De40=0.53(0.37) **							
$^{*92}\text{Rh}^n$	E : 55.3(0.3) keV above the (2+) isomer **							
$^{*92}\text{Rh}^n$	T : from 19Ha26=232(15); other 17Pa35=230(60) **							
$^{93}\text{Se}$	-40860#	400#	130# ms >300ns	1/2 <sup>+</sup> #	11 97Be70	I 1997	$\beta^- ?; \beta^-n ?; \beta^-2n ?$ *	
$^{93}\text{Se}^m$	-40180#	400#	420 ns 100		12Ka36	ET 2012	IT=100 *	
$^{93}\text{Br}$	-52890	430	152 ms 8	5/2 <sup>-</sup> #	11 13Mi13	TD 1981	$\beta^-=100; \beta^-n=64$ 6; $\beta^-2n ?$ *	
$^{93}\text{Kr}$	-64136.0	2.5	1.287 s 0.010	1/2 <sup>+</sup> *	11 13Mi13	TD 1951	$\beta^-=100; \beta^-n=1.95$ 11 *	
$^{93}\text{Rb}$	-72620	8	5.84 s 0.02	5/2 <sup>-</sup> *	11	1960	$\beta^-=100; \beta^-n=1.39$ 7 *	
$^{93}\text{Rb}^m$	-68197	8	111 ns 11	(27/2 <sup>-</sup> )	11	2010	IT=100 *	
$^{93}\text{Sr}$	-80086	8	7.43 m 0.03	5/2 <sup>+</sup> *	11	1959	$\beta^-=100$ *	
$^{93}\text{Y}$	-84227	10	10.18 h 0.08	1/2 <sup>-</sup> *	11	1948	$\beta^-=100$ *	
$^{93}\text{Y}^m$	-83468	10	820 ms 40	9/2 <sup>+</sup> *	11	1974	IT=100 *	
$^{93}\text{Zr}$	-87122.0	0.5	1.61 My0.05	5/2 <sup>+</sup>	11	1950	$\beta^-=100$ *	
$^{93}\text{Nb}$	-87212.8	1.5	STABLE	9/2 <sup>+</sup> *	11	1932	IS=100 *	
$^{93}\text{Nb}^m$	-87182.0	1.5	16.12 y 0.12	1/2 <sup>-</sup>	11 20Ho10	E 1965	IT=100 *	
$^{93}\text{Nb}^n$	-79753	17	1.5 $\mu$ s 0.5	33/2 <sup>-</sup> #	11	2007	IT=100 *	
$^{93}\text{Mo}$	-86807.08	0.18	4.0 ky 0.8	5/2 <sup>+</sup>	11	1946	$\epsilon=100$ *	
$^{93}\text{Mo}^m$	-84382.13	0.18	2424.95 0.04	6.85 h 0.07	21/2 <sup>+</sup>	11	1950	IT=99.88 1; $\beta^+=0.12$ 1 *
$^{93}\text{Mo}^n$	-77112	17	9695 17	1.8 $\mu$ s 1.0	(39/2 <sup>-</sup> )	11 05Fu01	T 2005	IT=100 *
$^{93}\text{Tc}$	-83606.1	1.0	2.75 h 0.05	9/2 <sup>+</sup>	11	1948	$\beta^+=100$ *	
$^{93}\text{Tc}^m$	-83214.3	1.0	391.84 0.08	43.5 m 1.0	1/2 <sup>-</sup>	11	1939	IT=77.4 6; $\beta^+=22.6$ 6 *
$^{93}\text{Tc}^n$	-81420.9	1.0	2185.16 0.15	10.2 $\mu$ s 0.3	(17/2 <sup>-</sup> )	11	1973	IT=100 *
$^{93}\text{Ru}$	-77216.7	2.1	59.7 s 0.6	(9/2 <sup>+</sup> )	11	1972	$\beta^+=100$ *	
$^{93}\text{Ru}^m$	-76482.3	2.1	734.40 0.10	10.8 s 0.3	(1/2 <sup>-</sup> )	11	1983	$\beta^+=78.0$ 23; IT=22.0 23; $\beta^+p=0.027$ 5 *
$^{93}\text{Ru}^n$	-75134.2	2.3	2082.5 0.9	2.30 $\mu$ s 0.07	(21/2 <sup>+</sup> )	11 17Pa35	T 1983	IT=100 *
$^{93}\text{Rh}$	-69011.8	2.6	13.9 s 1.6	9/2 <sup>+</sup> #	11	1994	$\beta^+=100$ *	
$^{93}\text{Pd}$	-58980	370	1.17 s 0.02	(9/2 <sup>+</sup> )	11 19Pa16	TD 1994	$\beta^+=100; \beta^+p=7.4$ 2 *	
$^{93}\text{Ag}$	-46400#	400#	228 ns 16	9/2 <sup>+</sup> #	16 16Ce02	T 1994	$p=?; \beta^+ ?; \beta^+p ?$ *	
$^{*93}\text{Se}^m$	E : 12Ka36=208.3(0.5) and 469.9(0.5) gamma rays in cascade to gs **							
$^{*93}\text{Se}^m$	T : symmetrized from 12Ka36=390(+120-80) **							
$^{*93}\text{Br}$	D : % $\beta^-n$ average 13Mi13=53(+11-8) 01Lh01=68(7) **							
$^{*93}\text{Kr}$	T : average 13Mi13=1.298(0.054) 12Qu01=1.245(0.070) stat(0.030) syst **							
$^{*93}\text{Kr}$	T : 76Ru01=1.33(0.05) 75As04=1.27(0.02) 69Ca03=1.289(0.012) **							
$^{*93}\text{Kr}$	J : 95Ke04=1/2 **							
$^{*93}\text{Kr}$	D : % $\beta^-n$ other (recent) 13Mi13=1.9(+0.6-0.2) **							

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
* <sup>93</sup> Sr	J : 90Li28=5/2						**	
* <sup>93</sup> Y	J : 07Ch07=1/2						**	
* <sup>93</sup> Y <sup>m</sup>	J : 07Ch07=9/2						**	
* <sup>93</sup> Nb	J : also 09Ch25=9/2						**	
* <sup>93</sup> Nb <sup>n</sup>	E : from 7435.3(2.1)+x keV in Ensdf2011; x<50 keV assumed in Nubase						**	
* <sup>93</sup> Nb <sup>n</sup>	J : week population (non-yrast) in 07Wa45, feeding the 7435.3 keV 37/2-						**	
* <sup>93</sup> Nb <sup>n</sup>	J : level; given the Weisskopf T1/2 estimates and the measured T1/2,						**	
* <sup>93</sup> Nb <sup>n</sup>	J : the depopulating 50 keV gamma transition is most likely E2						**	
* <sup>93</sup> Mo <sup>n</sup>	E : from 9670.0(2.3)+x keV in Ensdf2011; x<50 keV assumed in Nubase						**	
* <sup>93</sup> Mo <sup>n</sup>	T : symmetrized from 05Fu01=1.1(+1.5-0.4)						**	
* <sup>93</sup> Tc <sup>n</sup>	T : also 19Ha26=10(1)						**	
* <sup>93</sup> Ru <sup>n</sup>	T : average 17Pa35=2.36(0.12) 09Ga40=2.7(0.2) 83Gr33=2.6(0.3)						**	
* <sup>93</sup> Ru <sup>n</sup>	T : 83Ko07=2.6(0.2) 78Br25=2.05(0.10)						**	
* <sup>93</sup> Ag	T : estimated from the time of flight and the assumption that the ratio						**	
* <sup>93</sup> Ag	T : between the number of identified nuclei with the same Tz is identical						**	
<sup>94</sup> Se	-36800# 500#		50# ms >300ns	0 <sup>+</sup>	06 97Be70 I	1997	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?	*
<sup>94</sup> Se <sup>m</sup>	-34370# 500#	2430.0 0.6	680 ns 50	(7 <sup>-</sup> )	20Li05 EJT	2020	IT=100	*
<sup>94</sup> Br	-47650# 200#		70 ms 20	2 <sup>-</sup> #	06	1981	$\beta^-$ =100; $\beta^-$ n=68 16; $\beta^-$ 2n ?	*
<sup>94</sup> Br <sup>m</sup>	-47360# 200#	294.6 0.5	530 ns 15		12Ka36 ET	2012	IT=100	*
<sup>94</sup> Kr	-61348 12		212 ms 4	0 <sup>+</sup>	11 16Mi18 T	1972	$\beta^-$ =100; $\beta^-$ n=1.11 7	*
<sup>94</sup> Rb	-68562.8 2.0		2.702 s 0.005	3 <sup>-</sup> *	11 11Go37 D	1961	$\beta^-$ =100; $\beta^-$ n=10.3 3	*
<sup>94</sup> Rb <sup>m</sup>	-68458.6 2.0	104.2 0.2	130 ns 15	(0 <sup>-</sup> )	16Mi18 ETJ	2016	IT=100	*
<sup>94</sup> Rb <sup>n</sup>	-66487.9 2.4	2074.9 1.4	107 ns 16	(10 <sup>-</sup> )	11	2008	IT=100	*
<sup>94</sup> Sr	-78845.7 1.7		75.3 s 0.2	0 <sup>+</sup>	11	1959	$\beta^-$ =100	*
<sup>94</sup> Y	-82351 6		18.7 m 0.1	2 <sup>-</sup> *	06	1948	$\beta^-$ =100	*
<sup>94</sup> Y <sup>m</sup>	-81149 6	1202.3 1.0	1.304 $\mu$ s 0.012	(5 <sup>+</sup> )	06 17Ki09 T	1999	IT=100	*
<sup>94</sup> Zr	-87269.33 0.16		STABLE >110Py	0 <sup>+</sup>	06 99Ar25 T	1924	IS=17.38 4; 2 $\beta^-$ ?	*
<sup>94</sup> Nb	-86369.1 1.5		20.4 ky 0.4	6 <sup>+</sup>	06 12He11 T	1938	$\beta^-$ =100	*
<sup>94</sup> Nb <sup>m</sup>	-86328.2 1.5	40.892 0.012	6.263 m 0.004	3 <sup>+</sup>	06	1962	IT=99.50 6; $\beta^-$ =0.50 6	*
<sup>94</sup> Mo	-88414.08 0.14		STABLE	0 <sup>+</sup>	06	1930	IS=9.187 33	*
<sup>94</sup> Tc	-84158 4		293 m 1	7 <sup>+</sup> *	06	1948	$\beta^+$ =100	*
<sup>94</sup> Tc <sup>m</sup>	-84082 5	76 3	52 m 1	(2 <sup>+</sup> )	06 68Ar06 D	1948	$\beta^+$ $\approx$ 100; IT < 0.18	*
<sup>94</sup> Ru	-82584 3		51.8 m 0.6	0 <sup>+</sup>	06	1952	$\beta^+$ =100	*
<sup>94</sup> Ru <sup>m</sup>	-79940 3	2644.1 0.4	67.5 $\mu$ s 2.8	8 <sup>+</sup>	06 19Ha26 T	1971	IT=100	*
<sup>94</sup> Rh	-72908 3		70.6 s 0.6	(4 <sup>+</sup> )	06 06Ba55 J	1979	$\beta^+$ =100; $\beta^+$ p=1.8 5	*
<sup>94</sup> Rh <sup>m</sup>	-72853 3	54.60 0.20	480 ns 30	(2 <sup>+</sup> )	06	2004	IT=100	*
<sup>94</sup> Rh <sup>n</sup>	-72610# 200#	300# 200#	25.8 s 0.2	(8 <sup>+</sup> )	06	1973	$\beta^+$ =100	*
<sup>94</sup> Pd	-66102 4		9.1 s 0.3	0 <sup>+</sup>	06 19Pa16 TD	1982	$\beta^+$ =100; $\beta^+$ p < 0.13	*
<sup>94</sup> Pd <sup>m</sup>	-61219 4	4883.1 0.4	515 ns 1	(14 <sup>+</sup> )	06 19Ha26 T	1995	IT=100	*
<sup>94</sup> Pd <sup>n</sup>	-58892 4	7209.8 0.8	206 ns 18	(19 <sup>-</sup> )	11Br01 TJ	2011	IT=100	*
<sup>94</sup> Ag	-52400# 400#		27 ms 2	0 <sup>+</sup> #	06 19Pa16 TD	1994	$\beta^+$ =100; $\beta^+$ p < 0.2	*
<sup>94</sup> Ag <sup>m</sup>	-51050# 570#	1350# 400#	470 ms 10	(7 <sup>+</sup> )	06 19Pa16 TD	1994	$\beta^+$ =100; $\beta^+$ p=17.0 6	*
<sup>94</sup> Ag <sup>n</sup>	-45900 370	6500# 550#	400 ms 40	(21 <sup>+</sup> )	06	2002	$\beta^+$ =95.4 7; $\beta^+$ p $\approx$ 27; p=4.1 6; 2p=0.5 3	*
<sup>94</sup> Cd	-40440# 500#		80# ms >760ns	0 <sup>+</sup>	16 16Ce02 I	2016	$\beta^+$ ?; $\beta^+$ p ?	**
* <sup>94</sup> Se	I : 97Be70 > 300ns 95Cz.A > 300ns 17Ch18 observed excited states						**	
* <sup>94</sup> Se <sup>m</sup>	E : uncertainty of 0.3 keV is assumed for all gamma rays in the cascade						**	
* <sup>94</sup> Kr	T : average 16Mi18, 13Mi13=227(14) 03Be05=212(5) 72Am01=200(10)						**	
* <sup>94</sup> Kr	T : 75As04=220(20); other (not used) 96Me09=330(100)						**	
* <sup>94</sup> Rb	J : also 81Th04=3						**	
* <sup>94</sup> Y	J : 07Ch07=2						**	
* <sup>94</sup> Y <sup>m</sup>	T : average 17Ki09=1.33(0.01) 11Ru.A=1.295(0.005) 99Ge02=1.35(0.02);						**	
* <sup>94</sup> Y <sup>m</sup>	T : Birge ratio=2.77						**	
* <sup>94</sup> Ru <sup>m</sup>	T : average 19Ha26=64(4) 71Le19=71(4); other 17Ze02=102(17) for q=44+						**	
* <sup>94</sup> Ru <sup>m</sup>	T : (bare ion); 77Ha49=68(10)						**	
* <sup>94</sup> Pd	T : average 19Pa16=9.1(0.4) 82Ku15=9.0(0.5)						**	
* <sup>94</sup> Pd <sup>m</sup>	T : average 19Ha26=515(1) 17Pa35=495(7) 11Br01=499(13) 09Ga40=468(19)						**	
* <sup>94</sup> Pd <sup>m</sup>	T : 02La18=530(10), same as 98Gr.B=530(10) and supersedes 97Gr02=600(100);						**	
* <sup>94</sup> Pd <sup>m</sup>	T : other 95Go30=800(200)						**	
* <sup>94</sup> Pd <sup>n</sup>	E : from a least-squares fit to Eg						**	
* <sup>94</sup> Pd <sup>n</sup>	T : average 17Pa35=225(32) 11Br01=197(22)						**	

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
* <sup>94</sup> Ag <sup>n</sup>	D : %p=1.9(5) + 2.2(4) from 05Mu15, %2p from 06Mu03						**	
<sup>95</sup> Se	-30460#	500#	70# ms >400ns	3/2 <sup>+</sup> #	12 10Oh02 I	2010	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?	
<sup>95</sup> Br	-43850#	300#	80# ms >300ns	5/2 <sup>-</sup> #	10 97Be70 I	1997	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?	
<sup>95</sup> Br <sup>m</sup>	-43310#	300#	537.9 0.5	6.8 $\mu$ s 1.0	12Ka36 ET	2012	IT=100	
<sup>95</sup> Kr	-56159	19	114 ms 3	1/2 <sup>+</sup> *	10 95Ke04 J	1994	$\beta^-$ =100; $\beta^-$ n=2.87 18; $\beta^-$ 2n ?	
<sup>95</sup> Kr <sup>m</sup>	-55964	19	1.582 $\mu$ s 0.022	(7/2 <sup>+</sup> )	10 12Ka36 T	2006	IT=100	
<sup>95</sup> Rb	-65890	20	377.7 ms 0.8	5/2 <sup>-</sup> *	10	1967	$\beta^-$ =100; $\beta^-$ n=8.7 3	
<sup>95</sup> Rb <sup>m</sup>	-65055	20	< 500 ns	9/2 <sup>+</sup> #	10	2009	IT=100	
<sup>95</sup> Sr	-75117	6	23.90 s 0.14	1/2 <sup>+</sup> *	10	1961	$\beta^-$ =100	
<sup>95</sup> Y	-81208	7	10.3 m 0.1	1/2 <sup>-</sup> *	10	1959	$\beta^-$ =100	
<sup>95</sup> Y <sup>m</sup>	-80120	7	1087.6 0.6	48.6 $\mu$ s 0.5	10 11Ru.A T	1981	IT=100	
<sup>95</sup> Zr	-85659.9	0.9	64.032 d 0.006	5/2 <sup>+</sup>	10	1946	$\beta^-$ =100	
<sup>95</sup> Nb	-86786.3	0.5	34.991 d 0.006	9/2 <sup>+</sup>	10	1951	$\beta^-$ =100	
<sup>95</sup> Nb <sup>m</sup>	-86550.6	0.5	235.69 0.02	3.61 d 0.03	10	1969	IT=94.4 6; $\beta^-$ =5.6 6	
<sup>95</sup> Mo	-87711.87	0.12	STABLE	5/2 <sup>+</sup> *	10	1930	IS=15.873 30	
<sup>95</sup> Tc	-86021	5	19.258 h 0.026	9/2 <sup>+</sup> *	10 20Sz02 T	1947	$\beta^+$ =100	
<sup>95</sup> Tc <sup>m</sup>	-85982	5	38.91 0.04	61.96 d 0.24	10 20Sz02 T	1959	$\beta^+$ =96.1 3; IT=3.9 3	
<sup>95</sup> Ru	-83458	10	1.607 h 0.004	5/2 <sup>+</sup>	10 20Sz02 T	1948	$\beta^+$ =100	
<sup>95</sup> Rh	-78341	4	5.02 m 0.10	(9/2 <sup>+</sup> )	10	1967	$\beta^+$ =100	
<sup>95</sup> Rh <sup>m</sup>	-77798	4	543.3 0.3	1.96 m 0.04	10	1974	IT=88 5; $\beta^+$ =12 5	
<sup>95</sup> Pd	-69966	3	7.4 s 0.4	9/2 <sup>+</sup> #	10 19Pa16 TD	1980	$\beta^+$ =100; $\beta^+$ p=0.23 5	
<sup>95</sup> Pd <sup>m</sup>	-68091	3	1875.13 0.14	13.3 s 0.2	10 19Pa16 TD	1982	$\beta^+$ =89 3; IT=11 3; $\beta^+$ p=0.71 7	
<sup>95</sup> Ag	-59910#	400#	1.78 s 0.06	(9/2 <sup>+</sup> )	10 19Pa16 TD	1994	$\beta^+$ =100; $\beta^+$ p=2.3 2	
<sup>95</sup> Ag <sup>m</sup>	-59570#	400#	< 500 ms	(1/2 <sup>-</sup> )	10	2003	IT=100	
<sup>95</sup> Ag <sup>n</sup>	-57380#	400#	< 16 ms	(23/2 <sup>+</sup> )	10	2003	IT=100	
<sup>95</sup> Ag <sup>p</sup>	-55050#	400#	< 40 ms	(37/2 <sup>+</sup> )	10	2003	IT=100	
<sup>95</sup> Cd	-47060#	570#	32 ms 3	9/2 <sup>+</sup> #	18Pa20 TD	2011	$\beta^+$ =100; $\beta^+$ p=4.6 11	
* <sup>95</sup> Br <sup>m</sup>	T : symmetrized from 12Ka36=6.7(+1.1-0.9)						**	
* <sup>95</sup> Kr <sup>m</sup>	T : others 11Ru.A=1.28(0.05) 06Ge05=1.4(0.2)						**	
* <sup>95</sup> Rb	J : also 81Th04=5/2						**	
* <sup>95</sup> Sr	J : 90Li28=1/2						**	
* <sup>95</sup> Y	J : 07Ch07=1/2						**	
* <sup>95</sup> Ru	T : average 20Sz02=1.6033(0.0044) 70Bo22=1.632(0.021) 68Pi03=1.650(0.017)						**	
* <sup>95</sup> Pd	T : average 19Pa16=7.4(0.5) 12Lo08=7.5(0.5)						**	
* <sup>95</sup> Pd <sup>m</sup>	T : average 19Pa16=13.2(0.4) 82Ku15=13.3(0.3)						**	
* <sup>95</sup> Pd <sup>m</sup>	D : IT from 82Ku15=11(3)%						**	
* <sup>95</sup> Ag	T : average 19Pa16=1.80(0.07) 12Lo08=1.76(0.09)						**	
* <sup>95</sup> Ag	D : % $\beta^+$ p average 19Pa16=2.1(0.3)% 12Lo08=2.5(0.3)%						**	
* <sup>95</sup> Cd	T : others 17Da07=29(8) 10St.A=73(+53-28)						**	
* <sup>95</sup> Cd	D : % $\beta^+$ p symmetrized from 18Pa20=4.5(+1.2-1.0)						**	
<sup>96</sup> Br	-38210#	300#	20# ms >300ns		08 97Be70 I	1997	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?	
<sup>96</sup> Br <sup>m</sup>	-37900#	300#	3.0 $\mu$ s 0.9		12Ka36 ET	2012	IT=100	
<sup>96</sup> Kr	-53082	19	80 ms 8	0 <sup>+</sup>	12	1994	$\beta^-$ =100; $\beta^-$ n=3.7 4	
<sup>96</sup> Rb	-61354	3	201.5 ms 0.9	2 <sup>-</sup> *	08 93Ru01 TD	1967	$\beta^-$ =100; $\beta^-$ n=13.7 5; $\beta^-$ 2n ?	
<sup>96</sup> Rb <sup>m</sup>	-61350#	200#	200# ms >1ms	1(+#)	81Bo30 JI	1981	$\beta^-$ ?; IT ?; $\beta^-$ n ?; $\beta^-$ 2n ?	
<sup>96</sup> Rb <sup>n</sup>	-60219	3	1.80 $\mu$ s 0.04	(10 <sup>-</sup> )	08	1999	IT=100	
<sup>96</sup> Sr	-72918	8	1.059 s .008	0 <sup>+</sup>	08 12Qu01 T	1971	$\beta^-$ =100; $\beta^-$ n ?	
<sup>96</sup> Y	-78330	6	5.34 s 0.05	0 <sup>-</sup> *	08	1975	$\beta^-$ =100	
<sup>96</sup> Y <sup>m</sup>	-76790	6	1540 9 MD	9.6 s 0.2	08	1974	$\beta^-$ =100	
<sup>96</sup> Y <sup>n</sup>	-76675	6	1655.0 1.1	181 ns 9	(6 <sup>+</sup> )	20Is08 EJT	2017	IT=100
<sup>96</sup> Zr	-85438.86	0.11	23.4 Ey 1.7	0 <sup>+</sup>	08 18Ma51 T	1934	IS=2.80 2; $2\beta^-$ =100	
<sup>96</sup> Nb	-85602.83	0.15	23.35 h 0.05	6 <sup>+</sup>	08	1949	$\beta^-$ =100	
<sup>96</sup> Mo	-88794.89	0.12	STABLE	0 <sup>+</sup>	08	1930	IS=16.673 8	
<sup>96</sup> Tc	-85822	5	4.28 d 0.07	7 <sup>+</sup> *	08	1947	$\beta^+$ =100	
<sup>96</sup> Tc <sup>m</sup>	-85788	5	34.23 0.04	51.5 m 1.0	08	1950	IT=98.0 5; $\beta^+$ =2.0 5	
<sup>96</sup> Ru	-86080.39	0.17	STABLE	>80Ey	0 <sup>+</sup>	08 13Be09 T	1931	IS=5.54 14; $2\beta^+$ ?
<sup>96</sup> Rh	-79688	10	9.90 m 0.10	6 <sup>+</sup>	08	1967	$\beta^+$ =100	

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{96}\text{Rh}^m$	-79636	10	51.98 0.09	1.51 m 0.02	3 <sup>+</sup>	08	1966	IT=60 5; $\beta^+$ =40 5
$^{96}\text{Pd}$	-76183	4		122 s 2	0 <sup>+</sup>	08	1980	$\beta^+$ =100
$^{96}\text{Pd}^m$	-73652	4	2530.57 0.23	1.804 $\mu\text{s}$ 0.007	8 <sup>+</sup> #	08 17Pa35	T 1983	IT=100
$^{96}\text{Ag}$	-64510	90		4.45 s 0.03	(8) <sup>+</sup>	08 19Pa16	TD 1982	$\beta^+$ =100; $\beta^+$ p=4.2 4
$^{96}\text{Ag}^m$	-64510#	100#	0# 50#	6.9 s 0.5	(2 <sup>+</sup> )	08 12Lo08	TD 2003	$\beta^+$ =100; $\beta^+$ p=14.9 18
$^{96}\text{Ag}^n$	-62050	90	2461.4 0.3	103.2 $\mu\text{s}$ 4.5	(13 <sup>-</sup> )	11Bo23	TJD 2011	IT=100
$^{96}\text{Ag}^p$	-61820	90	2686.7 0.4	1.561 $\mu\text{s}$ 0.016	(15 <sup>+</sup> )	08 17Pa35	ETJ 2011	IT=100
$^{96}\text{Ag}^q$	-57560	90	6951.8 1.4	132 ns 17	(19 <sup>+</sup> )	17Pa35	ETJ 2011	IT=100
$^{96}\text{Cd}$	-55570#	410#		1003 ms 47	0 <sup>+</sup>	10 19Pa16	TD 2008	$\beta^+$ =100; $\beta^+$ p=1.6 3
$^{96}\text{Cd}^m$	-49540#	1450#	6030 1390	511 ms 26	16 <sup>+</sup>	10 19Pa16	ETD2011	$\beta^+$ =100; $\beta^+$ p=15.4 21
$^{96}\text{Cd}^n$	-49970#	410#	5605 5	198 ns 18	(12 <sup>-</sup> , 13 <sup>-</sup> )	19Da02	EJ 2019	IT=100
$^{96}\text{In}$	-38090#	500#		1# ms >400ns		16 16Ce02	I 2016	$\beta^+$ ?; p ?
$^{96}\text{Br}^m$	T : symmetrized from 12Ka36=2.7(+1.1-0.7)							**
$^{96}\text{Rb}$	J : 81Th04=2							**
$^{96}\text{Rb}$	D : % $\beta^-$ -n average 93Ru01=14.7(1.0) 81Ho07=14.7(1.2) 81En05=14.2(1.2)							**
$^{96}\text{Rb}$	D : 79Ri09=12.5(0.9) 69Am01=12.7(1.5)							**
$^{96}\text{Rb}$	T : average 12Qu01=212(17) 03Be05=197(6) 93Ru01=201(1) 79Ri09=197(5)							**
$^{96}\text{Rb}$	T : 78Wo09=203(4) 77Re05=205(4) 74Ro15=199(3.5) 71Tr02=207(3)							**
$^{96}\text{Rb}^m$	I : non-observation in 81Th04 is not in contradiction with 81Bo30							**
$^{96}\text{Rb}^n$	T : average 12Ka36=1.72(+0.16-0.14) 11Ru.A=1.77(0.05) 05Pi13=2.0(0.1)							**
$^{96}\text{Rb}^n$	T : 99Ge01=1.65(0.15)							**
$^{96}\text{Sr}$	T : average 12Qu01=0.950(0.035) 90Ma03=1.07(0.01) 79En02=1.10(0.02)							**
$^{96}\text{Sr}$	T : 78Wo09=1.015(0.019) 75Ba36=1.06(0.04)							**
$^{96}\text{Y}$	J : 07Ch07=0							**
$^{96}\text{Y}^m$	J : 07Ch07=8							**
$^{96}\text{Zr}$	T : 2v- $\beta\beta$ average 18Ma51=20.3(+4.6-0.31) 10Ar07=23.5(1.4.stat)(1.6.syst)							**
$^{96}\text{Zr}$	T : 99Ar25=21(+8-4.stat + 2,syst); others 93Ka12=39(9) and 01Wi17=9.4(3.2)							**
$^{96}\text{Zr}$	T : in geochemical exp., 16Fi01>24Ey; 20Ba.A,15Ba11=23(2) (evaluation)							**
$^{96}\text{Ru}$	T : 2nu- $\beta^+\epsilon$ >80 Ey (theor. most probable); 2nu $\beta^+\beta^+$ >140 Ey Onu2K>1 Zy							**
$^{96}\text{Pd}^m$	T : average 17Pa35=1.80(0.01) 98Gr.B=1.81(0.01), supersedes 97Gr02=1.7(0.1),							**
$^{96}\text{Pd}^m$	T : 09Ga40=1.76(0.05); others 07My02=2.10(0.21) 83Gr01=2.2(0.3)							**
$^{96}\text{Ag}$	T : average 19Pa16=4.46(0.04) 03Ba39=4.40(0.06) 97Sc30=4.50(0.06)							**
$^{96}\text{Ag}$	D : % $\beta^+$ p average 19Pa16=4.4(0.5) 96He25=3.7(0.9)							**
$^{96}\text{Ag}^m$	T : average 12Lo08=6.8(1.0) 03Ba39=6.9(0.6)							**
$^{96}\text{Ag}^m$	D : % $\beta^+$ p average 19Pa16=14.7(2.4) 12Lo08=14(3) 03Ba39=18(5)							**
$^{96}\text{Ag}^n$	E : from a least-squares fit to Eg using 11Bo23 level scheme							**
$^{96}\text{Ag}^n$	T : average 19Ha26=104(5) 11Bo23=100(10)							**
$^{96}\text{Ag}^p$	E : 43.7(0.2) keV above the 2643(0.3) keV, 13+ level							**
$^{96}\text{Ag}^p$	T : average 17Pa35=1.57(0.02) 11Bo23=1.56(0.03) 11Be34=1.45(0.07)							**
$^{96}\text{Ag}^q$	E : 4265(2) keV above 96Agp							**
$^{96}\text{Ag}^q$	T : average 19Ha26=120(20), supersedes 17Pa35=160(41), 11Bo23=160(30)							**
$^{96}\text{Cd}$	T : average 19Pa16=1020(60) 17Da07=970(90) 10St.A=990(130)							**
$^{96}\text{Cd}$	D : % $\beta^+$ p average 19Pa16=1.7(0.4) 17Da07=1.5(0.5)							**
$^{96}\text{Cd}^m$	E : symmetrized from 5810(+1560-1220) keV							**
$^{96}\text{Cd}^m$	T : average 19Pa16=530(30) 17Da07=450(+50-40); other 11Na34=290(+110-100)							**
$^{96}\text{Cd}^m$	D : % $\beta^+$ p average 19Pa16=19.5(2.9) 17Da07=11(3)							**
$^{96}\text{Cd}^n$	T : symmetrized from 19Da02=197(+19-17)							**
$^{96}\text{Cd}^n$	E : uncertainty of 1 keV is assumed for all gamma rays in the cascade							**
$^{97}\text{Br}$	-34000#	400#		40# ms >300ns	5/2 <sup>-</sup> #	10	1997	$\beta^-$ ?; $\beta^-$ -n ?; $\beta^-$ -2n ?
$^{97}\text{Kr}$	-47420	130		62.2 ms 3.2	3/2 <sup>+</sup> #	10 11Ni01	T 1997	$\beta^-$ =100; $\beta^-$ -n=6.7 6; $\beta^-$ -2n ?
$^{97}\text{Rb}$	-58519.1	1.9		169.1 ms 0.6	3/2 <sup>+</sup> *	15	1969	$\beta^-$ =100; $\beta^-$ -n=25.5 9; $\beta^-$ -2n ?
$^{97}\text{Rb}^m$	-58442.5	1.9	76.6 0.2	5.7 $\mu\text{s}$ 0.6	(1/2,3/2) <sup>-</sup>	15	2012	IT=100
$^{97}\text{Sr}$	-68581	3		432 ms 4	1/2 <sup>+</sup> *	10 02Pf04	D 1978	$\beta^-$ =100; $\beta^-$ -n=0.02 1
$^{97}\text{Sr}^m$	-68273	3	308.13 0.11	175.2 ns 2.1	7/2 <sup>+</sup>	10 19Es04	T 1990	IT=100
$^{97}\text{Sr}^n$	-67750	3	830.83 0.23	513 ns 5	(9/2 <sup>+</sup> )	10 19Es04	T 1974	IT=100
$^{97}\text{Y}$	-76115	7		3.75 s 0.03	1/2 <sup>-</sup> *	10	1970	$\beta^-$ =100; $\beta^-$ -n=0.055 4
$^{97}\text{Y}^m$	-75447	7	667.52 0.23	1.17 s 0.03	9/2 <sup>+</sup> *	10 83Re10	D 1970	$\beta^-$ >99.3;IT<0.7; $\beta^-$ -n=0.11 3
$^{97}\text{Y}^n$	-72592	7	3522.6 0.4	142 ms 8	(27/2 <sup>-</sup> )*	10	1986	IT=94.8 9; $\beta^-$ =5.2 9
$^{97}\text{Zr}$	-82936.69	0.12		16.749 h 0.008	1/2 <sup>+</sup> *	10	1951	$\beta^-$ =100
$^{97}\text{Zr}^m$	-81672.34	0.20	1264.35 0.16	104.8 ns 1.7	7/2 <sup>+</sup>	10 11Ru.A	T 1976	IT=100



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{97}\text{Nb}$	-85603	4			72.1 m 0.7	$9/2^+$	10	1951	$\beta^- = 100$
$^{97}\text{Nb}^m$	-84860	4	743.35	0.03	58.7 s 1.8	$1/2^-$	10	1950	IT=100
$^{97}\text{Mo}$	-87544.70	0.16			STABLE	$5/2^{+*}$	10	1930	IS=9.582 15
$^{97}\text{Tc}$	-87224	4			4.21 My0.16	$9/2^{+*}$	10 20Kr09 J	1946	$\epsilon = 100$
$^{97}\text{Tc}^m$	-87127	4	96.57	0.06	91.1 d 0.6	$1/2^-$	10 98Ko27 DT	1954	IT=96.06 18; $\epsilon = 3.94$ 18
$^{97}\text{Ru}$	-86120.6	2.8			2.8370 d 0.0014	$5/2^+$	10 09Go29 T	1946	$\beta^+ = 100$
$^{97}\text{Rh}$	-82600	40			30.7 m 0.6	$9/2^+$	10	1955	$\beta^+ = 100$
$^{97}\text{Rh}^m$	-82340	40	258.76	0.18	46.2 m 1.6	$1/2^-$	10	1971	$\beta^+ = 94.4$ 6;IT=5.6 6
$^{97}\text{Pd}$	-77806	5			3.10 m 0.09	$5/2^{+ \#}$	10	1969	$\beta^+ = 100$
$^{97}\text{Ag}$	-70904	12			25.5 s 0.3	$(9/2)^{+*}$	10	1978	$\beta^+ = 100$
$^{97}\text{Ag}^m$	-70290	40	620	40 MD	100# ms	$1/2^- \#$	20Ho03 E	2019	IT ?
$^{97}\text{Cd}$	-60730	420			1.16 s 0.05	$(9/2^+)$	10 19Pa16 TD	1978	$\beta^+ = 100; \beta^+ p = 7.4$ 2
$^{97}\text{Cd}^m$	-59480	420	1245.1	0.2	730 $\mu\text{s}$ 70	$(1/2^-)$	19Pa16 ETD2019		IT=100
$^{97}\text{Cd}^n$	-58110	720	2620	580	3.86 s 0.06	$(25/2^+)$	10 19Pa16 ETD1982		$\beta^+ = 100; \beta^+ p = 25.1$ 5
$^{97}\text{In}$	-47390#	400#			36 ms 6	$9/2^{+ \#}$	18Pa20 TD	2011	$\beta^+ = 100; \beta^+ p = 2.3$ 13; p ?
$^{97}\text{In}^m$	-46990#	410#	400#	100#	0.12 ms 0.07	$1/2^- \#$	20 18Pa20 TD	2018	p ?
$^{*97}\text{Kr}$	T : average 11Ni01=60(+6-5) 03Be05=63(4)								
$^{*97}\text{Sr}$	J : 90Li28=1/2								
$^{*97}\text{Sr}$	T : average 12Qu01=456(5,stat)(13,syst) 86Wa17=429(5) 87PFZX=420(20)								
$^{*97}\text{Sr}$	T : 82Ga24=420(40) 78Wo09=441(15); others (not used) 81En05=390(30)								
$^{*97}\text{Sr}$	T : 83Re10=403(4), superseded by 86Wa17 79En02=430(30), superseded by								
$^{*97}\text{Sr}$	T : 81En05								
$^{*97}\text{Sr}^m$	T : average 19Es04=174.7(6.9) 15Cz01=165(4) 11Ru.A=180.9(2.8) 06Hw01=165(25)								
$^{*97}\text{Sr}^m$	T : 83Kr11=170(10)								
$^{*97}\text{Sr}^n$	T : average 19Es04=526(17) 18Rz01=504(8) 13Ru07=515(10) 05Zl01=526(13);								
$^{*97}\text{Sr}^n$	T : others (not used) 12Ka36=520(+160-120) 80Mo.A=515(15) 06Hw01=255(56)								
$^{*97}\text{Sr}^n$	T : 03Hw03=265(27), non standard technique and conflicting								
$^{*97}\text{Y}$	J : 07Ch07=1/2								
$^{*97}\text{Y}^m$	J : 07Ch07=9/2								
$^{*97}\text{Y}^m$	D : % $\beta^-$ -n from 83Re10=0.11(0.03); other 86Wa17<0.08								
$^{*97}\text{Y}^n$	J : 07Ch07=(27/2)								
$^{*97}\text{Zr}$	J : 02Ca37=1/2								
$^{*97}\text{Zr}^m$	T : average 11Ru.A=106.1(2.1) 85Be20=102(3); others outweighed 06Hw01=97(16)								
$^{*97}\text{Zr}^m$	T : 96Lh03=106(7)								
$^{*97}\text{Tc}$	T : from 98Ko27								
$^{*97}\text{Tc}^m$	T : average 98Ko27=91.4(0.8), supersedes 93Ko64=92.2(1.8), 54Bo24=90.5(1.0)								
$^{*97}\text{Ag}$	J : 14Fe01=(9/2)								
$^{*97}\text{Cd}$	T : average 19Pa16=1.20(0.07) 11Lo09=1.10(0.08)								
$^{*97}\text{Cd}$	D : other % $\beta^+$ p 11Lo09=11.8(20)								
$^{*97}\text{Cd}^n$	J : 11Lo09=(25/2+)								
$^{*97}\text{In}$	T : other 10St.A=26(+47-10)								
$^{*97}\text{In}$	D : % $\beta^+$ p symmetrized from 18Pa20=1.7(+1.7-0.8)								
$^{*97}\text{In}^m$	T : from 1.3<T<230 us in 18Pa20								
$^{98}\text{Br}$	-28050#	400#			15# ms >400ns		20 10Oh02 I	2010	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$
$^{98}\text{Kr}$	-44120#	300#			42.8 ms 3.6	$0^+$	20 11Ni01 T	1997	$\beta^- = 100; \beta^- n = 7.0$ 10; $\beta^- 2n ?$
$^{98}\text{Rb}$	-54369	16			115 ms 6	$(0^-)^*$	20	1971	$\beta^- = 100; \beta^- n = 14.3$ 9; $\beta^- 2n = 0.054$ 8
$^{98}\text{Rb}^m$	-54296	20	73	26 BD	96 ms 3	$(3^+)^*$	20	1980	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$
$^{98}\text{Rb}^n$	-54191	16	178.5	0.4	358 ns 7	$(2^-)$	20 FGK205 J	2009	IT=100
$^{98}\text{Sr}$	-66422	3			653 ms 2	$0^+$	20	1971	$\beta^- = 100; \beta^- n = 0.23$ 3
$^{98}\text{Y}$	-72289	8			548 ms 2	$0^-*$	20	1970	$\beta^- = 100; \beta^- n = 0.33$ 3
$^{98}\text{Y}^m$	-72118	8	170.78	0.05	615 ns 8	$2^-$	20	1972	IT=100
$^{98}\text{Y}^n$	-71823	8	465.7	0.7 MD	2.32 s 0.08	$(6, 7)^+$	20 17Ur03 JED	1977	$\beta^- \approx 100; \text{IT} ?; \beta^- n = 3.44$ 95
$^{98}\text{Y}^p$	-71793	8	496.10	0.11	6.90 $\mu\text{s}$ 0.054	$(4)^-$	20	1970	IT=100
$^{98}\text{Y}^q$	-71695	13	594	10	180 ns 7	$(3^-, 4^-)$	20	2017	IT=100
$^{98}\text{Y}^r$	-71317	8	972.17	0.20	450 ns 150	$(8^+)$	20	2017	IT=100
$^{98}\text{Y}^s$	-71108	8	1181.50	0.18	762 ns 14	$(10^-)$	20	1972	IT=100
$^{98}\text{Zr}$	-81282	8			30.7 s 0.4	$0^+$	20	1967	$\beta^- = 100$
$^{98}\text{Zr}^m$	-74680	8	6601.9	1.1	1.9 $\mu\text{s}$ 0.2	$(17^-)$	20	2005	IT=100
$^{98}\text{Nb}$	-83525	5			2.86 s 0.06	$1^+$	20	1960	$\beta^- = 100$
$^{98}\text{Nb}^m$	-83441	6	84	4	51.1 m 0.4	$(5)^+$	20	1948	$\beta^- \approx 100; \text{IT} ?$

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>98</sup> Mo	-88115.98	0.17	STABLE >100Ty	0 <sup>+</sup>	20 52Fr23 T	1930	IS=24.292 80;2β <sup>-</sup> ? *	
<sup>98</sup> Tc	-86432	3	4.2 My0.3	6 <sup>+</sup> *	20 20Kr09 J	1955	β <sup>-</sup> =100;β <sup>+</sup> =0	
<sup>98</sup> Tc <sup>m</sup>	-86341	3	90.77 0.16	14.7 μs 0.5	(2,3) <sup>-</sup>	20	IT=100	
<sup>98</sup> Ru	-88225	6	STABLE	0 <sup>+</sup>	20	1944	IS=1.87 3	
<sup>98</sup> Rh	-83175	12	*	8.72 m 0.12	(2) <sup>+</sup>	20	1955 β <sup>+</sup> =100	
<sup>98</sup> Rh <sup>m</sup>	-83119	12	56.3 1.0 *	3.6 m 0.2	(5 <sup>+</sup> )	20	1966 IT=89 5;β <sup>+</sup> =11 5	
<sup>98</sup> Pd	-81321	5		17.7 m 0.4	0 <sup>+</sup>	20	1955 β <sup>+</sup> =100	
<sup>98</sup> Ag	-73070	30		47.5 s 0.3	(6) <sup>+</sup> *	20	1978 β <sup>+</sup> =100;β <sup>+</sup> p=0.0012 5 *	
<sup>98</sup> Ag <sup>m</sup>	-72960	30	107.28 0.10	161 ns 7	(4 <sup>+</sup> )	20	1998 IT=100	
<sup>98</sup> Cd	-67640	50		9.29 s 0.10	0 <sup>+</sup>	20	1978 β <sup>+</sup> =100;β <sup>+</sup> p<0.029 *	
<sup>98</sup> Cd <sup>m</sup>	-65210	50	2428.3 0.4	154 ns 16	(8 <sup>+</sup> )	20 17Pa35 T	1996 IT=100	
<sup>98</sup> Cd <sup>n</sup>	-61010	50	6635 2	224 ns 5	(12 <sup>+</sup> )	20	2004 IT=100	
<sup>98</sup> In	-53910#	300#	*	30 ms 1	(0 <sup>+</sup> )	20	1994 β <sup>+</sup> =100;β <sup>+</sup> p≤0.13	
<sup>98</sup> In <sup>m</sup>	-53090#	790#	820 730 *	890 ms 20	(9 <sup>+</sup> )	20	2001 β <sup>+</sup> =100;β <sup>+</sup> p=44 2	
* <sup>98</sup> Kr	T : average 11Ni01=42(4) 03Be05=46(8)							**
* <sup>98</sup> Rb <sup>n</sup>	J : 178.5 keV gamma to (0-) and Weisskopf estimates for E1,M1 and E2							**
* <sup>98</sup> Y	J : 07Ch07=0							**
* <sup>98</sup> Y <sup>m</sup>	T : average 17Ur03=640(20) 11Ru.A=610(9); other: 70Gr38=620(80)							**
* <sup>98</sup> Y <sup>n</sup>	J : other: 07Ch07=(4,5) hfs							**
* <sup>98</sup> Y <sup>n</sup>	D : %β <sup>-</sup> n from 81En05							**
* <sup>98</sup> Y <sup>p</sup>	T : average 17Ur03=6.95(0.06) 11Ru.At=6.87(0.05)							**
* <sup>98</sup> Y <sup>q</sup>	E : 564.0+x keV is proposed in 17Ur03; x=30(10) is estimated by Nubase							**
* <sup>98</sup> Y <sup>r</sup>	T : average 11Ru.At=806(21) 17Ur03=720(20) 70Gr38=830(100)							**
* <sup>98</sup> Mo	T : Onu-BB 52Fr23>100 Ty (theoretically faster, see text)							**
* <sup>98</sup> Ag	D : %β <sup>+</sup> p symmetrized from 96He25=0.0011(+5-4)							**
* <sup>98</sup> Ag	J : 14Fe01=(5,6)							**
* <sup>98</sup> Cd	T : average 92Pl01=9.2(0.3) 19Pa16=9.3 (0.1)							**
<sup>99</sup> Kr	-38400#	400#		40 ms 11	5/2 <sup>-</sup> #	17 03Be05 TD	1997 β <sup>-</sup> =100;β <sup>-</sup> n=11 7;β <sup>-</sup> 2n ? *	
<sup>99</sup> Rb	-51121	4		54 ms 4	(3/2 <sup>+</sup> )	17 02Pf04 D	1971 β <sup>-</sup> =100;β <sup>-</sup> n=17.3 25; β <sup>-</sup> 2n ?	
<sup>99</sup> Sr	-62519	5		269.2 ms 1.0	3/2 <sup>+</sup> *	17 93Ru01 D	1975 β <sup>-</sup> =100;β <sup>-</sup> n=0.100 19 *	
<sup>99</sup> Y	-70644	7		1.484 s 0.007	5/2 <sup>+</sup> *	17	1975 β <sup>-</sup> =100;β <sup>-</sup> n=1.77 19	
<sup>99</sup> Y <sup>m</sup>	-68502	7	2141.65 0.19	8.2 μs 0.4	(17/2 <sup>+</sup> )	17	1985 IT=100 *	
<sup>99</sup> Zr	-77617	10		2.1 s 0.1	1/2 <sup>+</sup> *	17	1970 β <sup>-</sup> =100 *	
<sup>99</sup> Zr <sup>m</sup>	-77365	10	251.96 0.09	336 ns 5	7/2 <sup>+</sup>	17 20Bo04 T	1970 IT=100 *	
<sup>99</sup> Nb	-82335	12		15.0 s 0.2	9/2 <sup>+</sup> *	17	1950 β <sup>-</sup> =100 *	
<sup>99</sup> Nb <sup>m</sup>	-81970	12	365.27 0.08	2.5 m 0.2	1/2 <sup>-</sup>	17	1960 β <sup>-</sup> ≈100;IT=?	
<sup>99</sup> Mo	-85970.11	0.23		65.932 h 0.005	1/2 <sup>+</sup> *	17 FGK209 T	1948 β <sup>-</sup> =100 *	
<sup>99</sup> Mo <sup>m</sup>	-85872.33	0.23	97.785 0.003	15.5 μs 0.2	5/2 <sup>+</sup>	17	1958 IT=100	
<sup>99</sup> Mo <sup>n</sup>	-85286.01	0.30	684.10 0.19	760 ns 60	11/2 <sup>-</sup>	17	1975 IT=100	
<sup>99</sup> Tc	-87327.9	0.9		211.1 ky 1.2	9/2 <sup>+</sup> *	17 20Kr09 J	1938 β <sup>-</sup> =100 *	
<sup>99</sup> Tc <sup>m</sup>	-87185.2	0.9	142.6836 0.0011	6.0066 h 0.0002	1/2 <sup>-</sup> *	17 FGK209 T	1958 IT≈100;β <sup>-</sup> =0.0037 6	
<sup>99</sup> Ru	-87625.4	0.3		STABLE	5/2 <sup>+</sup> *	17	1931 IS=12.76 14 *	
<sup>99</sup> Rh	-85585	19		16.1 d 0.2	1/2 <sup>-</sup>	17	1952 β <sup>+</sup> =100	
<sup>99</sup> Rh <sup>m</sup>	-85521	19	64.4 0.5	4.7 h 0.1	9/2 <sup>+</sup> *	17	1952 β <sup>+</sup> ≈100;IT ?	
<sup>99</sup> Pd	-82183	5		21.4 m 0.2	(5/2) <sup>+</sup>	17	1955 β <sup>+</sup> =100	
<sup>99</sup> Ag	-76712	6		2.07 m 0.05	(9/2) <sup>+</sup> *	17	1967 β <sup>+</sup> =100 *	
<sup>99</sup> Ag <sup>m</sup>	-76206	6	506.2 0.4	10.5 s 0.5	(1/2) <sup>-</sup> *	17	1978 IT=100 *	
<sup>99</sup> Cd	-69931.1	1.6		17 s 1	5/2 <sup>+</sup> #	17 19Pa16 TD	1978 β <sup>+</sup> =100;β <sup>+</sup> p=0.21 2; β <sup>+</sup> α<1e-4	
<sup>99</sup> In	-61380#	300#		3.11 s 0.06	9/2 <sup>+</sup> #	17 19Pa16 TD	1994 β <sup>+</sup> =100;β <sup>+</sup> p=0.29 3	
<sup>99</sup> In <sup>m</sup>	-60980#	340#	400# 150#	1# s	1/2 <sup>-</sup> #		β <sup>+</sup> ?;β <sup>+</sup> p ?;IT ?	
<sup>99</sup> Sn	-47980#	580#		24 ms 4	9/2 <sup>+</sup> #	17 18Pa20 TD	2011 β <sup>+</sup> =100;β <sup>+</sup> p=5 3 *	
* <sup>99</sup> Kr	T : other 11Ni01=13(+34-6)							**
* <sup>99</sup> Sr	J : 91Li05,90Li28=3/2							**
* <sup>99</sup> Sr	T : average 86ReZU=269(1) 83Re10=274(4) 83Wo10=266(6)							**
* <sup>99</sup> Y	J : 07Ch07=5/2							**
* <sup>99</sup> Y	T : other (recent) 19Do02=1.27(0.25)							**
* <sup>99</sup> Y <sup>m</sup>	T : average 13RuZX=8.0(0.5) 85Me09=8.6(0.8); other: 99Ge01=11(2)							**
* <sup>99</sup> Zr	J : 02Ca37=1/2							**
* <sup>99</sup> Zr <sup>m</sup>	J : 130.2-keV gamma ray, E2 to 3/2+							**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)		
* <sup>99</sup> Zr <sup>m</sup>	T : other 04Hw02=316(48) 99Ge01=400(20) 79Se01=292(20) 70Gr38=400(80)							**	
* <sup>99</sup> Nb	J : 09Ch25=9/2							**	
* <sup>99</sup> Mo	J : 74Ru05=1/2							**	
* <sup>99</sup> Tc	J : 20Kr09,58Lo62=9/2							**	
* <sup>99</sup> Ru	J : 13Ma15=5/2							**	
* <sup>99</sup> Ag	J : 14Fe01=(9/2)							**	
* <sup>99</sup> Ag <sup>m</sup>	J : 14Fe01=(1/2)							**	
* <sup>99</sup> Sn	D : %β <sup>+</sup> p symmetrized from 18Pa20=3.9(+3.4-1.7)							**	
<sup>100</sup> Kr	-34470#	400#	12 ms 8	0 <sup>+</sup>	11 11Ni01	TD 1997	β <sup>-</sup> =100;β <sup>-</sup> n ?;β <sup>-</sup> 2n ?	*	
<sup>100</sup> Rb	-46266	13	51.3 ms 1.6	4 <sup>-</sup> #	08 20Mu.A	T 1978	β <sup>-</sup> =100;β <sup>-</sup> n=5.6 12; β <sup>-</sup> 2n=0.15 5	*	
<sup>100</sup> Sr	-59818	7	202.1 ms 1.7	0 <sup>+</sup>	08 02Pf04	D 1978	β <sup>-</sup> =100;β <sup>-</sup> n=1.11 34	*	
<sup>100</sup> Sr <sup>m</sup>	-58199	7	1618.72 0.20	122 ns 9	(4 <sup>-</sup> )	12Ka36 T 1995	IT=100	*	
<sup>100</sup> Y	-67321	11	&	940 ms 30	4 <sup>+</sup> *	08 10Ba31 J 1977	β <sup>-</sup> =100;β <sup>-</sup> n ?	*	
<sup>100</sup> Y <sup>m</sup>	-67177	11	144 16 MD &	727 ms 6	1 <sup>+</sup> #	08 93Ru01 D 1977	β <sup>-</sup> =100;β <sup>-</sup> n=1.08 6	*	
<sup>100</sup> Zr	-76373	8	7.1 s 0.4	0 <sup>+</sup>	08	1970	β <sup>-</sup> =100		
<sup>100</sup> Nb	-79791	8	1.5 s 0.2	1 <sup>+</sup>	08	1967	β <sup>-</sup> =100		
<sup>100</sup> Nb <sup>m</sup>	-79478.5	2.0	313 8 MD	2.99 s 0.11	(5 <sup>+</sup> )	08	1967	β <sup>-</sup> =100	
<sup>100</sup> Nb <sup>n</sup>	-79444	11	347 8	460 ns 60	(4 <sup>-</sup> , 5 <sup>-</sup> )	08	1986	IT=100	*
<sup>100</sup> Nb <sup>p</sup>	-79057	11	734 8	12.43 μs 0.26	(8 <sup>-</sup> )	08 11Ru.A T 1980	IT=100	*	
<sup>100</sup> Mo	-86193.0	0.3	7.07 Ey 0.14	0 <sup>+</sup>	08 20Ba.A T 1930	1930	IS=9.744 65;2β <sup>-</sup> =100	*	
<sup>100</sup> Tc	-86021.0	1.4	15.46 s 0.19	1 <sup>+</sup>	08	1952	β <sup>-</sup> ≈100;ε=0.0018 9		
<sup>100</sup> Tc <sup>m</sup>	-85820.3	1.4	200.67 0.04	8.32 μs 0.14	(4 <sup>+</sup> )	08	1958	IT=100	
<sup>100</sup> Tc <sup>n</sup>	-85777.1	1.4	243.95 0.04	3.2 μs 0.2	(6 <sup>+</sup> )	08	1967	IT=100	
<sup>100</sup> Ru	-89227.4	0.3	STABLE	0 <sup>+</sup>	08	1931	IS=12.60 7		
<sup>100</sup> Rh	-85591	18	20.8 h 0.1	1 <sup>-</sup> *	08	1948	β <sup>+</sup> =100;ε=95.1 5; e <sup>+</sup> =4.9 5		
<sup>100</sup> Rh <sup>m</sup>	-85516	18	74.782 0.014	214.0 ns 2.0	(2 <sup>+</sup> )	08	1965	IT=100	
<sup>100</sup> Rh <sup>n</sup>	-85483	18	107.6 0.2	4.6 m 0.2	(5 <sup>+</sup> )	08	1973	IT≈98.3;β <sup>+</sup> ≈1.7	
<sup>100</sup> Rh <sup>p</sup>	-85371	18	219.61 0.22	130 ns 10	(7 <sup>+</sup> )	08	1984	IT=100	
<sup>100</sup> Pd	-85213	18	3.63 d 0.09	2.01 m 0.09	0 <sup>+</sup>	08	1948	ε=100	
<sup>100</sup> Ag	-78138	5	2.01 m 0.09	2.01 m 0.09	(5 <sup>+</sup> ) <sup>+</sup> *	08	1970	β <sup>+</sup> =100	*
<sup>100</sup> Ag <sup>m</sup>	-78122	5	15.52 0.16	2.24 m 0.13	(2 <sup>+</sup> )	08	1980	β <sup>+</sup> =?;IT ?	
<sup>100</sup> Cd	-74194.6	1.7	49.1 s 0.5	0 <sup>+</sup>	10	1970	β <sup>+</sup> =100		
<sup>100</sup> In	-64178.1	2.2	5.62 s 0.06	6 <sup>+</sup> #	14 19Pa16	TD 1982	β <sup>+</sup> =100;β <sup>+</sup> p=1.66 3		
<sup>100</sup> Sn	-57150	240	1.18 s 0.08	0 <sup>+</sup>	14 19Lu08	T 1994	β <sup>+</sup> =100;β <sup>+</sup> p<17	*	
<sup>100</sup> Sn <sup>m</sup>	-52650#	310#	4500# 200#	100# ns	6 <sup>+</sup> #	11Hi.A ETJ	p ?		
* <sup>100</sup> Kr	T : symmetrized from 11Ni01=7(+11-3)							**	
* <sup>100</sup> Rb	T : average 20Mu.A=50(5) 11Ni01=48(3) 87PfZX=53(2)							**	
* <sup>100</sup> Rb	D : %β <sup>-</sup> n from 93Ru01; %β <sup>-</sup> 2n from P2n/Pn=0.027 7 in 81JoZV and %β <sup>-</sup> n							**	
* <sup>100</sup> Sr	T : average 11Ni01=181(+16-13) 93Ru01=165(24) 87PfZX=207(10) 86Wa17=204(2)							**	
* <sup>100</sup> Sr	T : 86Wo01=193(4) 83Mu19=214(8) 78Ko29=170(80) 85IaZZ=200(20)							**	
* <sup>100</sup> Sr <sup>m</sup>	T : other 95Pf04=85(7)							**	
* <sup>100</sup> Y	J : 10Ba31=4+ and p5/2[422] n3/2[411], K=4+ configuration by the measured							**	
* <sup>100</sup> Y	J : magnetic moment; other 07Ch07=(3), Ensdf2008 assigns J=(3,4,5) and							**	
* <sup>100</sup> Y	J : associate this state with an excited isomer							**	
* <sup>100</sup> Y	T : from β - γ(t) in 77Kh03, where low- and high-spin β <sup>-</sup> decaying							**	
* <sup>100</sup> Y	T : isomers were resolved; recent (from β(t)) 09Pe06=660(+150-120)							**	
* <sup>100</sup> Y	T : and 12Qu01=845(80,stat)(55,syst) include both gs and isomer							**	
* <sup>100</sup> Y <sup>m</sup>	T : average 96Me09=710(30) 86Wo01=735(7) 83Mu14=682(18) 73Kh03=550(150)							**	
* <sup>100</sup> Y <sup>m</sup>	T : from data dominated by the <sup>100</sup> Sr (J <sup>π</sup> =0+) isobar and hence							**	
* <sup>100</sup> Y <sup>m</sup>	T : a preferable feeding to the low-spin isomer							**	
* <sup>100</sup> Y <sup>m</sup>	J : direct β <sup>-</sup> feeding to 0+ states in <sup>100</sup> Zr in 86Wo01;							**	
* <sup>100</sup> Y <sup>m</sup>	J : p5/2[422] n3/2[411], K=1+ configuration							**	
* <sup>100</sup> Y <sup>m</sup>	D : %β <sup>-</sup> n other (after 93Ru01) 96Me09=1.8(0.6) 02Pf04=1.16(0.32), compilation							**	
* <sup>100</sup> Nb <sup>n</sup>	E : 34.3 keV above the 5+ isomer							**	
* <sup>100</sup> Nb <sup>n</sup>	J : a cascade of two M1 gamma rays from 6- and absence of gamma from 8-							**	
* <sup>100</sup> Nb <sup>p</sup>	E : 420.7 keV above the 5+ isomer							**	
* <sup>100</sup> Nb <sup>p</sup>	J : 28 keV gamma, (E2) to (6-)							**	
* <sup>100</sup> Mo	T : 2v-ββ symmetrized from 20Ba.A=7.06(+0.15-0.13) (evaluation); others							**	
* <sup>100</sup> Mo	T : (recent) 20Ar09=7.12(+0.18-0.14,stat)(0.10,syst)							**	

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)			Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
* <sup>100</sup> Mo	T : 19Ar04=6.81(0.01,stat)(+0.38-0.40,syst)										
* <sup>100</sup> Mo	T : 17Ar18=6.90(0.15,stat)(0.37,syst) 15Ba11=7.1(0.4) (evaluation)										
* <sup>100</sup> Ag	J : 14Fe01=(5)										
* <sup>100</sup> Sn	T : also 12Hi07=1.16(0.20) 08Ba53=0.55(+0.70-0.31) 96Ki23=0.94(+0.54-0.26)										
<sup>101</sup> Kr	-28580#	500#			9# ms >400ns	5/2 <sup>+</sup> #	10	100h02	I	2010	$\beta^-$ ?; $\beta^-_n$ ?; $\beta^-_n 2n$ ?
<sup>101</sup> Rb	-42567	20			31.8 ms 3.3	3/2 <sup>+</sup> #	06	11Ni01	T	1992	$\beta^-$ =100; $\beta^-_n$ =28 4; $\beta^-_n 2n$ ?
<sup>101</sup> Sr	-55325	8			113.7 ms 1.7	(5/2 <sup>-</sup> )	06	02Pf04	D	1983	$\beta^-$ =100; $\beta^-_n$ =2.75 35
<sup>101</sup> Y	-65055	7			426 ms 20	5/2 <sup>+</sup> *	06	02Pf04	D	1983	$\beta^-$ =100; $\beta^-_n$ =2.3 8
<sup>101</sup> Y <sup>m</sup>	-63850	7	1205.0	1.0	870 ns 90	13/2 <sup>-</sup> #		09Fo05	ETD	2009	IT=100
<sup>101</sup> Zr	-73161	8			2.29 s 0.08	3/2 <sup>+</sup> *	06			1972	$\beta^-$ =100
<sup>101</sup> Nb	-78891	4			7.1 s 0.3	5/2 <sup>+</sup> *	06			1970	$\beta^-$ =100
<sup>101</sup> Mo	-83520.0	0.3			14.61 m 0.03	1/2 <sup>+</sup>	06			1941	$\beta^-$ =100
<sup>101</sup> Mo <sup>m</sup>	-83506.5	0.3	13.497	0.009	226 ns 7	3/2 <sup>+</sup>	06			1977	IT=100
<sup>101</sup> Mo <sup>o</sup>	-83463.0	0.3	57.015	0.011	133 ns 70	5/2 <sup>+</sup>	06			1977	IT=100
<sup>101</sup> Tc	-86345	24			14.22 m 0.01	9/2 <sup>+</sup>	06			1941	$\beta^-$ =100
<sup>101</sup> Tc <sup>m</sup>	-86137	24	207.526	0.020	636 $\mu$ s 8	1/2 <sup>-</sup>	06			1964	IT=100
<sup>101</sup> Ru	-87958.1	0.4			STABLE	5/2 <sup>+</sup> *	06			1931	IS=17.06 2
<sup>101</sup> Ru <sup>m</sup>	-87430.5	0.4	527.56	0.10	17.5 $\mu$ s 0.4	11/2 <sup>-</sup>	06			1974	IT=100
<sup>101</sup> Rh	-87412	6			4.07 y 0.05	1/2 <sup>-</sup>	06	18Sh09	T	1948	$\epsilon$ =100
<sup>101</sup> Rh <sup>m</sup>	-87255	6	157.32	0.03	4.343 d 0.010	9/2 <sup>+</sup> *	06			1944	$\epsilon$ =92.80 25; IT=7.20 25
<sup>101</sup> Pd	-85432	5			8.47 h 0.06	5/2 <sup>+</sup>	06			1948	$\beta^+$ =100
<sup>101</sup> Ag	-81334	5			11.1 m 0.3	9/2 <sup>+</sup> *	06			1966	$\beta^+$ =100
<sup>101</sup> Ag <sup>m</sup>	-81060	5	274.1	0.3	3.10 s 0.10	(1/2 <sup>-</sup> )*	06			1975	IT=100
<sup>101</sup> Cd	-75836.5	1.5			1.36 m 0.05	5/2 <sup>+</sup> *	06	18Yo07	J	1969	$\beta^+$ =100
<sup>101</sup> In	-68545	12			15.1 s 1.1	(9/2 <sup>+</sup> )	06	19Pa16	D	1988	$\beta^+$ =100; $\beta^+$ p<1.7
<sup>101</sup> In <sup>m</sup>	-67910	40	640	40	MD	10# s		20Ho03	E	2019	$\beta^+$ ?; IT ?
<sup>101</sup> Sn	-60310	300			2.22 s 0.05	(7/2 <sup>+</sup> )	07	19Pa16	TD	1994	$\beta^+$ =100; $\beta^+$ p=21.0 7
* <sup>101</sup> Rb	T : average 11Ni01=31(+5-4) 95Lh04=32(5)										
* <sup>101</sup> Sr	T : average 11Ni01=113(2) 86Wa17=114(4) 83Wo10=121(6) 87PfZX=104(15)										
* <sup>101</sup> Y	T : average 96Me09=400(20) 86Wa17=440(20) 83Wo10=500(50); others										
* <sup>101</sup> Y	T : 09Pe06=510(+76-67) 12Qu01=480(+143-114) 93Ru01=279(9), outlier										
* <sup>101</sup> Y	J : 07Ch07=5/2										
* <sup>101</sup> Y <sup>m</sup>	T : symmetrized from 09Fo05=860(+90-80); other 12Ka36=187(+49-38)										
* <sup>101</sup> Y <sup>m</sup>	E : E(13/2 <sup>+</sup> )=724.98(10) keV from 05Lu21 + 480(1) keV from 09Fo05										
* <sup>101</sup> Y <sup>m</sup>	I : 09Fo05=129,164,204,230 and 480 gamma rays in a cascade to gs, the first										
* <sup>101</sup> Y <sup>m</sup>	I : four in agreement with SF data of 05Lu21; other 12Ka36=128.0(0.5) and										
* <sup>101</sup> Y <sup>m</sup>	I : 203.5(0.5) gamma rays in a cascade to gs, but limited statistics										
* <sup>101</sup> Zr	T : average 19Do02=2.27(0.12) 72Th08=2.3(0.1)										
* <sup>101</sup> Zr	J : 02Ca37=3/2										
* <sup>101</sup> Nb	J : 09Ch25=5/2										
* <sup>101</sup> Rh <sup>m</sup>	T : average 68Li08=4.39(0.08) 66Ar05=4.34(0.01)										
* <sup>101</sup> Rh <sup>m</sup>	T : 65Er04=4.43(0.06); Birge ratio=3.22										
* <sup>101</sup> Ag	J : other 14Fe01=9/2										
* <sup>101</sup> Ag <sup>m</sup>	J : 14Fe01=(1/2)										
* <sup>101</sup> In	T : average 97Sz04=14.9(1.2) 88Hu07=16(3)										
* <sup>101</sup> In <sup>m</sup>	E : average 20Ho03=608(57) 19Xu13=659(50)										
* <sup>101</sup> Sn	D : % $\beta^+$ p average 19Pa16=23.6(0.8) 12Lo08=22(1) 10St.A=20(1)										
* <sup>101</sup> Sn	J : from 10Da17										
<sup>102</sup> Rb	-37250	80			37 ms 4	(4 <sup>+</sup> )	09	16Wa16	JD	1995	$\beta^-$ =100; $\beta^-_n$ =65 22; $\beta^-_n 2n$ ?
<sup>102</sup> Sr	-52160	70			69 ms 6	0 <sup>+</sup>	09			1986	$\beta^-$ =100; $\beta^-_n$ =5.5 15
<sup>102</sup> Y	-61173	4		*	360 ms 40	(5 <sup>-</sup> )	09	17Br12	ID	1980	$\beta^-$ =100; $\beta^-_n$ <2.6
<sup>102</sup> Y <sup>m</sup>	-61070#	100#	100#	100#	300 ms 100	(0 <sup>-</sup> , 1 <sup>-</sup> )	09			1983	$\beta^-$ =100; $\beta^-_n$ <2.6; IT ?
<sup>102</sup> Zr	-71581	9			2.01 s 0.08	0 <sup>+</sup>	09	19Do02	T	1970	$\beta^-$ =100
<sup>102</sup> Nb	-76298.3	2.5			4.3 s 0.4	(4 <sup>+</sup> )	09			1972	$\beta^-$ =100
<sup>102</sup> Nb <sup>m</sup>	-76204	8	94	7	MD	(1 <sup>+</sup> )	09	19Do02	T	1976	$\beta^-$ =100; IT ?
<sup>102</sup> Mo	-83561	8			11.3 m 0.2	0 <sup>+</sup>	09			1954	$\beta^-$ =100
<sup>102</sup> Tc	-84573	9		*	5.28 s 0.15	1 <sup>+</sup>	09			1954	$\beta^-$ =100
<sup>102</sup> Tc <sup>m</sup>	-84520#	50#	50#	50#	4.35 m 0.07	(4 <sup>+</sup> )	09			1954	$\beta^-$ ≈100; IT ?
<sup>102</sup> Ru	-89106.4	0.4			STABLE	0 <sup>+</sup>	09			1931	IS=31.55 14

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{102}\text{Rh}$	-86783	6	207.0 d 1.5	$2^-*$	09 98Sh21 T	1941	$\beta^+=78.5; \beta^-=22.5$	*
$^{102}\text{Rh}^m$	-86642	6	140.73 0.09	3.742 y 0.010	09 99Gi14 J	1962	$\beta^+\approx 100; IT=0.233.24$	
$^{102}\text{Pd}$	-87903.0	0.4	STABLE	$>7.6\text{E}y$	09 16Le16 T	1935	$IS=1.02.1; 2\beta^+?$	
$^{102}\text{Ag}$	-82247	8	12.9 m 0.3	$0^+$	09	1960	$\beta^+=100$	
$^{102}\text{Ag}^m$	-82238	8	9.40 0.07	7.7 m 0.5	09	1967	$\beta^+=51.5; IT=49.5$	
$^{102}\text{Cd}$	-79659.7	1.7	5.5 m 0.5	$0^+$	09	1969	$\beta^+=100$	
$^{102}\text{In}$	-70695	5	23.3 s 0.1	$(6^+)$	09 95Sz01 J	1981	$\beta^+=100; \beta^+p=0.0093.13$	
$^{102}\text{Sn}$	-64930	100	3.8 s 0.2	$0^+$	09	1994	$\beta^+=100$	*
$^{102}\text{Sn}^m$	-62910	100	2017 2	367 ns 8	09 98Li50 E	1996	$IT=100$	*
$^{102}\text{Sb}$	-51100#	400#					p?	
* $^{102}\text{Rb}$	T : average 15Lo04=37(10) 11Ni01=35(+15-8) 87PfZX=37(5)							**
* $^{102}\text{Rb}$	D : other $\% \beta^-n=18(8)$ in 87PfZX							**
* $^{102}\text{Sr}$	T : also 11Ni01=85(15)							**
* $^{102}\text{Y}$	J : direct $\beta^-$ feeding of 4- and 5- levels in $^{102}\text{Zr}$ in 17Br12;							**
* $^{102}\text{Y}$	J : p5/2[422]n5/2[532], K=5- configuration from systematics							**
* $^{102}\text{Y}$	T : from 91Hi02 for the high-spin $\beta^-$ decaying state							**
* $^{102}\text{Y}$	D : from $\% \beta^-n=4.9.12$ , average 86ReZS=6.0(1.7) 96Me09=4.0(1.5),							**
* $^{102}\text{Y}$	D : and by splitting equally between gs and isomer							**
* $^{102}\text{Y}^m$	J : direct $\beta^-$ feeding of $^{102}\text{Sr}$ ( $0^+$ ) - see discussion in 91Hi02;							**
* $^{102}\text{Y}^m$	J : p5/2[422]n5/2[532], K=0- configuration from systematics; other							**
* $^{102}\text{Y}^m$	J : 07Ch07=(2,3)							**
* $^{102}\text{Y}^m$	T : from 91Hi02 for the low-spin $\beta^-$ decaying state							**
* $^{102}\text{Y}^m$	D : from $\% \beta^-n=4.9.12$ , average 86ReZS=6.0(1.7) 96Me09=4.0(1.5),							**
* $^{102}\text{Y}^m$	D : and by splitting equally between gs and isomer							**
* $^{102}\text{Nb}^m$	T : average 19Do02=1.33(0.27) 76Ah06=1.3(0.2)							**
* $^{102}\text{Tc}^m$	J : direct $\beta^-$ feeding of J=4 and 5 levels and the lack of such to the 6+							**
* $^{102}\text{Tc}^m$	J : levels in $^{102}\text{Ru}$ in 70Hu02 and 69B116							**
* $^{102}\text{Rh}$	T : average 98Sh21=207.3(1.7) 61Hi06=206(3)							**
* $^{102}\text{Pd}$	T : 16Le16 (supersedes 13Le10) $>8.8\text{E}y$ , $>7.6\text{E}y$ for the first excited							**
* $^{102}\text{Pd}$	T : $0^+$ and $2^+$ states, and $>14\text{E}y$ for the second excited $2^+$							**
* $^{102}\text{Sn}$	T : 95Fa.A=4.6(1.4), supersedes 95Sc28=4.5(0.7) from the same group							**
* $^{102}\text{Sn}^m$	T : from 11Hi.A							**
$^{103}\text{Rb}$	-33160#	400#	26 ms 11	$3/2^+\#$	15 15Lo04 TD	2010	$\beta^-=100; \beta^-n?; \beta^-2n?$	*
$^{103}\text{Sr}$	-47280#	200#	53 ms 10	$5/2^+\#$	15	1997	$\beta^-=100; \beta^-n?; \beta^-2n?$	
$^{103}\text{Y}$	-58457	11	239 ms 12	$5/2^+\#$	09 11Ni01 T	1994	$\beta^-=100; \beta^-n=8.0.17$	*
$^{103}\text{Zr}$	-67809	9	1.38 s 0.07	$(5/2^-)$	09 09Pe06 TD	1987	$\beta^-=100; \beta^-n<1$	
$^{103}\text{Nb}$	-75029	4	1.34 s 0.07	$5/2^+*$	09 19Do02 T	1971	$\beta^-=100; \beta^-n?$	*
$^{103}\text{Mo}$	-80954	9	67.5 s 1.5	$3/2^+$	09 09Ch09 J	1963	$\beta^-=100$	
$^{103}\text{Tc}$	-84604	10	54.2 s 0.8	$5/2^+$	09	1957	$\beta^-=100$	
$^{103}\text{Ru}$	-87267.2	0.4	39.245 d 0.008	$3/2^+$	09 FGK204 T	1945	$\beta^-=100$	
$^{103}\text{Ru}^m$	-87029.0	0.8	238.2 0.7	11/2 $^-$	09	1964	$IT=100$	
$^{103}\text{Rh}$	-88031.7	2.3	STABLE	$1/2^-*$	09	1934	$IS=100$	
$^{103}\text{Rh}^m$	-87991.9	2.3	39.753 0.006	56.114 m 0.009	09	1943	$IT=100$	
$^{103}\text{Pd}$	-87457.0	0.9	16.991 d 0.019	$5/2^+$	09	1950	$\epsilon=100$	
$^{103}\text{Ag}$	-84803	4	65.7 m 0.7	$7/2^+*$	09	1954	$\beta^+=100$	
$^{103}\text{Ag}^m$	-84669	4	134.45 0.04	5.7 s 0.3	1/2 $^-$	09	$IT=100$	
$^{103}\text{Cd}$	-80651.6	1.8	7.3 m 0.1	$5/2^+*$	09 18Yo07 J	1960	$\beta^+=100$	
$^{103}\text{In}$	-74632	9	60 s 1	$(9/2^+)$	09 97Sz04 T	1978	$\beta^+=100$	
$^{103}\text{In}^m$	-74000	9	631.7 0.1	34 s 2	$(1/2^-)$	09	$\beta^+=67; IT=33$	*
$^{103}\text{Sn}$	-67090#	100#	7.0 s 0.2	$5/2^+\#$	09	1981	$\beta^+=100; \beta^+p=1.2.1$	
$^{103}\text{Sb}$	-56670#	300#	$<49\text{ns}$	$5/2^+\#$	15 13Su23 I	2010	p?	
* $^{103}\text{Rb}$	T : symmetrized from 15Lo04=23(+13-9)							**
* $^{103}\text{Y}$	T : average 11Ni01=234(+18-15) 09Pe06=260(+40-20) 96Me09=230(20)							**
* $^{103}\text{Y}$	T : 96Lh04=190(50)							**
* $^{103}\text{Y}$	D : $\% \beta^-n$ average 09Pe06=8(2) 96Me09=8(3)							**
* $^{103}\text{Nb}$	J : 09Ch25=5/2							**
* $^{103}\text{In}^m$	E : other 20Ho03=689(77)							**
$^{104}\text{Rb}$	-27450#	500#	35# ms $>550\text{ns}$		18Sh11 IT	2018	$\beta^-?; \beta^-n?; \beta^-2n?$	
$^{104}\text{Sr}$	-43760#	300#	50.6 ms 4.2	$0^+$	15 15Lo04 T	1997	$\beta^-=100; \beta^-n?; \beta^-2n?$	*

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{104}\text{Y}$	-54080#	200#	197 ms 4	$(0^+, 1^+)^\#$	15 09Pe06	D 1994	$\beta^- = 100; \beta^-_n = 34\ 10; \beta^- 2n ?$ *	
$^{104}\text{Zr}$	-65718	9	920 ms 28	$0^+$	07 09Pe06	TD 1990	$\beta^- = 100; \beta^-_n < 1$	
$^{104}\text{Nb}$	-71811.0	1.8	0.98 s 0.05	$(5^-)$	07	1976	$\beta^- = 100; \beta^-_n = 0.05\ 3$ *	
$^{104}\text{Nb}^m$	-71801.2	1.9	4.9 s 0.3	$(0^-, 1^-)$	07 FGK207	J 1971	$\beta^- = 100; \beta^-_n = 0.06\ 3$ *	
$^{104}\text{Mo}$	-80344	9	60 s 2	$0^+$	07	1962	$\beta^- = 100$	
$^{104}\text{Tc}$	-82499	25	18.3 m 0.3	$(3^-)$	07	1956	$\beta^- = 100$ *	
$^{104}\text{Tc}^m$	-82429	25	69.7 0.2	$(5^-)$	07	1981	IT=100 *	
$^{104}\text{Tc}^n$	-82393	25	106.1 0.3	400 ns 20	4#	07	1999	IT=100
$^{104}\text{Ru}$	-88095.8	2.5	STABLE	$0^+$	07	1931	IS=18.62 27; $2\beta^- ?$ *	
$^{104}\text{Rh}$	-86959.3	2.3	42.3 s 0.4	$1^+$	07	1939	$\beta^- = 99.55\ 10; \beta^+ = 0.45\ 10$	
$^{104}\text{Rh}^m$	-86830.3	2.3	128.9679 0.0005	$5^+$	07	1939	IT=99.87 1; $\beta^- = 0.13\ 1$	
$^{104}\text{Pd}$	-89395.1	1.3	STABLE	$0^+$	07	1935	IS=11.14 8	
$^{104}\text{Ag}$	-85116	4	69.2 m 1.0	$5^+*$	07	1955	$\beta^+ = 100$	
$^{104}\text{Ag}^m$	-85109	4	6.90 0.22	$2^+*$	07	1959	$\beta^+ \approx 100; IT < 0.07$	
$^{104}\text{Cd}$	-83968.4	1.7	57.7 m 1.0	$0^+$	07	1955	$\beta^+ = 100$	
$^{104}\text{In}$	-76183	6	1.80 m 0.03	$(5^+)*$	07	1977	$\beta^+ = 100$	
$^{104}\text{In}^m$	-76090	6	93.48 0.10	$(3^+)$	07 89Va05	D 1988	IT=80 5; $\beta^+ = 20\ 5$	
$^{104}\text{Sn}$	-71627	6	20.8 s 0.5	$0^+$	07	1985	$\beta^+ = 100$	
$^{104}\text{Sb}$	-59300#	100#	470 ms 130		07 96FaZZ	TD 1995	$\beta^+ = ?; \beta^+ p < 7; p < 7; \alpha ?$ *	
$^{104}\text{Te}$	-49630	320	< 4 ns	$0^+$	18Au04	D 2018	$\alpha = 100$ *	
* $^{104}\text{Sr}$	T : average 15Lo04=53(5) 11Ni01=43(+9-7) **							
* $^{104}\text{Y}$	T : average 15Lo04=198(20) 11Ni01=197(4) 99Wa09=180(60); other **							
* $^{104}\text{Y}$	T : 09Pe06=260(+60-50) 99Wa09=180(60) **							
* $^{104}\text{Nb}$	T : average 19Do02=0.97(0.10) 96Me09=1.0(0.1) 82Ke05=0.99(0.07) **							
* $^{104}\text{Nb}$	T : 76Ah06=0.8(0.2); other 80BaZL=0.91, no uncertainty quoted **							
* $^{104}\text{Nb}^m$	D : % $\beta^-_n$ other 83En03=0.71%, conflicting (not used) **							
* $^{104}\text{Tc}$	J : strong $\beta^-$ feeding to 2+, 2- and 4+ levels in $^{104}\text{Ru}$ ; **							
* $^{104}\text{Tc}$	J : expected conf=p3/2[301] n3/2[411], K=3- **							
* $^{104}\text{Tc}^m$	J : E2 gamma to (3-) level (from Ensdf2007) **							
* $^{104}\text{Ru}$	T : Onu-BB to 1st exc. state : 13Be09>650Ey 12An08>190Ey **							
* $^{104}\text{Sb}$	T : symmetrized from 96FaZZ=440(+150-110), supersedes 95Sc28, **							
* $^{104}\text{Sb}$	T : 95Sc33=520(+180,-130) **							
* $^{104}\text{Sb}$	D : %p from 96FaZZ, supersedes 95Sc28<1% **							
* $^{104}\text{Te}$	T : from 19Xi06; other 18Au04<18 ns **							
$^{105}\text{Sr}$	-38190#	500#	39 ms 5	$5/2^+^\#$	19	1997	$\beta^- = 100; \beta^-_n ?; \beta^- 2n ?$	
$^{105}\text{Y}$	-50570#	400#	95 ms 9	$5/2^+^\#$	19 09Pe06	D 1994	$\beta^- = 100; \beta^-_n < 82; \beta^- 2n ?$ *	
$^{105}\text{Zr}$	-61458	12	670 ms 28	$1/2^+^\#$	19 09Pe06	TD 1992	$\beta^- = 100; \beta^-_n < 2$ *	
$^{105}\text{Nb}$	-69916	4	2.91 s 0.05	$(5/2^+)$	19	1984	$\beta^- = 100; \beta^-_n = 1.7\ 9$	
$^{105}\text{Mo}$	-77331	9	36.3 s 0.8	$(5/2^-)$	19	1962	$\beta^- = 100$	
$^{105}\text{Tc}$	-82290	40	7.64 m 0.06	$(3/2^-)$	19	1955	$\beta^- = 100$	
$^{105}\text{Ru}$	-85934.5	2.5	4.439 h 0.011	$3/2^+$	19	1945	$\beta^- = 100$	
$^{105}\text{Ru}^m$	-85913.9	2.5	20.606 0.014	340 ns 15	5/2+	19	1974	IT=100
$^{105}\text{Rh}$	-87851.3	2.5	35.341 h 0.019	$7/2^+$	19	1945	$\beta^- = 100$	
$^{105}\text{Rh}^m$	-87721.6	2.5	129.742 0.004	42.8 s 0.3	1/2-	19	1950	IT=100
$^{105}\text{Pd}$	-88417.9	1.1	STABLE	$5/2^+*$	19	1935	IS=22.33 8	
$^{105}\text{Pd}^m$	-87928.8	1.1	489.1 0.3	35.5 $\mu\text{s}$ 0.5	11/2-	19	1970	IT=100
$^{105}\text{Ag}$	-87071	5	41.29 d 0.07	$1/2^-*$	19	1939	$\beta^+ = 100$	
$^{105}\text{Ag}^m$	-87046	5	25.468 0.016	7.23 m 0.16	7/2+	19	1969	IT=99.66 7; $\beta^+ = 0.34\ 7$
$^{105}\text{Cd}$	-84333.8	1.4	55.5 m 0.4	$5/2^+*$	19	1950	$\beta^+ = 100$ *	
$^{105}\text{Cd}^m$	-81816.2	1.5	2517.6 0.5	4.5 $\mu\text{s}$ 0.5	(21/2+)	19	1976	IT=100
$^{105}\text{In}$	-79641	10	5.07 m 0.07	$9/2^+*$	19	1975	$\beta^+ = 100$	
$^{105}\text{In}^m$	-78967	10	674.09 0.25	48 s 6	(1/2)-	19	1975	IT $\approx$ 100; $\beta^+ ?$ *
$^{105}\text{Sn}$	-73338	4	32.7 s 0.5	$(5/2^+)$	19	1981	$\beta^+ = 100; \beta^+ p = 0.011\ 4$	
$^{105}\text{Sb}$	-64015	22	1.12 s 0.16	$(5/2^+)$	05 96FaZZ	T 1994	$\beta^+ = 100; p < 0.1; \beta^+ p ?$ *	
$^{105}\text{Te}$	-52810	300	633 ns 66	$(7/2^+)$	06 06Se08	T 2006	$\alpha \approx 100$ *	
* $^{105}\text{Y}$	T : symmetrized from 15Lo04=107(+6-9); others 11Ni01=83(+5-4) **							
* $^{105}\text{Y}$	T : 09Pe06=160(+85-60) **							
* $^{105}\text{Zr}$	J : 20Ur02=1/2+, 1/2+[411] **							
* $^{105}\text{Cd}$	J : also 18Yo07=5/2 **							
* $^{105}\text{In}^m$	E : other 20Ho03=702(27) **							
* $^{105}\text{Sb}$	T : from 96FaZZ, supersedes 95Sc28=1.30(0.15) (preliminary, the same group) **							

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)		
* <sup>105</sup> Sb	D : %p from 05Li47<0.1% above 430 keV, disagrees with 96FaZZ,94Ti03 1%							**	
* <sup>105</sup> Te	T : average 06Li41=620(70) 06Se08=700(+250-170)							**	
* <sup>105</sup> Te	J : favorite $\alpha$ decay to the 171.7-keV state [ $J=(7/2^+)$ ] in <sup>101</sup> Sn							**	
<sup>106</sup> Sr	-34300#	600#	21 ms 8	0 <sup>+</sup>	15 15Lo04 T	2010	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$	*	
<sup>106</sup> Y	-45790#	500#	75 ms 6	2 <sup>+</sup> #	15 15Lo04 T	1997	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$	*	
<sup>106</sup> Zr	-58750#	200#	179 ms 6	0 <sup>+</sup>	15 15Lo04 T	1994	$\beta^- = 100; \beta^- n < 7$	*	
<sup>106</sup> Nb	-66202.7	1.4	900 ms 20	1 <sup>-</sup> #	15 14Lu07 J	1976	$\beta^- = 100; \beta^- n = 4.5 \ 3$	*	
<sup>106</sup> Nb <sup>m</sup>	-66100#	50#	100# 50#	1.20 s 0.06	(4 <sup>-</sup> )	20Ha14 TJ	1976	$\beta^- = 100; IT ?$	*
<sup>106</sup> Nb <sup>n</sup>	-65997.9	1.5	204.8 0.5	820 ns 38	(3 <sup>+</sup> )	14Lu07 EJ	1999	IT=100	*
<sup>106</sup> Mo	-76128	9	8.73 s 0.12	0 <sup>+</sup>	08	1969	$\beta^- = 100$		
<sup>106</sup> Tc	-79776	12	35.6 s 0.6	(1,2)( <sup>+</sup> #)	08	1965	$\beta^- = 100$		
<sup>106</sup> Ru	-86323	5	371.8 d 0.18	0 <sup>+</sup>	08	1948	$\beta^- = 100$		
<sup>106</sup> Rh	-86363	5	30.07 s 0.35	1 <sup>+</sup>	08	1947	$\beta^- = 100$		
<sup>106</sup> Rh <sup>m</sup>	-86231	10	132 11 BD	131 m 2	(6 <sup>+</sup> )	08	1955	$\beta^- = 100$	
<sup>106</sup> Pd	-89907.5	1.1	STABLE	0 <sup>+</sup>	08	1935	IS=27.33 3		
<sup>106</sup> Ag	-86942	3	23.96 m 0.04	1 <sup>+</sup> *	08	1937	$\beta^+ \approx 100; \beta^- ?$		
<sup>106</sup> Ag <sup>m</sup>	-86852	3	89.66 0.07	8.28 d 0.02	6 <sup>+</sup> *	08	1938	$\beta^+ = 100; IT ?$	
<sup>106</sup> Cd	-87132.2	1.1	STABLE	>1.1Zy	0 <sup>+</sup>	08 16Be11 T	1935	IS=1.245 22; $2\beta^+ ?$	*
<sup>106</sup> In	-80608	12	6.2 m 0.1	7 <sup>+</sup> *	08	1962	$\beta^+ = 100$		
<sup>106</sup> In <sup>m</sup>	-80579	12	28.6 0.3	5.2 m 0.1	(2 <sup>+</sup> )	08	1966	$\beta^+ = 100$	
<sup>106</sup> Sn	-77354	5	1.92 m 0.08	0 <sup>+</sup>	08	1975	$\beta^+ = 100$		
<sup>106</sup> Sb	-66473	7	600 ms 200	(2 <sup>+</sup> )	08	1981	$\beta^+ = 100$		
<sup>106</sup> Sb <sup>m</sup>	-66370	7	103.5 0.3	226 ns 14	(4 <sup>+</sup> )	08 99So08 T	1998	IT=100	*
<sup>106</sup> Te	-58220	100	78 $\mu$ s 11	0 <sup>+</sup>	08 16Ca33 T	1981	$\alpha = 100$	*	
<sup>106</sup> I	-43300#	400#					$\alpha ?$		
* <sup>106</sup> Sr	T : symmetrized from 15Lo04=20(+8-7)							**	
* <sup>106</sup> Y	T : average 15Lo04=82(+10-5) 15NiZZ=62(9); other 11Ni01=62(+25-14)							**	
* <sup>106</sup> Zr	T : average 15Lo04=175(7) 11Ni01=186(+11-10)							**	
* <sup>106</sup> Nb	T : from 96Me09 using $\beta^-$ (0), predominantly from the low-spin $\beta^-$							**	
* <sup>106</sup> Nb	T : decaying state							**	
* <sup>106</sup> Nb <sup>m</sup>	T : average 20Ha14=1.10(0.05) 09Pe06=1.24(0.02) 83Sh06=1.02(0.05);							**	
* <sup>106</sup> Nb <sup>m</sup>	T : Birge ratio=3.2; contain contributions from the shorter gs							**	
* <sup>106</sup> Nb <sup>n</sup>	T : average 12Ka36=660(+110-100) 99Ge01=840(40)							**	
* <sup>106</sup> Cd	T : for $2\nu\text{-}\epsilon\beta^+$ , theoretically fastest channel; others 12Be14>210Ey							**	
* <sup>106</sup> Cd	T : 02Tr04>410Ey							**	
* <sup>106</sup> Sb <sup>m</sup>	T : average 99So08=232(21) 98Li50=220(20)							**	
* <sup>106</sup> Te	T : average 16Ca33=70(+20-15) 05Ja03=85(+25-15) 94Pa11=60(+40-20) and							**	
* <sup>106</sup> Te	T : 81Sc17=60(+30-10)							**	
<sup>107</sup> Sr	-28250#	700#	25# ms >400ns	1/2 <sup>+</sup> #	10 10Oh02 I	2010	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$		
<sup>107</sup> Y	-41970#	500#	33.5 ms 0.3	5/2 <sup>+</sup> #	15 15Lo04 T	1997	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$	*	
<sup>107</sup> Zr	-54020#	300#	145.7 ms 2.4	5/2 <sup>+</sup> #	15 15Lo04 T	1994	$\beta^- = 100; \beta^- n < 23$	*	
<sup>107</sup> Nb	-63724	8	286 ms 8	(5/2 <sup>+</sup> )	08 19Ku16 TJ	1992	$\beta^- = 100; \beta^- n = 7.4 \ 8$	*	
<sup>107</sup> Mo	-72545	9	3.5 s 0.5	(1/2 <sup>+</sup> )	08 19Ku16 J	1972	$\beta^- = 100$	*	
<sup>107</sup> Mo <sup>m</sup>	-72480	9	65.4 0.2	445 ns 21	(5/2 <sup>+</sup> )	08 19Ku16 J	1976	IT=100	*
<sup>107</sup> Tc	-78750	9	21.2 s 0.2	(3/2 <sup>-</sup> )	08 09Gu11 J	1965	$\beta^- = 100$		
<sup>107</sup> Tc <sup>m</sup>	-78720	9	30.1 0.1	3.85 $\mu$ s 0.05	(1/2 <sup>+</sup> )	08	2007	IT=100	
<sup>107</sup> Tc <sup>n</sup>	-78684	9	65.72 0.14	184 ns 3	(5/2 <sup>+</sup> )	08	1974	IT=100	
<sup>107</sup> Ru	-83863	9	3.75 m 0.05	(5/2 <sup>+</sup> )	08	1951	$\beta^- = 100$		
<sup>107</sup> Rh	-86864	12	21.7 m 0.4	7/2 <sup>+</sup>	08	1951	$\beta^- = 100$		
<sup>107</sup> Rh <sup>m</sup>	-86596	12	268.36 0.04	> 10 $\mu$ s	1/2 <sup>-</sup>	08	1986	IT=100	
<sup>107</sup> Pd	-88372.7	1.2	6.5 My 0.3	5/2 <sup>+</sup>	08	1958	$\beta^- = 100$		
<sup>107</sup> Pd <sup>m</sup>	-88257.0	1.2	115.74 0.12	850 ns 100	1/2 <sup>+</sup>	08	1969	IT=100	
<sup>107</sup> Pd <sup>n</sup>	-88158.1	1.2	214.6 0.3	21.3 s 0.5	11/2 <sup>-</sup>	08	1952	IT=100	
<sup>107</sup> Ag	-88406.7	2.4	STABLE	1/2 <sup>-</sup> *	08	1924	IS=51.839 8	*	
<sup>107</sup> Ag <sup>m</sup>	-88313.6	2.4	93.125 0.019	44.3 s 0.2	7/2 <sup>+</sup>	08	1940	IT=100	
<sup>107</sup> Cd	-86990.3	1.7	6.50 h 0.02	5/2 <sup>+</sup> *	08	1946	$\beta^+ = 100$	*	
<sup>107</sup> In	-83567	10	32.4 m 0.3	9/2 <sup>+</sup> *	08	1949	$\beta^+ = 100$		
<sup>107</sup> In <sup>m</sup>	-82889	10	678.5 0.3	50.4 s 0.6	1/2 <sup>-</sup>	08	1973	IT=100	*
<sup>107</sup> Sn	-78512	5	2.90 m 0.05	(5/2 <sup>+</sup> )	08	1976	$\beta^+ = 100$		

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{107}\text{Sb}$	-70653	4	4.0 s 0.2	$5/2^+\#$	08	1994	$\beta^+=100$	
$^{107}\text{Te}$	-60660#	100#	3.22 ms 0.09	$5/2^+\#$	08 19Au02	T 1979	$\alpha=70\ 30;\beta^+\ ?; \beta^+\text{p}\ ?$	*
$^{107}\text{I}$	-49430#	300#	20# $\mu\text{s}$	$5/2^+\#$			$\alpha\ ?$	
* $^{107}\text{Y}$	T : other 11Ni01=41(+15-9)							**
* $^{107}\text{Zr}$	T : average 15Lo04=150(3) 11Ni01=138(4); not used 09Pe06=150(+40-30)							**
* $^{107}\text{Nb}$	T : average 19Ku16=270(20) 15Lo04=280(20) 09Pe06=290(11) 96Me09=300(30)							**
* $^{107}\text{Nb}$	D : % $\beta^-$ -n average 09Pe06=8(1) 96Me09=6.0(1.5)							**
* $^{107}\text{Mo}$	J : 20Ur02,19Ku16=1/2+, 1/2+[411]							**
* $^{107}\text{Mo}^m$	T : average 06Pi14=420(30) 99Ge01=470(30); others 20Ur02=600(150)							**
* $^{107}\text{Mo}^m$	T : $^{76}\text{ChZD}=238(7)$							**
* $^{107}\text{Mo}^m$	J : 20Ur02,19Ku16=5/2+, 5/2+[413]							**
* $^{107}\text{Ag}$	J : also 14Fe01=1/2							**
* $^{107}\text{Cd}$	J : also 18Yo07,13Yo02=5/2							**
* $^{107}\text{In}^m$	E : other 20Ho03=663(22)							**
* $^{107}\text{Te}$	T : average 19Au02=3.6(0.2) 14Pa11=3.1(0.1) 79Sc22=3.6(+0.6-0.4)							**
$^{108}\text{Y}$	-36780#	600#	30 ms 5	$6^-$	15 15Lo04	T 2010	$\beta^-=100;\beta^-n\ ?; \beta^-2n\ ?$	*
$^{108}\text{Zr}$	-50950#	400#	78.5 ms 2.0	$0^+$	15	1997	$\beta^-=100;\beta^-n\ ?$	
$^{108}\text{Zr}^m$	-48880#	400#	2074.5 0.8	$540\ \text{ns}\ 30$	( $6^+$ )	15 12Ka36	T 2011	IT=100
$^{108}\text{Nb}$	-59545	8	201 ms 4	$(2^+)$	15 20Ha14	T 1994	$\beta^-=100;\beta^-n=6.3\ 5;\beta^-2n\ ?$	*
$^{108}\text{Nb}^m$	-59378	8	166.6 0.5	109 ns 2	$6^-$	15 12Ka36	DT 2012	IT=100
$^{108}\text{Mo}$	-70749	9	1.105 s 0.010	$0^+$	08 09Pe06	TD 1972	$\beta^-=100;\beta^-n<0.5$	*
$^{108}\text{Tc}$	-75923	9	5.17 s 0.07	( $2^+$ )	08	1970	$\beta^-=100$	
$^{108}\text{Ru}$	-83661	9	4.55 m 0.05	$0^+$	08	1955	$\beta^-=100$	
$^{108}\text{Rh}$	-85031	14	16.8 s 0.5	$1^+$	08	1955	$\beta^-=100$	
$^{108}\text{Rh}^m$	-84917	12	115 18 MD	6.0 m 0.3	( $5^+$ )	08	1969	$\beta^-=100$
$^{108}\text{Pd}$	-89524.2	1.1	STABLE	$0^+$	08	1935	IS=26.46 9	
$^{108}\text{Ag}$	-87606.8	2.4	2.382 m 0.011	$1^+$ *	08	1937	$\beta^-=97.15\ 20;\beta^+=2.85\ 20$	
$^{108}\text{Ag}^m$	-87497.3	2.4	109.466 0.007	439 y 9	$6^+$ *	08 18Sh09	T 1969	$\beta^+=91.3\ 9;\text{IT}=8.7\ 9$
$^{108}\text{Cd}$	-89252.4	1.1	STABLE	>410Py	$0^+$	08 95Ge14	T 1935	IS=0.888 11; $2\beta^+\ ?$
$^{108}\text{In}$	-84120	9	58.0 m 1.2	$7^+$ *	08	1949	$\beta^+=100$	
$^{108}\text{In}^m$	-84090	9	29.75 0.05	39.6 m 0.7	$2^+$ *	08	1955	$\beta^+=100$
$^{108}\text{Sn}$	-82070	5	10.30 m 0.08	$0^+$	08	1968	$\beta^+=100$	
$^{108}\text{Sb}$	-72445	5	7.4 s 0.3	( $4^+$ )	08	1976	$\beta^+=100$	
$^{108}\text{Te}$	-65782	5	2.1 s 0.1	$0^+$	08 85Ti02	D 1974	$\beta^+=51\ 4;\alpha=49\ 4;$ $\beta^+\text{p}=2.4\ 10;\beta^+\alpha<0.065$	
$^{108}\text{I}$	-52770#	100#	26.4 ms 0.8	$1^+\#$	08 19Au02	TD 1991	$\alpha\approx 99.50\ 21;\text{p}=0.50\ 21;$ $\beta^+\ ?; \beta^+\text{p}\ ?$	
$^{108}\text{Xe}$	-42630	380	72 $\mu\text{s}$ 35	$0^+$	18Au04	TD 2018	$\alpha=100$	*
* $^{108}\text{Y}$	T : other 11Ni01=25(+66-10)							**
* $^{108}\text{Zr}^m$	T : symmetrized from 12Ka36=536(+26-25); other 11Su11=620(150)							**
* $^{108}\text{Nb}$	T : average 20Ha14=186(8) 15Lo04=195(6) 09Pe06=210(5)							**
* $^{108}\text{Nb}$	D : % $\beta^-$ -n other 20Ha14=18(11)							**
* $^{108}\text{Mo}$	T : average 09Pe06=1.110(0.011) 95Jo02=1.090(0.020)							**
* $^{108}\text{Ag}^m$	T : average 18Sh09=448(27) 04Sh04=438(9)							**
* $^{108}\text{Xe}$	T : average 19Xi06=30(+57-12) 18Au04=58(+106-23)							**
$^{109}\text{Y}$	-32480#	700#	25 ms 5	$5/2^+\#$	16 15Lo04	T 2010	$\beta^-=100;\beta^-n\ ?; \beta^-2n\ ?$	
$^{109}\text{Zr}$	-45730#	500#	56 ms 3	$5/2^+\#$	16	1997	$\beta^-=100;\beta^-n\ ?; \beta^-2n\ ?$	
$^{109}\text{Nb}$	-56690	430	106.9 ms 4.9	$3/2^-$	16 15Lo04	T 1994	$\beta^-=100;\beta^-n=31\ 5$	*
$^{109}\text{Nb}^m$	-56380	430	312.5 0.4	115 ns 8	$7/2^+\#$	16 12Ka36	T 2011	IT=100
$^{109}\text{Mo}$	-66659	11	700 ms 14	( $1/2^+$ )	16 09Pe06	TD 1992	$\beta^-=100;\beta^-n=1.3\ 6$	*
$^{109}\text{Mo}^m$	-66589	11	69.7 0.5	210 ns 60	$5/2^+\#$	16 12Ka36	ET 2012	IT=100
$^{109}\text{Tc}$	-74283	10	905 ms 21	( $5/2^+$ )	16 19Do02	T 1976	$\beta^-=100;\beta^-n=0.08\ 2$	*
$^{109}\text{Ru}$	-80738	9	34.4 s 0.2	( $5/2^+$ )	16	1967	$\beta^-=100$	
$^{109}\text{Ru}^m$	-80642	9	96.14 0.15	680 ms 30	( $5/2^-$ )	16	1976	IT=100
$^{109}\text{Rh}$	-84999	4	80.8 s 0.7	$7/2^+$	16	1972	$\beta^-=100$	
$^{109}\text{Rh}^m$	-84773	4	225.873 0.019	1.66 $\mu\text{s}$ 0.04	$3/2^+$	16	1987	IT=100
$^{109}\text{Pd}$	-87606.5	1.1	13.59 h 0.12	$5/2^+$	16	1937	$\beta^-=100$	
$^{109}\text{Pd}^m$	-87493.1	1.1	113.4000 0.0014	380 ns 50	$1/2^+$	16	1978	IT=100
$^{109}\text{Pd}^m$	-87417.5	1.1	188.9903 0.0010	4.703 m 0.009	$11/2^-$	16	1957	IT=100



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{109}\text{Ag}$	-88719.4	1.3	STABLE	$1/2^-*$	16	1924	IS=48.161 8 *
$^{109}\text{Ag}^m$	-88631.4	1.3	88.0337 0.0010	$7/2^+*$	16	1967	IT=100
$^{109}\text{Cd}$	-88504.3	1.5	461.3 d 0.5	$5/2^+*$	16 FGK209 T	1950	$\varepsilon=100$ *
$^{109}\text{Cd}^m$	-88444.7	1.5	59.60 0.07	$1/2^+$	16	1956	IT=100
$^{109}\text{Cd}^n$	-88041.2	1.5	463.10 0.11	$11/2^-$	16	1964	IT=100
$^{109}\text{In}$	-86490	4	4.159 h 0.010	$9/2^+*$	16	1948	$\beta^+=100$ *
$^{109}\text{In}^m$	-85840	4	649.79 0.10	$1/2^-$	16	1966	IT=100 *
$^{109}\text{In}^n$	-84388	4	2101.86 0.11	$19/2^+$	16	1963	IT=100[gs=100,m=0] *
$^{109}\text{Sn}$	-82630	8	18.1 m 0.2	$5/2^+*$	16	1966	$\beta^+=100$ *
$^{109}\text{Sb}$	-76251	5	17.2 s 0.5	$5/2^+\#$	16	1976	$\beta^+=100$
$^{109}\text{Te}$	-67715	4	4.4 s 0.2	$(5/2^+)$	16	1967	$\beta^+=96.1\ 13;\alpha=3.9\ 13;$ $\beta^+p=9.4\ 31;\beta^+\alpha<0.0049$ *
$^{109}\text{I}$	-57673	7	92.8 $\mu\text{s}$ 0.8	$(1/2^+,3/2^+)$	16	1984	$p=99.986\ 4;\alpha=0.014\ 4$ *
$^{109}\text{Xe}$	-46170	300	13 ms 2	$(7/2^+)$	16	2006	$\alpha\approx 100;\beta^+ ?;\beta^+p ?$ *
* $^{109}\text{Nb}$	T : average 15Lo04=110(6) 11Ni01=100(+9-8); other 09Pe06=130(20) **						
* $^{109}\text{Nb}$	D : % $\beta^-$ n other 09Pe06<15 conflicting **						
* $^{109}\text{Nb}^m$	T : symmetrized from 12Ka36=114(+8-7); other 11Wa03=150(30) **						
* $^{109}\text{Nb}^n$	J : from 11Wa03, based on conf=p7/2[413],K=7/2+ and oblate shape **						
* $^{109}\text{Mo}$	T : others 15Lo04=700(+40-60), 92Ay02=530(60) **						
* $^{109}\text{Mo}$	J : 20Ur02=1/2+, 1/2+[411] **						
* $^{109}\text{Mo}^m$	T : symmetrized from 12Ka36=194(+76-49) **						
* $^{109}\text{Mo}^n$	J : 20Ur02=5/2+, 5/2+[413] **						
* $^{109}\text{Tc}$	T : average 19Do02=870(70) 09Pe06=1040(110) 96Me09=820(100) 92PeZX=870(40) **						
* $^{109}\text{Tc}$	T : 69WiZX=930(30) 90Al43=900(100) **						
* $^{109}\text{Ag}$	J : 50Cr26,37Ja01=1/2 **						
* $^{109}\text{Cd}$	J : also 18Yo07,13Yo02=5/2 **						
* $^{109}\text{In}$	J : 58Ma43,59Ma19=9/2 **						
* $^{109}\text{In}^m$	E : other 20Ho03=651(27) **						
* $^{109}\text{In}^n$	E : other 20Ho03=2098(11) **						
* $^{109}\text{Sn}$	J : 87Eb01=5/2, but in conflict with 74Ho17=7/2 **						
* $^{109}\text{I}$	T : other (not used) 19Xi06=89.3(6.0) **						
$^{110}\text{Zr}$	-42220#	500#		$37.5\ \text{ms}\ 2.0$	$0^+$	12 15Lo04 T	1997 $\beta^-=100;\beta^-n ?;\beta^-2n ?$
$^{110}\text{Nb}$	-52310	840	*	75 ms 1	$5^+\#$	12 20Ha14 TD	1994 $\beta^-=100;\beta^-n=40\ 8;\beta^-2n ?$
$^{110}\text{Nb}^m$	-52210#	840#	100# 50#	94 ms 9	$2^+\#$	12 20Ha14 TD	2020 $\beta^-=100;IT ?;\beta^-n=40\ 8;$ $\beta^-2n ?$ *
$^{110}\text{Mo}$	-64536	24		292 ms 7	$0^+$	12 15Lo04 T	1992 $\beta^-=100;\beta^-n=2.0\ 7$
$^{110}\text{Tc}$	-71035	9		900 ms 13	$(2^+,3^+)$	12	1976 $\beta^-=100;\beta^-n=0.04\ 2$
$^{110}\text{Ru}$	-80073	9		12.04 s 0.17	$0^+$	12	1970 $\beta^-=100$
$^{110}\text{Rh}$	-82829	18	*	3.35 s 0.12	$(1^+)$	12	1963 $\beta^-=100$
$^{110}\text{Rh}^m$	-82610#	150#	220# 150#	28.5 s 1.3	$(6^+)$	12	1969 $\beta^-=100$
$^{110}\text{Pd}$	-88330.9	0.6		STABLE >290Ey	$0^+$	12 16Le16 T	1935 IS=11.72 9;2 $\beta^- ?$ *
$^{110}\text{Ag}$	-87457.3	1.3		24.56 s 0.11	$1^+*$	12	1937 $\beta^-\approx 100;\varepsilon=0.30\ 6$
$^{110}\text{Ag}^m$	-87456.2	1.3	1.112 0.016	660 ns 40	$2^-$	12	1975 IT=100
$^{110}\text{Ag}^n$	-87339.7	1.3	117.59 0.05	249.863 d 0.024	$6^+*$	12 FGK209 T	1938 $\beta^-=98.67\ 8;IT=1.33\ 8$
$^{110}\text{Cd}$	-90348.0	0.4		STABLE	$0^+$	12	1925 IS=12.470 61
$^{110}\text{In}$	-86470	12		4.92 h 0.08	$7^+*$	12	1939 $\beta^+=100$
$^{110}\text{In}^m$	-86408	12	62.08 0.04	69.1 m 0.5	$2^+*$	12	1962 $\beta^+=100$
$^{110}\text{Sn}$	-85842	14		4.154 h 0.004	$0^+$	12	1965 $\varepsilon=100$
$^{110}\text{Sb}$	-77450	6		23.6 s 0.3	$(3^+)$	12	1972 $\beta^+=100$
$^{110}\text{Te}$	-72230	7		18.6 s 0.8	$0^+$	12	1977 $\beta^+\approx 100;\alpha ?$
$^{110}\text{I}$	-60470	60		664 ms 24	$(1^+)$	12	1977 $\beta^+=83\ 4;\alpha=17\ 4;\beta^+p=11\ 3;$ $\beta^+\alpha=1.1\ 3$
$^{110}\text{Xe}$	-51920	100		93 ms 3	$0^+$	12	1981 $\alpha=64\ 35;\beta^+=36\ 35;\beta^+p ?$
* $^{110}\text{Nb}$	T : 20Ha14 $\beta - \gamma(t)$ gated on gamma's depopulating 5+ and 6+ levels; **						
* $^{110}\text{Nb}$	T : others 15Lo04=82(2) 11Ni01=86(6) 11Wa26=81(6), 75(9) **						
* $^{110}\text{Nb}$	T : 96Me09=170(20) both for the gs and isomer **						
* $^{110}\text{Nb}$	D : % $\beta^-$ n from 96Me09 includes both gs and isomer **						
* $^{110}\text{Nb}^m$	T : 20Ha14 beta-gamma time gated on gamma depopulating 2+ following **						
* $^{110}\text{Nb}^n$	T : 110Zr->110Nb->110Mo decay; only low spin levels are populated **						
* $^{110}\text{Nb}^m$	D : % $\beta^-$ n from 96Me09 includes both gs and isomer **						
* $^{110}\text{Pd}$	T : 16Le16 supersedes 13Le10 **						

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)
$^{111}\text{Zr}$	-36480#	600#		24.0 ms	0.5		15 15Lo04	T 2010	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$
$^{111}\text{Nb}$	-48960#	300#		54 ms	2		15	1997	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$
$^{111}\text{Mo}$	-59940	13		193.6 ms	4.4		15 15Lo04	T 1994	$\beta^- = 100; \beta^- n < 12$
$^{111}\text{Mo}^m$	-59840#	50#	100# 50#	~ 200 ms			15	2011	$\beta^- = 100; \beta^- n ?$
$^{111}\text{Tc}$	-69025	11		350 ms	11		09 09Pe06	T 1988	$\beta^- = 100; \beta^- n = 0.85$
$^{111}\text{Ru}$	-76785	10		2.12 s	0.07		09	1971	$\beta^- = 100$
$^{111}\text{Rh}$	-82304	7		11 s	1		09	1975	$\beta^- = 100$
$^{111}\text{Pd}$	-85985.9	0.7		23.56 m	0.09		09 15Kr07	T 1937	$\beta^- = 100$
$^{111}\text{Pd}^m$	-85813.7	0.7	172.18 0.08	5.563 h	0.013		09 15Kr07	TD 1952	IT=76.8 10; $\beta^- = 23.2$
$^{111}\text{Ag}$	-88215.4	1.5		7.433 d	0.010		09 16Co01	T 1937	$\beta^- = 100$
$^{111}\text{Ag}^m$	-88155.6	1.5	59.82 0.04	64.8 s	0.8		09	1957	IT=99.3 2; $\beta^- = 0.7$
$^{111}\text{Cd}$	-89252.2	0.4		STABLE			09	1925	IS=12.795 12
$^{111}\text{Cd}^m$	-88856.0	0.4	396.214 0.021	48.50 m	0.09		09	1945	IT=100
$^{111}\text{In}$	-88392	3		2.8048 d	0.0001		09 FGK209	T 1947	$\epsilon = 100$
$^{111}\text{In}^m$	-87855	3	536.99 0.07	7.7 m	0.2		09	1966	IT=100
$^{111}\text{Sn}$	-85939	5		35.3 m	0.6		09	1949	$\beta^+ = 100$
$^{111}\text{Sn}^m$	-85684	5	254.71 0.04	12.5 $\mu\text{s}$	1.0		09	1972	IT=100
$^{111}\text{Sb}$	-80837	9		75 s	1		09	1972	$\beta^+ = 100$
$^{111}\text{Te}$	-73587	6		26.2 s	0.6		09 05Sh24	T 1967	$\beta^+ = 100; \beta^+ p ?$
$^{111}\text{I}$	-64954	5		2.5 s	0.2		09	1977	$\beta^+ \approx 100; \alpha \approx 0.088$
$^{111}\text{Xe}$	-54520#	120#		740 ms	200		09 12Ca03	D 1979	$\beta^+ = 89.6$ 1.9; $\alpha = 10.4$ 1.9; $\beta^+ p ?$
$^{111}\text{Cs}$	-42900#	200#		1# $\mu\text{s}$					p ?
* $^{111}\text{Mo}$	T : average 15Lo04=196(5) 11Ku16=186(9); other 09Pe06=200(+41-36)								
* $^{111}\text{Tc}$	T : other 96Me09=290(20), supersedes 88Pe13=300(30)								
* $^{111}\text{Pd}$	T : average 15Kr07=23.6(0.1) 77Kr14=23.4(0.2)								
* $^{111}\text{Ag}$	T : average 16Co01=7.423(0.013) 74Ro18=7.450(0.017)								
* $^{111}\text{Cd}$	J : also 13Yo02=1/2								
* $^{111}\text{Cd}^m$	J : also 13Yo02=11/2								
* $^{111}\text{Te}$	T : others (not used) 67Ka01=19.0(7) 67Bo41=19.5(5), outliers								
* $^{111}\text{I}$	D : % $\alpha$ from 78Ro19								
$^{112}\text{Zr}$	-32420#	700#		43 ms	21		15 15Lo04	T 2010	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$
$^{112}\text{Nb}$	-44070#	300#		38 ms	2		15 15Lo04	T 1997	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$
$^{112}\text{Mo}$	-57480#	200#		125 ms	5		15 15Lo04	T 1994	$\beta^- = 100; \beta^- n ?$
$^{112}\text{Tc}$	-65259	6		323 ms	6		15 15Lo04	T 1990	$\beta^- = 100; \beta^- n = 1.5$
$^{112}\text{Tc}^m$	-64907	6	352.3 0.7	150 ns	17		15 10Br15	T 2010	IT=100
$^{112}\text{Ru}$	-75631	10		1.75 s	0.07		15	1970	$\beta^- = 100$
$^{112}\text{Rh}$	-79730	40		3.4 s	0.4		15 99Lh01	T 1972	$\beta^- = 100$
$^{112}\text{Rh}^m$	-79390	60	340 70 BD	6.73 s	0.15		15 99Lh01	T 1987	$\beta^- = 100$
$^{112}\text{Pd}$	-86321	7		21.04 h	0.17		15	1951	$\beta^- = 100$
$^{112}\text{Ag}$	-86583.7	2.4		3.130 h	0.008		15	1938	$\beta^- = 100$
$^{112}\text{Cd}$	-90574.86	0.25		STABLE			15	1925	IS=24.109 7
$^{112}\text{In}$	-87990	4		14.88 m	0.15		15	1947	$\beta^+ = 62$ 4; $\beta^- = 38$ 4
$^{112}\text{In}^m$	-87833	4	156.592 0.025	20.67 m	0.08		15	1953	IT=100
$^{112}\text{In}^n$	-87639	4	350.80 0.05	690 ns	50		15	1976	IT=100
$^{112}\text{In}^p$	-87376	4	613.82 0.06	2.81 $\mu\text{s}$	0.03		15	1976	IT=100
$^{112}\text{Sn}$	-88655.05	0.29		STABLE			15	1927	IS=0.97 1; $\beta^+ ?$
$^{112}\text{Sb}$	-81599	18		53.5 s	0.6		15	1959	$\beta^+ = 100$
$^{112}\text{Sb}^m$	-80773	18	825.9 0.4	536 ns	22		15	1976	IT=100
$^{112}\text{Te}$	-77568	8		2.0 m	0.2		15	1976	$\beta^+ = 100$
$^{112}\text{I}$	-67063	10		3.34 s	0.08		15 78Ro19	D 1977	$\beta^+ \approx 100; \alpha \approx 0.0012$ ;
$^{112}\text{Xe}$	-60026	8		2.7 s	0.8		15	1978	$\beta^+ = 98.8$ 8; $\alpha = 1.2$ 8; $\beta^+ p ?$
$^{112}\text{Cs}$	-46420#	120#		490 $\mu\text{s}$	30		15	1994	p $\approx 100; \alpha < 0.26$
* $^{112}\text{Zr}$	T : symmetrized from 15Lo04=30(+30-10)								
* $^{112}\text{Nb}$	T : other 11Ni01=33(+9-6) same group								
* $^{112}\text{Tc}$	D : % $\beta^- n$ from 99Wa09=1.5(0.2), supersedes 96Me09=2.6(0.5); other 09Pe06=4(1)								
* $^{112}\text{Tc}$	T : others 09Pe06=290(11); 99Wa09=290(20), supersedes 96Me09=230(20)								
* $^{112}\text{Tc}^m$	E : 12Ka36=93.1(0.5) keV and 259.2(0.5) keV gamma rays in cascade to gs								

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>112</sup> Tc <sup>m</sup>	T : from 10Br15=150(17); other 12Ka36=218(+60-43)						**
* <sup>112</sup> Rh	T : 99Lh01=3.45(0.37) supersedes 91Jo11=2.1(0.3), 88Ay02=3.8(0.6) same group						**
* <sup>112</sup> Rh <sup>m</sup>	T : supersedes 88Ay02=6.8(0.2) of the same group						**
* <sup>112</sup> Xe	D : % $\alpha$ symmetrized from 94Pa11=0.8(+1.1-0.5); other 78Ro19~0.84						**
<sup>113</sup> Zr	-26340# 300#		15# ms >550 ns	3/2 <sup>+</sup>	18Sh11 IT	2018	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
<sup>113</sup> Nb	-40210# 400#		32 ms 4	3/2 <sup>-</sup> #	15	1997	$\beta^-$ =100; $\beta^-n$ ?; $\beta^-2n$ ?
<sup>113</sup> Mo	-52650# 300#		80 ms 2	5/2 <sup>+</sup> #	15	1994	$\beta^-$ =100; $\beta^-n$ ?
<sup>113</sup> Tc	-62812 3		152 ms 8	5/2 <sup>+</sup> #	15	1992	$\beta^-$ =100; $\beta^-n=2.1$ 3
<sup>113</sup> Tc <sup>m</sup>	-62698 3	114.4 0.5	527 ns 16	5/2 <sup>-</sup> #	15 12Ka36 T	2010	IT=100 *
<sup>113</sup> Ru	-71870 40		800 ms 50	(1/2 <sup>+</sup> )	10	1988	$\beta^-$ =100
<sup>113</sup> Ru <sup>m</sup>	-71740 50	131 33	510 ms 30	(7/2 <sup>-</sup> )	10 98Ku17 E	1998	$\beta^-$ = ?; IT= ? *
<sup>113</sup> Rh	-78767 7		2.80 s 0.12	(7/2 <sup>+</sup> )	10 93Pe11 J	1971	$\beta^-$ =100
<sup>113</sup> Pd	-83590 7		93 s 5	(5/2 <sup>+</sup> )	10	1954	$\beta^-$ =100
<sup>113</sup> Pd <sup>m</sup>	-83509 7	81.1 0.3	300 ms 100	(9/2 <sup>-</sup> )	10	1993	IT=100
<sup>113</sup> Ag	-87027 17		5.37 h 0.05	1/2 <sup>-</sup> *	10	1949	$\beta^-$ =100
<sup>113</sup> Ag <sup>m</sup>	-86984 17	43.50 0.10	68.7 s 1.6	7/2 <sup>+</sup>	10	1958	IT=64 7; $\beta^-$ =36 7
<sup>113</sup> Cd	-89043.29 0.24		8.04 Py 0.05	1/2 <sup>+</sup> *	10	1925	IS=12.227 7; $\beta^-$ =100 *
<sup>113</sup> Cd <sup>m</sup>	-88779.75 0.24	263.54 0.03	13.89 y 0.11	11/2 <sup>-</sup> *	10 11Ko01 TD	1965	$\beta^-$ =99.9036 19; IT=0.0964 19 *
<sup>113</sup> In	-89367.12 0.19		STABLE	9/2 <sup>+</sup> *	10	1934	IS=4.281 52
<sup>113</sup> In <sup>m</sup>	-88975.42 0.19	391.699 0.003	1.6579 h 0.0004	1/2 <sup>-</sup> *	10	1939	IT=100 *
<sup>113</sup> Sn	-88328.1 1.6		115.08 d 0.04	1/2 <sup>+</sup> *	10 FGK209 T	1939	$\beta^+$ =100
<sup>113</sup> Sn <sup>m</sup>	-88250.7 1.6	77.389 0.019	21.4 m 0.4	7/2 <sup>+</sup> *	10	1961	IT=91.1 23; $\beta^+$ =8.9 23
<sup>113</sup> Sb	-84417 17		6.67 m 0.07	5/2 <sup>+</sup>	10	1958	$\beta^+$ =100
<sup>113</sup> Te	-78347 28		1.7 m 0.2	(7/2 <sup>+</sup> )	10	1974	$\beta^+$ =100
<sup>113</sup> I	-71120 8		6.6 s 0.2	5/2 <sup>+</sup> #	10	1977	$\beta^+$ =100; $\alpha=3.310e-5$ #; $\beta^+ \alpha$ ?
<sup>113</sup> Xe	-62204 7		2.74 s 0.08	5/2 <sup>+</sup> #	10 85Ti02 D	1973	$\beta^+ \approx 100$ ; $\alpha=?$ ; $\beta^+p=7$ 4; $\beta^+ \alpha \approx 0.007$ 4 *
<sup>113</sup> Xe <sup>m</sup>	-61800 7	403.6 1.4	6.9 $\mu$ s 0.3	(11/2 <sup>-</sup> )	13Pr01 ETJ	2013	IT=100
<sup>113</sup> Cs	-51765 9		16.94 $\mu$ s 0.09	(3/2 <sup>+</sup> )	15 16Ho16 T	1984	p=100 *
<sup>113</sup> Ba	-39710# 300#		30# ms	5/2 <sup>+</sup> #			p ?; $\alpha$ ?
* <sup>113</sup> Tc <sup>m</sup>	T : symmetrized from 12Ka36=526(+16-15); other 10Br15=500(100)						**
* <sup>113</sup> Ru <sup>m</sup>	E : above the 98-keV level and below 164-keV level						**
* <sup>113</sup> Cd	T : from 07Be61=8.037(0.005,stat)(0.05,sys);						**
* <sup>113</sup> Cd	T : other (recent) 09Da03=8.00(0.11)(syt 0.24) outweighed						**
* <sup>113</sup> Cd	J : also 13Yo02=1/2						**
* <sup>113</sup> Cd <sup>m</sup>	T : average 11Ko01=13.97(0.13) 72Wa11=14.6(0.5) 65Fl02=13.6(0.2)						**
* <sup>113</sup> Cd <sup>m</sup>	J : also 13Yo02=11/2						**
* <sup>113</sup> In <sup>m</sup>	T : from Ensdf12010=99.476(0.023) m						**
* <sup>113</sup> Xe	D : % $\alpha=0.0024-0.0204$ from estimated limit for the reduced width in 85Ti02;						**
* <sup>113</sup> Xe	D : % $\beta^+p$ and % $\beta^+ \alpha$ derived from $\beta^+p/\alpha=605(35)$ and						**
* <sup>113</sup> Xe	D : $\beta^+p/\beta^+ \alpha=500-1500$ in 85Ti02						**
* <sup>113</sup> Cs	T : average 16Ho16=16.9(0.1) (>10000 events) 15Wa02=17.1(0.2) (18000 events)						**
<sup>114</sup> Nb	-34960# 500#		17 ms 5	2 <sup>-</sup> #	15	2010	$\beta^-$ =100; $\beta^-n$ ?; $\beta^-2n$ ?
<sup>114</sup> Mo	-49680# 300#		58 ms 2	0 <sup>+</sup>	15	1997	$\beta^-$ =100; $\beta^-n$ ?
<sup>114</sup> Tc	-58600 430		121 ms 9	5 <sup>+</sup> #	12 15Lo04 T	1994	$\beta^-$ =100; $\beta^-n=1.3$ 4 *
<sup>114</sup> Tc <sup>m</sup>	-58437 13	160 430 MD*	90 ms 20	1 <sup>+</sup> #	12 11Ri01 TD	2011	$\beta^- \approx 100$ ; IT ?; $\beta^-n=1.3$ 4 *
<sup>114</sup> Ru	-70221 4		540 ms 30	0 <sup>+</sup>	12 06Mo07 T	1991	$\beta^-$ =100; $\beta^-n$ ?; $\beta^-2n$ ? *
<sup>114</sup> Rh	-75710 70		1.85 s 0.05	1 <sup>+</sup>	12	1988	$\beta^-$ =100
<sup>114</sup> Rh <sup>m</sup>	-75510# 170#	200# 150# *	1.85 s 0.05	(7 <sup>-</sup> )	12	1987	$\beta^-$ =100
<sup>114</sup> Pd	-83490 7		2.42 m 0.06	0 <sup>+</sup>	12	1958	$\beta^-$ =100
<sup>114</sup> Ag	-84931 5		4.6 s 0.1	1 <sup>+</sup>	12	1958	$\beta^-$ =100
<sup>114</sup> Ag <sup>m</sup>	-84732 5	198.9 1.0	1.50 ms 0.05	(6 <sup>+</sup> )	12 90Pe10 TED	1990	IT=100 *
<sup>114</sup> Cd	-90014.93 0.28		STABLE >92Py	0 <sup>+</sup>	12 95Ge14 T	1925	IS=28.754 81; $2\beta^-$ ?
<sup>114</sup> In	-88569.8 0.3		71.9 s 0.1	1 <sup>+</sup>	12	1937	$\beta^-$ =99.50 15; $\beta^+=0.50$ 15
<sup>114</sup> In <sup>m</sup>	-88379.5 0.3	190.2682 0.0008	49.51 d 0.01	5 <sup>+</sup> *	12	1939	IT=96.75 24; $\beta^+=3.25$ 24
<sup>114</sup> In <sup>m</sup>	-88067.9 0.3	501.948 0.003	43.1 ms 0.6	8 <sup>-</sup>	12	1958	IT=100[gs=0,m=100]
<sup>114</sup> Sn	-90559.735 0.029		STABLE	0 <sup>+</sup>	12	1927	IS=0.66 1
<sup>114</sup> Sn <sup>m</sup>	-87472.37 0.08	3087.37 0.07	733 ns 14	7 <sup>-</sup>	12	1980	IT=100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{114}\text{Sb}$	-84497	20	3.49 m 0.03	$3^+$	12	1959	$\beta^+=100$	
$^{114}\text{Sb}^m$	-84002	20	219 $\mu\text{s}$ 12	$(8^-)$	12	1973	IT=100	
$^{114}\text{Te}$	-81890	24	15.2 m 0.7	$0^+$	12	1968	$\beta^+=100$	
$^{114}\text{I}$	-72639	20	2.01 s 0.15	$1^+$	12 20Ay05	TD 1977	$\beta^+=100; \beta^+p ?; \alpha \approx 7.7e-9\#$	
$^{114}\text{I}^m$	-72373	20	6.2 s 0.5	$(7^-)$	12	1995	$\beta^+ = ?; IT = ?$	
$^{114}\text{Xe}$	-67086	11	10.0 s 0.4	$0^+$	12	1977	$\beta^+=100$	
$^{114}\text{Cs}$	-54690	90	570 ms 20	$(1^+)$	12	1978	$\beta^+ \approx 100; \alpha = 0.018\ 6;$ $\beta^+p = 8.7\ 13; \beta^+ \alpha = 0.19\ 3$	
$^{114}\text{Ba}$	-45910	100	460 ms 125	$0^+$	12 16Ca33	T 1995	$\beta^+ \approx 100; \beta^+p = 20\ 10; \alpha = 0.9\ 3;$ 12C < 0.0034	
* $^{114}\text{Tc}$	T : average 15Lo04=120(10) 11Ro01=110(20) 99Wa09=150(30)							**
* $^{114}\text{Tc}$	T : other: 06Mo07=91(+62-35) probably mixture of gs and isomer							**
* $^{114}\text{Tc}$	D : % $\beta^-n$ from 99Wa09, value in a mixture of gs and isomer							**
* $^{114}\text{Tc}^m$	D : % $\beta^-n$ from 99Wa09, value in a mixture of gs and isomer							**
* $^{114}\text{Ru}$	T : average 06Mo07=510(+69-65) 92Jo05=530(60) 91Le09=570(50)							**
* $^{114}\text{Ag}^m$	E : 34.5(0.5), 43.9(0.5), 47.4(0.5), 73.1(0.5) in a cascade to gs in 90Pe10							**
* $^{114}\text{I}$	T : average 20Ay05=1.89(0.23) 77Ki11=2.1(0.2)							**
* $^{114}\text{I}^m$	J : from M3 to (4-) following by E2 to (2)- following by E1 to 1+							**
* $^{114}\text{Ba}$	T : average 16Ca33=380(+190-110) 97Ja12=430(+300-150)							**
$^{115}\text{Nb}$	-30880#	500#	23 ms 8	$3/2^- \#$	15	2010	$\beta^- = 100; \beta^-n ?; \beta^-2n ?$	
$^{115}\text{Mo}$	-44550#	400#	45.5 ms 2.0	$3/2^+ \#$	15	2010	$\beta^- = 100; \beta^-n ?; \beta^-2n ?$	
$^{115}\text{Tc}$	-55800#	200#	78 ms 2	$5/2^+ \#$	15	1994	$\beta^- = 100; \beta^-n ?$	
$^{115}\text{Ru}$	-66105	25	318 ms 19	$(1/2^+)$	12	1992	$\beta^- = 100; \beta^-n ?$	
$^{115}\text{Ru}^m$	-66110	90	82	6	76 ms 6	12 10Ku25	ETJ 2010	
$^{115}\text{Rh}$	-74229	7	1.03 s 3	$(7/2^+)$	12 92PeZX	T 1988	$\beta^- = 100; \beta^-n ?$	
$^{115}\text{Pd}$	-80426	14	25 s 2	$(1/2^+)$	12	1958	$\beta^- = 100$	
$^{115}\text{Pd}^m$	-80337	14	89.21	0.16	50 s 3	12	1987	$\beta^- = 92.0\ 20; IT = 8.0\ 20$
$^{115}\text{Ag}$	-84983	18	20.0 m 0.5	$1/2^-$	12	1949	$\beta^- = 100$	
$^{115}\text{Ag}^m$	-84942	18	41.16	0.10	18.0 s 0.7	12	1958	$\beta^- = 79.0\ 3; IT = 21.0\ 3$
$^{115}\text{Cd}$	-88084.5	0.7	53.46 h 0.05	$1/2^+ *$	12	1939	$\beta^- = 100$	
$^{115}\text{Cd}^m$	-87903.5	0.9	181.0	0.5	44.56 d 0.24	12	1959	$\beta^- \approx 100; IT ?$
$^{115}\text{In}$	-89536.357	0.012	441 Ty 25		441 Ty 25	12	1924	IS=95.719 52; $\beta^- = 100$
$^{115}\text{In}^m$	-89200.113	0.021	336.244	0.017	4.486 h 0.004	12	1961	IT=95.0 7; $\beta^- = 5.0\ 7$
$^{115}\text{Sn}$	-90033.846	0.015	STABLE		$1/2^+ *$	12	1927	IS=0.34 1
$^{115}\text{Sn}^m$	-89421.04	0.04	612.81	0.04	3.26 $\mu\text{s}$ 0.08	12	1967	IT=100
$^{115}\text{Sn}^n$	-89320.21	0.12	713.64	0.12	159 $\mu\text{s}$ 1	12	1958	IT=100
$^{115}\text{Sb}$	-87003	16	32.1 m 0.3	$5/2^+$	12	1958	$\beta^+ = 100$	
$^{115}\text{Sb}^m$	-84207	16	2796.26	0.09	159 ns 3	12	1977	IT=100
$^{115}\text{Te}$	-82063	28	*		5.8 m 0.2	12	1961	$\beta^+ = 100$
$^{115}\text{Te}^m$	-82053	30	10	6	6.7 m 0.4	12 74Ch51	E 1974	$\beta^+ \approx 100; IT ?$
$^{115}\text{Te}^n$	-81783	28	280.05	0.20	7.5 $\mu\text{s}$ 0.2	12	1972	IT=100
$^{115}\text{I}$	-76338	29	1.3 m 0.2	$5/2^+ \#$	12	1969	$\beta^+ = 100$	
$^{115}\text{Xe}$	-68657	12	18 s 3	$(5/2^+)$	12	1969	$\beta^+ = 100; \beta^+p = 0.34\ 6$	
$^{115}\text{Cs}$	-59700#	100#	1.4 s 0.8	$9/2^+ \#$	12	1978	$\beta^+ = 100; \beta^+p \approx 0.07$	
$^{115}\text{Ba}$	-48920#	200#	450 ms 50	$5/2^+ \#$	12 97Ja12	D 1997	$\beta^+ = 100; \beta^+p > 15$	
* $^{115}\text{Ru}^m$	E : 20 keV above the 61.7-keV level in 10Ku25							**
* $^{115}\text{Rh}$	T : average 92PeZX=1.04(0.03) 88Ay01=0.99(0.05)							**
* $^{115}\text{Cd}$	J : also 13Yo02=1/2							**
* $^{115}\text{Cd}^m$	J : also 13Yo02=11/2							**
* $^{115}\text{Te}^m$	E : less than 20 keV in 74Ch51							**
* $^{115}\text{Xe}$	T : average 71Ho07=18(4) 69Ha03=19(5)							**
$^{116}\text{Nb}$	-25230#	300#	12# ms > 550ns	$1^- \#$	18Sh11	I 2018	$\beta^- ?; \beta^-n ?; \beta^-2n ?$	
$^{116}\text{Mo}$	-41210#	500#	32 ms 4	$0^+$	15	2010	$\beta^- = 100; \beta^-n ?; \beta^-2n ?$	
$^{116}\text{Tc}$	-51210#	300#	57 ms 3	$2^+ \#$	15	1997	$\beta^- = 100; \beta^-n ?; \beta^-2n ?$	
$^{116}\text{Ru}$	-64069	4	204 ms 6	$0^+$	15	1994	$\beta^- = 100; \beta^-n ?$	
$^{116}\text{Rh}$	-70740	70	685 ms 39	$1^+$	10 06Mo07	TD 1970	$\beta^- = 100; \beta^-n < 2.1$	
$^{116}\text{Rh}^m$	-70540#	170#	200#	150#	570 ms 50	10 01Wa04	T 1987	$\beta^- = 100; \beta^-n < 2.1$
$^{116}\text{Pd}$	-79831	7	11.8 s 0.4	$0^+$	10	1970	$\beta^- = 100$	
$^{116}\text{Ag}$	-82543	3	3.83 m 0.08	$(0^-)$	10 09Ba52	TJ 1958	$\beta^- = 100$	

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{116}\text{Ag}^m$	-82495	3	47.90 0.10	20 s 1	(3 <sup>+</sup> )	10 05Ba94 TJD 2005	$\beta^- = 93$ 4;IT=7 4
$^{116}\text{Ag}^n$	-82413	3	129.80 0.22	9.3 s 0.3	(6 <sup>-</sup> )	10 05Ba94 TJD 1970	$\beta^- = 92$ 4;IT=8 4
$^{116}\text{Cd}$	-88712.49	0.16		26.9 Ey 0.9	0 <sup>+</sup>	10 20Ba.A T 1925	IS=7.512 54;2 $\beta^- = 100$
$^{116}\text{In}$	-88249.76	0.22		14.10 s 0.03	1 <sup>+</sup>	10 13Wr01 D 1937	$\beta^- \approx 100$ ; $\epsilon = 0.0237$ 43
$^{116}\text{In}^m$	-88122.49	0.22	127.267 0.006	54.29 m 0.17	5 <sup>+</sup> *	10 1945	$\beta^- = 100$
$^{116}\text{In}^n$	-87960.10	0.22	289.660 0.006	2.18 s 0.04	8 <sup>-</sup> *	10 1950	IT=100[gs=0,m=100]
$^{116}\text{Sn}$	-91525.98	0.10		STABLE	0 <sup>+</sup>	10 1922	IS=14.54 9
$^{116}\text{Sn}^m$	-89160.00	0.10	2365.975 0.021	348 ns 19	5 <sup>-</sup>	10 1964	IT=100
$^{116}\text{Sn}^n$	-87978.82	0.20	3547.16 0.17	833 ns 30	10 <sup>+</sup>	10 1978	IT=100
$^{116}\text{Sb}$	-86822	5		15.8 m 0.8	3 <sup>+</sup> *	10 1949	$\beta^+ = 100$
$^{116}\text{Sb}^m$	-86728	5	93.99 0.05	194 ns 4	1 <sup>+</sup>	10 1976	IT=100
$^{116}\text{Sb}^n$	-86440	40	390 40 BD	60.3 m 0.6	8 <sup>-</sup> *	10 1949	$\beta^+ = 100$
$^{116}\text{Te}$	-85264	24		2.49 h 0.04	0 <sup>+</sup>	10 1958	$\beta^+ = 100$
$^{116}\text{I}$	-77420	80		2.91 s 0.15	1 <sup>+</sup>	10 1976	$\beta^+ = 100$
$^{116}\text{I}^m$	-76990	80	430.4 0.5	3.27 $\mu$ s 0.16	(7 <sup>-</sup> )	10 1990	IT=100
$^{116}\text{Xe}$	-73047	13		59 s 2	0 <sup>+</sup>	10 1969	$\beta^+ = 100$
$^{116}\text{Cs}$	-62040#	100#	*	700 ms 40	(1 <sup>+</sup> )	10 77Bo28 D 1975	$\beta^+ = 100$ ; $\beta^+ p = 0.28$ 7; $\beta^+ \alpha = 0.049$ 25
$^{116}\text{Cs}^m$	-61940#	120#	100# 60# *	3.85 s 0.13	(7 <sup>+</sup> )	10 1975	$\beta^+ = 100$ ; $\beta^+ p = 0.44$ 7; $\beta^+ \alpha = 0.0034$ 23
$^{116}\text{Ba}$	-54380#	200#		1.3 s 0.2	0 <sup>+</sup>	10 1997	$\beta^+ = 100$ ; $\beta^+ p = 3$ 1
$^{116}\text{La}$	-40050#	320#		10# ms		10	$\beta^+ ?$ ; $\beta^+ p ?$ ; $p ?$
* $^{116}\text{Rh}$	T : average 06Mo07=688(+52-50) 88Ay02=680(60)						
* $^{116}\text{Rh}$	D : % $\beta^- n$ from 06Mo07, a mixture of gs and isomer						
* $^{116}\text{Rh}^m$	D : % $\beta^- n$ from 06Mo07, a mixture of gs and isomer						
* $^{116}\text{Ag}$	T : from 09Ba52=230(5) s						
* $^{116}\text{Cd}$	T : value for 2 $\nu$ - $\beta\beta$ ; others (recent) 18Ba44=26.3(0.1,stat)(+1.1-1.2,syst)						
* $^{116}\text{Cd}$	T : 17Ar01=27.4(0.4,stat)(1.8,syst) 15Ba11=28.7(1.3) (evaluation)						
* $^{116}\text{In}$	D : % $\epsilon$ average 13Wr01=0.0246(44stat)(39syst) 98Bh04=0.0227(0.0063)						
* $^{116}\text{In}$	T : also 13Wr01=14.9(0.8)						
* $^{116}\text{Cs}$	D : % $\beta^+ p$ from 77Bo28; Ensdf2010=2.8(0.7)% in error						
* $^{116}\text{Cs}^m$	D : % $\beta^+ p$ average 77Bo28=0.66(0.13) 78Da07=0.36(0.08)%; Birge ratio=1.97						
* $^{116}\text{Cs}^m$	D : % $\beta^+ \alpha$ average 78Da07=0.008(0.002)% and 0.0022(0.0010), from						
* $^{116}\text{Cs}^m$	D : % $\beta^+ p = 0.44(0.07)$ and $\beta^+ p/\beta^+ \alpha = 200(80)$ in 85Ti02;						
* $^{116}\text{Cs}^m$	D : Birge ratio=2.6						
* $^{116}\text{Cs}^m$	J : direct $\beta^+$ feedings to 6+ and 8+ levels in $^{116}\text{Xe}$ in 80Ma16						
* $^{116}\text{La}$	T : estimate for $\beta^+$ decay; no p decay within 20 us-20 ms						
$^{117}\text{Mo}$	-35690#	500#		22 ms 5	3/2 <sup>+</sup> #	15 2010	$\beta^- = 100$ ; $\beta^- n ?$ ; $\beta^- 2n ?$
$^{117}\text{Tc}$	-48140#	400#		44.5 ms 3.0	5/2 <sup>+</sup> #	15 1997	$\beta^- = 100$ ; $\beta^- n ?$ ; $\beta^- 2n ?$
$^{117}\text{Ru}$	-59490	430		151 ms 3	3/2 <sup>+</sup> #	15 1994	$\beta^- = 100$ ; $\beta^- n ?$
$^{117}\text{Ru}^m$	-59310	430	185.0 0.4	2.49 $\mu$ s 0.06	7/2 <sup>-</sup> #	15 2012	IT=100
$^{117}\text{Rh}$	-68897	9		421 ms 30	7/2 <sup>+</sup> #	11 06Mo07 TD 1991	$\beta^- = 100$ ; $\beta^- n < 7.6$
$^{117}\text{Rh}^m$	-68576	9	321.2 1.0	138 ns 17	3/2 <sup>+</sup> #	13La25 ET 2013	IT=100
$^{117}\text{Pd}$	-76424	7		4.3 s 0.3	(3/2 <sup>+</sup> )	11 04Ur04 J 1968	$\beta^- = 100$
$^{117}\text{Pd}^m$	-76221	7	203.3 0.3	19.1 ms 0.7	(9/2 <sup>-</sup> )	11 04Ur04 J 1990	IT=100
$^{117}\text{Ag}$	-82182	14		73.6 s 1.4	1/2 <sup>-</sup> #	11 1958	$\beta^- = 100$
$^{117}\text{Ag}^m$	-82153	14	28.6 0.2	5.34 s 0.05	7/2 <sup>+</sup> #	11 1990	$\beta^- = 94.0$ 15;IT=6.0 15
$^{117}\text{Cd}$	-86418.4	1.0		2.503 h 0.005	1/2 <sup>+</sup> *	11 19Gi09 T 1939	$\beta^- = 100$
$^{117}\text{Cd}^m$	-86282.0	1.0	136.4 0.2	3.441 h 0.009	11/2 <sup>-</sup> *	11 19Gi09 T 1966	$\beta^- = 100$
$^{117}\text{In}$	-88943	5		43.2 m 0.3	9/2 <sup>+</sup> *	11 1937	$\beta^- = 100$
$^{117}\text{In}^m$	-88628	5	315.303 0.011	116.2 m 0.3	1/2 <sup>-</sup> *	11 1940	$\beta^- = 52.9$ 15;IT=47.1 15
$^{117}\text{Sn}$	-90397.7	0.5		STABLE	1/2 <sup>+</sup> *	11 20Yo.A J 1923	IS=7.68 7
$^{117}\text{Sn}^m$	-90083.1	0.5	314.58 0.04	13.939 d 0.024	11/2 <sup>-</sup> *	12 20Yo.A J 1950	IT=100
$^{117}\text{Sn}^n$	-87991.3	0.6	2406.4 0.4	1.75 $\mu$ s 0.07	(19/2 <sup>+</sup> )	11 1979	IT=100
$^{117}\text{Sb}$	-88640	8		2.97 h 0.02	5/2 <sup>+</sup> *	11 21Da02 T 1947	$\beta^+ = 100$
$^{117}\text{Sb}^m$	-85509	8	3130.76 0.19	355 $\mu$ s 17	(25/2 <sup>+</sup> )	11 1970	IT=100
$^{117}\text{Sb}^n$	-85409	8	3230.7 0.2	290 ns 5	(23/2 <sup>-</sup> )	11 1987	IT=100
$^{117}\text{Te}$	-85096	13		62 m 2	1/2 <sup>+</sup> *	11 1958	$\beta^+ = 100$ ; $\epsilon = 75$ 1; $e^+ = 25$ 1
$^{117}\text{Te}^m$	-84800	13	296.1 0.5	103 ms 3	(11/2 <sup>-</sup> )	11 99Mo30 J 1963	IT=100
$^{117}\text{I}$	-80439	26		2.22 m 0.04	(5/2 <sup>+</sup> )	11 1969	$\beta^+ = 100$ ; $e^+ \approx 77$
$^{117}\text{Xe}$	-74185	10		61 s 2	5/2 <sup>+</sup> *	11 90NeZY J 1969	$\beta^+ = 100$ ; $\beta^+ p = 0.0029$ 6

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{117}\text{Cs}$	-66490	60			8.4 s 0.6	$9/2^+\#$	11	1972	$\beta^+=100$
$^{117}\text{Cs}^m$	-66340#	100#	150#	80#	6.5 s 0.4	$3/2^+\#$	11	1978	$\beta^+=100$
$^{117}\text{Cs}^x$	-66440	80	50	50	$R=?$	$spmix$			
$^{117}\text{Ba}$	-57460	250			1.75 s 0.07	$(3/2^+)$	11 97Ja12	D 1977	$\beta^+=100;\beta^+p=13.3$ ; $\beta^+\alpha=0.0248$
$^{117}\text{La}$	-46270#	200#			21.7 ms 1.8	$(3/2^+)$	11 11Li28	TJ 2001	$p\approx 100;\beta^+ ?;\beta^+p ?$
$^{117}\text{La}^m$		<i>non-exist</i>		RN	10 ms 5	$(9/2^+)$	11 01So02	I	
$^{117}\text{Ru}^m$		T: symmetrized from 12Ka36=2.487(+0.058-0.055); other 12LaZT=2.0(0.3)							**
$^{117}\text{Rh}$		T: average 06Mo07=394(+47-43) 91Pe10=440(40)							**
$^{117}\text{Ag}$		T: symmetrized from 72.8(+2.0-0.7)							**
$^{117}\text{Ag}^m$		J: E3 to 1/2-#							**
$^{117}\text{Cd}$		J: 13Yo02=1/2							**
$^{117}\text{Cd}^m$		J: 13Yo02=1/2							**
$^{117}\text{Sn}^m$		T: average 16Do10=13.91(0.03) 14Un01=14.00 (0.05) 03Po21=13.98 (0.07)							**
$^{117}\text{Xe}$		J: 90NeZY=5/2							**
$^{117}\text{Ba}$		D: % $\beta^+p$ from 97Ja12. $\beta^+p/\beta^+\alpha=350-1200$ from 85Ti02 yields							**
$^{117}\text{Ba}$		D: % $\beta^+\alpha=0.011\%-0.037\%$							**
$^{117}\text{La}$		T: average 11Li28=20.1(2.5) 01Ma69=24(3) 01So02=22(5)							**
$^{117}\text{La}^m$		I: reported in 01So02 with E=121(10) keV, but not confirmed in 11Li28							**
$^{118}\text{Mo}$	-32370#	500#			21 ms 6	$0^+$	15 15Lo04	TD 2015	$\beta^-=100;\beta^-n ?;\beta^-2n ?$
$^{118}\text{Tc}$	-43290#	400#			30 ms 4	$2^+\#$	15	2010	$\beta^-=100;\beta^-n ?;\beta^-2n ?$
$^{118}\text{Ru}$	-57000#	200#			99 ms 3	$0^+$	15	1994	$\beta^-=100;\beta^-n ?$
$^{118}\text{Rh}$	-64887	24			282 ms 9	$1^+\#$	06 15Lo04	T 1994	$\beta^-=100;\beta^-n=3.114$
$^{118}\text{Rh}^m$	-64690#	150#	200#	150#	310 ms 30	$6^-\#$	06 00Jo18	T 1994	$\beta^-\approx 100;IT ?;\beta^-n=3.114$
$^{118}\text{Pd}$	-75388.4	2.5			1.9 s 0.1	$0^+$	06	1969	$\beta^-=100$
$^{118}\text{Ag}$	-79553.8	2.5			3.76 s 0.15	$(2^-)$	95 93Ja03	J 1967	$\beta^+=100$
$^{118}\text{Ag}^m$	-79508.0	2.5	45.79	0.09	$\sim 0.1 \mu\text{s}$	$(1,2)^-$	95 93Ja03	J 1989	IT=100
$^{118}\text{Ag}^n$	-79426.2	2.5	127.63	0.10	2.0 s 0.2	$(5^+)$	95 FGK208	JD 1971	$\beta^-=59.3;IT=41.3$
$^{118}\text{Ag}^p$	-79274.4	2.5	279.37	0.20	$\sim 0.1 \mu\text{s}$	$(3^+)$	95 93Ja03	TJ 1989	IT=100
$^{118}\text{Cd}$	-86702	20			50.3 m 0.2	$0^+$	95	1961	$\beta^-=100$
$^{118}\text{In}$	-87228	8			5.0 s 0.5	$1^+$	95	1949	$\beta^-=100$
$^{118}\text{In}^m$	-87130#	50#	100#	50#	4.364 m 0.007	$5^+*$	95 94It.A	T 1964	$\beta^+=100$
$^{118}\text{In}^n$	-86990#	50#	240#	50#	8.5 s 0.3	$8^-*$	95	1969	IT=98.6 3[gs=0,m=98.6]; $\beta^-=1.43$
$^{118}\text{Sn}$	-91652.8	0.5			STABLE	$0^+$	95	1924	IS=24.22 9
$^{118}\text{Sn}^m$	-89077.9	0.5	2574.91	0.04	230 ns 10	$7^-$	95	1961	IT=100
$^{118}\text{Sn}^n$	-88544.7	0.5	3108.06	0.22	2.52 $\mu\text{s}$ 0.06	$(10^+)$	95 11Fo15	J 1973	IT=100
$^{118}\text{Sb}$	-87996	3			3.6 m 0.1	$1^+*$	95	1947	$\beta^+=100$
$^{118}\text{Sb}^m$	-87945	3	50.814	0.021	20.6 $\mu\text{s}$ 0.6	$3^+$	95	1975	IT=100
$^{118}\text{Sb}^n$	-87746	5	250	6	5.01 h 0.03	$8^-*$	95 21Da02	T 1947	$\beta^+=100$
$^{118}\text{Te}$	-87691	18			6.00 d 0.02	$0^+$	95	1948	$\epsilon=100$
$^{118}\text{I}$	-80971	20			13.7 m 0.5	$(2^-)$	95	1957	$\beta^+=100$
$^{118}\text{I}^m$	-80782	20	188.8	0.7	8.5 m 0.5	$(7^-)$	95 03Mo36	E 1968	$\beta^+\approx 100;IT ?$
$^{118}\text{Xe}$	-78079	10			3.8 m 0.9	$0^+$	95	1965	$\beta^+=100$
$^{118}\text{Cs}$	-68409	13			14 s 2	$2^+*$	95	1969	$\beta^+=100;\beta^+p=0.02114$ ; $\beta^+\alpha=0.00125$
$^{118}\text{Cs}^m$	-68310#	60#	100#	60#	17 s 3	$(7^-)$	95 93Be46	J 1972	$\beta^+=100;\beta^+p=0.02114$ ; $\beta^+\alpha=0.00125$
$^{118}\text{Cs}^x$	-68404	12	5	4	$R < 0.1$	$spmix$			
$^{118}\text{Ba}$	-62200#	200#			5.2 s 0.2	$0^+$	06 97Ja12	T 1997	$\beta^+=100$
$^{118}\text{La}$	-49620#	300#			200# ms	$1^-\#$			$\beta^+ ?;\beta^+p ?$
$^{118}\text{Mo}$		T: symmetrized from 15Lo04=19(+7-4)							**
$^{118}\text{Rh}$		T: average 15Lo04=285(10) 06Mo07=266(+22-21) from $\beta^-(t)$ ; probably contain							**
$^{118}\text{Rh}$		T: contributions from both the low- and high-spin $\beta^-$ decaying states							**
$^{118}\text{Rh}$		J: direct $\beta^-$ feeding to $0^+$ state in $^{118}\text{Pd}$ in 06Wa10							**
$^{118}\text{Rh}$		D: % $\beta^-n$ from 06Mo07, probably a mixture of gs and isomer							**
$^{118}\text{Rh}^m$		T: from $\beta^- - \gamma(t)$ using 575-keV gamma ray, depopulating the $4^+$							**
$^{118}\text{Rh}^m$		T: level in $^{118}\text{Pd}$ , in 00Jo18; most-likely dominated by the							**
$^{118}\text{Rh}^m$		T: high-spin $\beta^-$ decaying state							**
$^{118}\text{Rh}^m$		J: direct $\beta^-$ feeding to 6- level in $^{118}\text{Pd}$ in 06Wa10							**
$^{118}\text{Rh}^m$		D: % $\beta^-n$ from 06Mo07, probably a mixture of gs and isomer							**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
* <sup>118</sup> Ag <sup>n</sup>	J : 127.6-keV gamma-ray E3 to (2-)						**	
* <sup>118</sup> Ag <sup>n</sup>	D : from Ig(127keV, <sup>118</sup> Ag)/Ig(48 keV, <sup>118</sup> Cd)=0.124(0.015) in 73FoZF						**	
* <sup>118</sup> In <sup>n</sup>	E : 138.2(0.5) keV above <sup>118</sup> In <sup>m</sup>						**	
* <sup>118</sup> Sb <sup>m</sup>	J : E2 to 1+						**	
* <sup>118</sup> Sb <sup>n</sup>	T : average 21Da02=5.18(0.05) 74Ca06=5.00(0.02) 72Pa13=5.11(0.06)						**	
* <sup>118</sup> Sb <sup>n</sup>	T : 68Ki06=5.15(0.05) 67Ha27=4.96(0.02); Birge ratio=2.68						**	
* <sup>118</sup> Im	E : from a least-squares fit to level scheme of 03Mo36						**	
* <sup>118</sup> Cs	D : from %β <sup>+</sup> p=0.042(6), %β <sup>+</sup> α=0.0024(4) for mixture of gs and isomer						**	
<sup>119</sup> Mo	-26580#	300#	12# ms >550ns	3/2 <sup>+</sup> #	18Sh11 I	2018	β <sup>-</sup> ?; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?	
<sup>119</sup> Tc	-40170#	500#	22 ms 3	5/2 <sup>+</sup> #	15	2010	β <sup>-</sup> =100; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?	
<sup>119</sup> Ru	-52080#	300#	69.5 ms 2.0	3/2 <sup>+</sup> #	15	1997	β <sup>-</sup> =100; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?	
<sup>119</sup> Ru <sup>m</sup>	-51850#	300#	227.1 0.7	384 ns 22	15	2012	IT=100	
<sup>119</sup> Rh	-62823	9	190 ms 6	7/2 <sup>+</sup> #	09 15Lo04 T	1994	β <sup>-</sup> =100; β <sup>-</sup> n=6.4 16	
<sup>119</sup> Pd	-71407	8	920 ms 80	3/2 <sup>+</sup> #	09 06Mo07 TD	1991	β <sup>-</sup> =100; β <sup>-</sup> n ?	
<sup>119</sup> Pd <sup>m</sup>	-71110#	150#	300# 150#	3# ms			IT ?; β <sup>-</sup> ?	
<sup>119</sup> Ag	-78646	15	6.0 s 0.5	1/2 <sup>-</sup> #	09	1975	β <sup>-</sup> =100	
<sup>119</sup> Ag <sup>m</sup>	-78626#	25#	20# 20#	2.1 s 0.1	7/2 <sup>+</sup> #	09	1975	β <sup>-</sup> =100
<sup>119</sup> Cd	-83980	40	2.69 m 0.02	1/2 <sup>+</sup> *	09 13Yo02 J	1961	β <sup>-</sup> =100	
<sup>119</sup> Cd <sup>m</sup>	-83830	40	146.54 0.11	2.20 m 0.02	11/2 <sup>-</sup> *	09 13Yo02 J	1974	β <sup>-</sup> =100
<sup>119</sup> In	-87699	7	2.4 m 0.1	9/2 <sup>+</sup> *	09	1949	β <sup>-</sup> =100	
<sup>119</sup> In <sup>m</sup>	-87388	7	311.37 0.03	18.0 m 0.3	1/2 <sup>-</sup> *	09 76Sc30 D	1973	β <sup>-</sup> =97.4 4; IT=2.6 4
<sup>119</sup> In <sup>n</sup>	-87045	7	654.27 0.07	130 ns 15	(3/2) <sup>+</sup>	09	1974	IT=100
<sup>119</sup> In <sup>p</sup>	-85042	7	2656.9 1.8	265 ns 10	(25/2 <sup>+</sup> )	09 20Bi06 T	2002	IT=100
<sup>119</sup> Sn	-90065.0	0.7	STABLE	1/2 <sup>+</sup> *	09	1925	IS=8.59 4	
<sup>119</sup> Sn <sup>m</sup>	-89975.5	0.7	89.531 0.013	293.1 d 0.7	11/2 <sup>-</sup> *	09 20Yo.A J	1950	IT=100
<sup>119</sup> Sn <sup>n</sup>	-87938.0	1.2	2127.0 1.0	9.6 μs 1.2	(19/2 <sup>+</sup> )	09	1992	IT=100
<sup>119</sup> Sn <sup>p</sup>	-87696.0	0.8	2369.0 0.3	96 ns 9	23/2 <sup>+</sup>	16Is03 ETJ	2016	IT=100
<sup>119</sup> Sb	-89476	7	38.19 h 0.22	5/2 <sup>+</sup> *	09	1947	ε=100	
<sup>119</sup> Sb <sup>m</sup>	-86922	7	2553.6 0.3	130 ns 3	19/2 <sup>-</sup>	09 91Io02 J	1991	IT=100
<sup>119</sup> Sb <sup>n</sup>	-86634	7	2841.7 0.4	835 ms 81	25/2 <sup>+</sup>	09 19Mi18 ET	1979	IT=100
<sup>119</sup> Te	-87183	7	16.05 h 0.05	1/2 <sup>+</sup> *	09	1948	β <sup>+</sup> =100; ε=97.94 5; e <sup>+</sup> =2.06 5	
<sup>119</sup> Te <sup>m</sup>	-86922	7	260.96 0.05	4.70 d 0.04	11/2 <sup>-</sup> *	09	1960	β <sup>+</sup> =100; ε=99.59 4; e <sup>+</sup> =0.41 4
<sup>119</sup> I	-83778	22	19.1 m 0.4	5/2 <sup>+</sup>	09	1954	β <sup>+</sup> =100; e <sup>+</sup> =51 4; ε=49 4	
<sup>119</sup> Xe	-78794	10	5.8 m 0.3	5/2 <sup>+</sup> *	09 90NeZY J	1965	β <sup>+</sup> =100; e <sup>+</sup> =79 5; ε=21 5	
<sup>119</sup> Cs	-72305	14	43.0 s 0.2	9/2 <sup>+</sup> *	09 75Ho09 D	1969	β <sup>+</sup> =100; β <sup>+</sup> α < 2e-6	
<sup>119</sup> Cs <sup>m</sup>	-72260#	30#	50# 30#	30.4 s 0.1	3/2 <sup>+</sup> *	09	1978	β <sup>+</sup> =100
<sup>119</sup> Cs <sup>s</sup>	-72289	9	16 11	R = 0.50 0.25	spmix			
<sup>119</sup> Ba	-64590	200	5.4 s 0.3	(5/2 <sup>+</sup> )	09	1974	β <sup>+</sup> =100; β <sup>+</sup> p=25 2	
<sup>119</sup> La	-55020#	300#	1# s	11/2 <sup>-</sup> #			β <sup>+</sup> ?	
<sup>119</sup> Ce	-43820#	500#	200# ms	5/2 <sup>+</sup> #			β <sup>+</sup> ?; β <sup>+</sup> p ?	
* <sup>119</sup> Ru <sup>m</sup>	T : symmetrized from 12Ka36=383(+22-21)						**	
* <sup>119</sup> Pd	T : average 06Mo07=918(111) 91Pe04=920(130)						**	
* <sup>119</sup> Ag <sup>m</sup>	E : estimated from 7/2+ levels in <sup>113</sup> Ag=43 keV <sup>115</sup> Ag=41 keV and						**	
* <sup>119</sup> Ag <sup>m</sup>	E : <sup>117</sup> Ag=28 keV						**	
* <sup>119</sup> In <sup>m</sup>	D : %IT symmetrized from 76Sc30=2.5(+0.5-0.3); other 61Gi06~5						**	
* <sup>119</sup> In <sup>p</sup>	T : average 20Bi06=270(11) 02Lu15=240(25)						**	
* <sup>119</sup> Sb <sup>n</sup>	T : average 19Mi18=776(181) 79Sh03=850(90)						**	
* <sup>119</sup> Sb <sup>n</sup>	E : based on 19Mi18=2799(30) keV and known states in <sup>119</sup> Sn see 87Lu06;						**	
* <sup>119</sup> Sb <sup>n</sup>	E : other Ensdf2009=x keV above 2841.7(0.4)-keV level, conflicting						**	
* <sup>119</sup> Sb <sup>n</sup>	J : from 87Lu06						**	
* <sup>119</sup> Xe	J : 90NeZY=5/2						**	
<sup>120</sup> Tc	-35000#	500#	21 ms 5	3 <sup>+</sup> #	17	2010	β <sup>-</sup> =100; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?	
<sup>120</sup> Ru	-49720#	400#	45 ms 2	0 <sup>+</sup>	17	2010	β <sup>-</sup> =100; β <sup>-</sup> n ?	
<sup>120</sup> Rh	-58620#	200#	129.6 ms 4.2	8 <sup>-</sup> #	17 15Lo04 T	1994	β <sup>-</sup> =100; β <sup>-</sup> n < 9.3; β <sup>-</sup> 2n ?	
<sup>120</sup> Rh <sup>m</sup>	-58460#	200#	157.2 0.7	295 ns 16	6#	17 12Ka36 ETD2012	IT=100	
<sup>120</sup> Pd	-70279.6	2.3	492 ms 33	0 <sup>+</sup>	17 06Mo07 TD	1993	β <sup>-</sup> =100; β <sup>-</sup> n < 0.7	
<sup>120</sup> Ag	-75652	4	1.52 s 0.07	4(+)	02 12Ba58 TJ	1971	β <sup>-</sup> =100; β <sup>-</sup> n < 0.003	





**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{121}\text{Te}$	-88543	26	19.31 d 0.07	$1/2^+$	10 19Jo03 T	1939	$\beta^+=100$ *
$^{121}\text{Te}^m$	-88249	26	293.974 0.022	$11/2^-$	10 19Jo03 T	1940	IT=88.6 11; $\beta^+=11.4$ 11 *
$^{121}\text{I}$	-86246	5	2.12 h 0.01	$5/2^+*$	10	1950	$\beta^+=100$ *
$^{121}\text{I}^m$	-83869	5	2376.9 0.4	$21/2^+\#$	10	1982	IT=100 *
$^{121}\text{Xe}$	-82481	10	40.1 m 2.0	$5/2^+*$	10 90NeZY J	1952	$\beta^+=100$ *
$^{121}\text{Cs}$	-77102	14	155 s 4	$3/2^+*$	10	1969	$\beta^+=100$ *
$^{121}\text{Cs}^m$	-77034	14	68.5 0.3	$9/2^+*$	10 91Ge02 D	1981	$\beta^+\approx 83$ ;IT $\approx 17$ *
$^{121}\text{Cs}^x$	-77056	16	46 8	$R = 2.1$			<i>spmix</i>
$^{121}\text{Ba}$	-70740	140	29.7 s 1.5	$5/2^+*$	10 75Bo11 D	1975	$\beta^+=100$ ; $\beta^+p=0.02$ 1 *
$^{121}\text{La}$	-62190#	300#	5.3 s 0.2	$11/2^- \#$	10	1988	$\beta^+=100$ ; $\beta^+p?$ *
$^{121}\text{Ce}$	-52690#	400#	1.1 s 0.1	$5/2^+(\#)$	10	1997	$\beta^+=100$ ; $\beta^+p\approx 1$ *
$^{121}\text{Pr}$	-41550#	500#	12 ms 5	$(3/2)^+(\#)$	10	2005	$p\approx 100$ *
* $^{121}\text{Rh}$	T : average 15Lo04=76(5) 20Sh.A=71(7) **						
* $^{121}\text{Rh}$	D : % $\beta^-n$ from 20Sh.A>11.5(0.6) **						
* $^{121}\text{Pd}$	T : other 06Mo07=285(24) **						
* $^{121}\text{Pd}^m$	T : symmetrized from 12Ka36=460(+85-92) assuming two isomers in a cascade; **						
* $^{121}\text{Pd}^m$	T : other 12LaZT=630(50) assuming a single-decaying isomer **						
* $^{121}\text{Pd}^n$	T : symmetrized from 12Ka36=463(+83-94) assuming two isomers in a cascade; **						
* $^{121}\text{Pd}^n$	E : x keV above $^{121}\text{Pd}^m$ , with x<50 keV estimated by nubase **						
* $^{121}\text{Ag}$	T : average 06Mo07=661(+75-72) 83Re05=780(10); others 82Fo10=720(100) **						
* $^{121}\text{Ag}$	T : 95Fe12=1043(80) **						
* $^{121}\text{In}^n$	T : others (not used) 10Re01=17(2)us 02Lu15=350(50)ns **						
* $^{121}\text{Sn}^n$	E : from a least-squares fit to the level scheme in 16Is03 **						
* $^{121}\text{Sn}^p$	E : from a least-squares fit to the level scheme in 16Is03 **						
* $^{121}\text{Sn}^q$	E : from a least-squares fit to the level scheme in 16Is03; other **						
* $^{121}\text{Sn}^q$	E : 2832.7(0.5) from a least-squares fit to the level scheme in 12As05 **						
* $^{121}\text{Sb}^m$	E : x keV above the 2721.6(0.4) level with x<60 in 08Jo03 **						
* $^{121}\text{Te}$	T : average 19Jo03=19.38(0.03) 08Ea01=19.2 (0.1) 95Si30=19.16(0.05); **						
* $^{121}\text{Te}$	T : Birge ratio=2.8 **						
* $^{121}\text{Te}^m$	T : average 19Jo03=165.1(0.7) 08Ea01=164.2 (0.8) **						
* $^{121}\text{Xe}$	J : 90NeZY=5/2 **						
* $^{121}\text{Ba}$	J : 88We14=5/2 **						
* $^{121}\text{Pr}$	T : symmetrized from 05Ro19=10(+6-3) **						
$^{122}\text{Tc}$	-26310#	300#	13# ms >550ns	$1^+\#$	18Sh11 I	2018	$\beta^- ?$ ; $\beta^-n ?$ ; $\beta^-2n ?$
$^{122}\text{Ru}$	-41780#	500#	25 ms 1	$0^+$	15	2010	$\beta^-=100$ ; $\beta^-n ?$ ; $\beta^-2n ?$
$^{122}\text{Rh}$	-51880#	300#	51 ms 6	$7^- \#$	13 15Lo04 T	1997	$\beta^+=100$ ; $\beta^-n < 3.9$ ; $\beta^-2n ?$ *
$^{122}\text{Rh}^m$	-51610#	300#	271.0 0.7	$4^+\#$	13 12Ka36 ETD2012		IT=100 *
$^{122}\text{Pd}$	-64616	20	193 ms 5	$0^+$	14 15Lo04 T	1994	$\beta^-=100$ ; $\beta^-n < 2.5$ *
$^{122}\text{Ag}$	-71110	40	529 ms 13	$(3^+)$	07	1978	$\beta^-=100$ ; $\beta^-n = 0.186$ 10 *
$^{122}\text{Ag}^m$	-71030#	60#	80# 50#	$(1^-)$	07	2000	$\beta^-=100$ ;IT ?; $\beta^-n=?$ *
$^{122}\text{Ag}^n$	-71030#	60#	80# 50#	$(9^-)$	07 95Za01 D	2000	$\beta^+=100$ ;IT ?; $\beta^-n ?$ *
$^{122}\text{Ag}^p$	-70940#	60#	171# 50#	$(1^+)$	13La11 TJE	2013	IT=100 *
$^{122}\text{Cd}$	-80612.4	2.3	5.24 s 0.03	$0^+$	07	1973	$\beta^-=100$ *
$^{122}\text{In}$	-83570	50	1.5 s 0.3	$1^+$	07	1963	$\beta^-=100$ *
$^{122}\text{In}^m$	-83530#	80#	40# 60#	$10.3$ s 0.6	07	1979	$\beta^-=100$ *
$^{122}\text{In}^n$	-83280	130	290 140 BD	10.8 s 0.4	8-*	07	1979 $\beta^-=100$ *
$^{122}\text{Sn}$	-89940.0	2.4	STABLE	$0^+$	07	1928	IS=4.63 3; $2\beta^- ?$ *
$^{122}\text{Sn}^m$	-87531.0	2.4	2409.03 0.04	7.5 $\mu$ s 0.9	07	1979	IT=100 *
$^{122}\text{Sn}^n$	-87174.5	2.4	2765.5 0.3	62 $\mu$ s 3	10 $^+$	07 14Is04 EJ	1992 IT=100 *
$^{122}\text{Sn}^p$	-85218.8	2.4	4721.2 0.3	139 ns 9	15-	14Is04 EJT	2012 IT=100 *
$^{122}\text{Sb}$	-88334.2	2.5	2.7238 d 0.0002	$2^-*$	07	1939	$\beta^-=97.59$ 12; $\beta^+=2.41$ 12 *
$^{122}\text{Sb}^m$	-88272.8	2.5	61.4131 0.0005	1.86 $\mu$ s 0.08	3 $^+$	07	1962 IT=100 *
$^{122}\text{Sb}^n$	-88196.7	2.5	137.4726 0.0008	530 $\mu$ s 30	5 $^+$	07	1963 IT=100 *
$^{122}\text{Sb}^p$	-88170.6	2.5	163.5591 0.0017	4.191 m 0.003	8-	07	1947 IT=100 *
$^{122}\text{Te}$	-90313.3	1.4	STABLE	$0^+$	07	1932	IS=2.55 12 *
$^{122}\text{I}$	-86079	5	3.63 m 0.06	$1^+$	07 12At01 D	1950	$\beta^+=100$ ; $e^+=78$ 2; $e=22$ 2 *
$^{122}\text{I}^m$	-85764	5	314.9 0.4	193.3 ns 0.9	7-	07 19Mo28 TJ	2004 IT=100 *
$^{122}\text{I}^n$	-85700	5	379.4 0.5	79.1 $\mu$ s 1.2	7-	07 19Mo28 TJ	2004 IT=100 *
$^{122}\text{I}^p$	-85685	5	394.1 0.5	78.2 $\mu$ s 0.4	$(8^+)$	07 19Mo28 TJ	2004 IT=100 *
$^{122}\text{I}^q$	-85635	5	444.1 0.5	146.5 ns 1.2	8-	07 19Mo28 TJ	2004 IT=100 *
$^{122}\text{Xe}$	-85355	11	20.1 h 0.1	$0^+$	07	1952	$\epsilon=100$ *

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{122}\text{Cs}$	-78140	30		21.18 s	0.19	1+*	07 75Ho09 D	1969 $\beta^+=100; \beta^+ \alpha < 2e-7$
$^{122}\text{Cs}^m$	-78090	30	45.87 0.12	> 1 $\mu\text{s}$		3+	07	1987 IT=100
$^{122}\text{Cs}^n$	-78005	9	140 30 MD	3.70 m	0.11	8-*	07	1969 $\beta^+=100$
$^{122}\text{Cs}^p$	-78010	30	127.07 0.16	360 ms	20	5-	07	1969 IT=100
$^{122}\text{Cs}^x$	-78130	30	14 7	$R = 0.10$		0.05		<i>spmix</i>
$^{122}\text{Ba}$	-74609	28		1.95 m	0.15	0+	07	1974 $\beta^+=100$
$^{122}\text{La}$	-64540#	300#		8.6 s	0.5	(1-)	07	1984 $\beta^+=100; \beta^+ p=?$
$^{122}\text{Ce}$	-57870#	400#		2# s		0+	07	2005 $\beta^+ ?; \beta^+ p ?$
$^{122}\text{Pr}$	-44780#	500#		500# ms				$\beta^+ ?; \beta^+ p ?$
* $^{122}\text{Rh}$	D : $\% \beta^- n$ from 20Sh.A < 3.87(6)							
* $^{122}\text{Rh}^m$	E : 12Ka36=63.9(0.5) and 207.1(0.5) gamma rays in a cascade to gs							
* $^{122}\text{Rh}^m$	T : symmetrized from 12Ka36=820(+130-110)							
* $^{122}\text{Pd}$	T : average 15Lo04=195(5) 06Mo07=175(16)							
* $^{122}\text{Ag}$	D : $\% \beta^- n$ from 83Re05, probably includes gs and isomers							
* $^{122}\text{Ag}^m$	D : $\beta^- n$ was observed by 00Kr18, but it was not quantified							
* $^{122}\text{Ag}^n$	J : direct $\beta^-$ decay feeding of 8- level in $^{122}\text{Cd}$ in 95Za01;							
* $^{122}\text{Ag}^n$	J : 00Kr18=9 from hfs							
* $^{122}\text{Ag}^p$	E : 91-keV above $^{122}\text{Ag}^m$ in 13La11							
* $^{122}\text{Sn}$	T : Onu-BB 18No01 > 13 Ty							
* $^{122}\text{Sn}^n$	E : from a least-squares fit to the level scheme of 14Is04; other							
* $^{122}\text{Sn}^n$	E : 2765.3(0.4) from a least-squares fit to the level scheme of 12As05							
* $^{122}\text{Sn}^n$	T : other 17Ki09=60.8(+8.3-7.0)							
* $^{122}\text{Sn}^p$	T : average 14Is04=134(12) 12As05=146(15)							
* $^{122}\text{Sn}^p$	E : from a least-squares fit to the level scheme in 14Is04; other							
* $^{122}\text{Sn}^p$	E : 4720.2(0.45) from a least-squares fit to the level scheme in 12As05							
* $^{122}\text{I}$	T : others 12At01=4.15(+0.30-0.25) for $^{122}\text{I}^{+53}$ and							
* $^{122}\text{I}$	T : 3.13(+0.15-0.13) for $^{122}\text{I}^{+52}$							
* $^{122}\text{La}$	J : significant direct $\beta^+$ feeding to 2+ in $^{122}\text{Ba}$ in 92Mo13							
$^{123}\text{Ru}$	-36550#	500#		19 ms	2	3/2+#	15	2010 $\beta^- = 100; \beta^- n ?; \beta^- 2n ?$
$^{123}\text{Rh}$	-49190#	400#		42 ms	4	7/2+#	15 20Sh.A D	2010 $\beta^- = 100; \beta^- n > 24; \beta^- 2n ?$
$^{123}\text{Pd}$	-60430	790		108 ms	1	3/2+#	15 14SmZZ TD	1994 $\beta^- = 100; \beta^- n = 10.6$
$^{123}\text{Pd}^m$	-60330#	790#	100# 50#	100# ms		11/2-#	15 19Ch24 ID	2019 $\beta^- \approx 100; IT ?$
$^{123}\text{Ag}$	-69570	30		294 ms	5	(7/2+)*	17 06Mo07 TD	1976 $\beta^- = 100; \beta^- n = 0.56.5$
$^{123}\text{Ag}^m$	-69510	30	59.5 0.5	100# ms		(1/2-)*	19Ch24 E	2019 $\beta^- = 100; \beta^- n ?$
$^{123}\text{Ag}^n$	-68120#	30#	1450# 14#	202 ns	20		17 13La11 ETD	2013 IT=100
$^{123}\text{Ag}^p$	-68100	30	1472.8 0.8	393 ns	16	(17/2-)	17 13La11 ET	2009 IT=100
$^{123}\text{Cd}$	-77414.2	2.7		2.10 s	0.02	3/2+*	04 13Yo02 J	1983 $\beta^- = 100$
$^{123}\text{Cd}^m$	-77271	3	143 4	1.82 s	0.03	11/2-*	04 13Yo02 J	1986 $\beta^- = ?; IT ?$
$^{123}\text{In}$	-83429	20		6.17 s	0.05	9/2+*	04	1960 $\beta^- = 100$
$^{123}\text{In}^m$	-83102	20	327.21 0.04	47.4 s	0.4	1/2-*	04	1960 $\beta^- = 100$
$^{123}\text{In}^n$	-81351	20	2078.1 0.6	1.4 $\mu\text{s}$	0.2	(17/2-)	04Sc42 ETJ	2004 IT=100
$^{123}\text{In}^p$	-81326#	24#	2103# 14#	> 100 $\mu\text{s}$		(21/2-)	10 10Re01 EJT	2010 IT=100
$^{123}\text{Sn}$	-87814.7	2.5		129.2 d	0.4	11/2-*	04	1948 $\beta^- = 100$
$^{123}\text{Sn}^m$	-87790.1	2.5	24.6 0.4	40.06 m	0.01	3/2+*	04	1948 $\beta^- = 100$
$^{123}\text{Sn}^n$	-85869.8	2.5	1944.90 0.12	7.4 $\mu\text{s}$	2.6	19/2+	04 16Is03 EJ	1992 IT=100
$^{123}\text{Sn}^p$	-85662.0	2.5	2152.66 0.19	6 $\mu\text{s}$		23/2+	04 16Is03 EJ	1994 IT=100
$^{123}\text{Sn}^q$	-85102.2	2.5	2712.47 0.21	34 $\mu\text{s}$		27/2-	04 16Is03 EJ	1994 IT=100
$^{123}\text{Sb}$	-89222.9	1.4		STABLE		7/2+*	04	1922 IS=42.79.5
$^{123}\text{Sb}^m$	-86985.1	1.4	2237.8 0.3	214 ns	3	19/2-	09Wa02 ETJ	2005 IT=100
$^{123}\text{Sb}^n$	-86609.5	1.5	2613.4 0.4	65 $\mu\text{s}$	1	23/2+	09Wa02 ETJ	2007 IT=100
$^{123}\text{Te}$	-89171.0	1.4		STABLE		> 2Py	04 03Al02 T	1932 IS=0.89.3; $\epsilon = 100$
$^{123}\text{Te}^m$	-88923.5	1.4	247.47 0.04	119.2 d	0.1	11/2-	04	1951 IT=100
$^{123}\text{I}$	-87943	4		13.2232 h	0.0015	5/2+*	04 FGK209 T	1949 $\beta^+ = 100$
$^{123}\text{Xe}$	-85248	10		2.08 h	0.02	1/2+*	04 90NeZY J	1952 $\beta^+ = 100$
$^{123}\text{Xe}^m$	-85063	10	185.18 0.11	5.49 $\mu\text{s}$	0.26	7/2-	04	1981 IT=100
$^{123}\text{Cs}$	-81044	12		5.88 m	0.03	1/2+*	04	1954 $\beta^+ = 100$
$^{123}\text{Cs}^m$	-80888	12	156.27 0.05	1.64 s	0.12	11/2-	04	1972 IT=100
$^{123}\text{Cs}^n$	-80792	13	252 6	114 ns	5	(9/2+)	04 GAu127 E	2000 IT=100
$^{123}\text{Cs}^x$	-81037	13	7 4	$R < 0.1$		<i>spmix</i>		
$^{123}\text{Ba}$	-75655	12		2.7 m	0.4	5/2+*	04	1962 $\beta^+ = 100$
$^{123}\text{Ba}^m$	-75534	12	120.95 0.08	830 ns	60	1/2+#	04	1991 IT=100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{123}\text{La}$	-68650#	200#	17 s 3	11/2 <sup>-</sup> #	04	1978	$\beta^+=100$
$^{123}\text{Ce}$	-60290#	300#	3.8 s 0.2	(5/2) <sup>(+)</sup> #	04	1984	$\beta^+=100;\beta^+p=?$
$^{123}\text{Pr}$	-50230#	400#	800# ms	3/2 <sup>+</sup> #			$\beta^+ ?;\beta^+p ?$
* $^{123}\text{Pd}$	T : from 15Lo04=108(1); others 06Mo07=174(+38-34) 14SmZZ=170(+49-52)						
* $^{123}\text{Ag}$	J : 00Kr18=(7/2), $g_{9/2}$ ; hfs comparison to $^{107,107m}\text{Ag}$						
* $^{123}\text{Ag}$	D : % $\beta^-$ -n average 06Mo07=1.0(0.5) 93Ru01=0.55(0.05) 14TeZY=0.60(0.25);						
* $^{123}\text{Ag}$	D : probably includes gs and isomer						
* $^{123}\text{Ag}$	T : average 06Mo07=272(24) 95Fe12=293(7) 86Ma42=300(20) 83Re05=300(10);						
* $^{123}\text{Ag}$	T : others 89Hu10=350(40) 76Lu02=390(30) 14TeZY=396(15)						
* $^{123}\text{Ag}^m$	J : 00Kr18=(1/2), $p_{1/2}$ ; hfs comparison to $^{107,107m}\text{Ag}$						
* $^{123}\text{Ag}^n$	E : 13La11=1365+x keV above $^{123}\text{Ag}^m$ ; x=50# keV estimated by Nubase						
* $^{123}\text{Ag}^p$	T : average 13La11=393(16) 09St28=396(37); other 05WaZY=330(20)						
* $^{123}\text{In}^n$	E : from a least-squares fit to gamma-ray energies in 04Sc42						
* $^{123}\text{In}^p$	E : from 2078.1+x keV; x=50# keV estimated by Nubase						
* $^{123}\text{Sn}^n$	E : from a least-squares fit to the level scheme in 16Is03						
* $^{123}\text{Sn}^p$	E : from a least-squares fit to the level scheme in 16Is03						
* $^{123}\text{Sn}^q$	E : from a least-squares fit to the level scheme in 16Is03						
* $^{123}\text{Sb}^m$	E : from a least-squares fit to gamma-ray energies in 09Wa02						
* $^{123}\text{Sb}^n$	E : from a least-squares fit to gamma-ray energies in 09Wa02						
* $^{123}\text{Xe}$	J : 90NeZY=1/2						
* $^{123}\text{Cs}^m$	J : E3 to 5/2+ followed by E2 to 1/2+						
* $^{123}\text{Cs}^n$	E : from 231.63 + x; x<40 keV estimated by Nubase						
* $^{123}\text{Ba}$	J : also 88We14=5/2						
$^{124}\text{Ru}$	-33590#	600#	15 ms 3	0 <sup>+</sup>	15	2010	$\beta^-=100;\beta^-n ?;\beta^-2n ?$
$^{124}\text{Rh}$	-44710#	400#	30 ms 2	2 <sup>+</sup> #	15 20Sh.A D	2010	$\beta^-=100;\beta^-n<31;\beta^-2n ?$
$^{124}\text{Pd}$	-58400#	300#	88 ms 15	0 <sup>+</sup>	14 14SmZZ TD	1997	$\beta^-=100;\beta^-n=17.5$
$^{124}\text{Pd}^m$	-57400#	850#	1000# 800#	> 20 $\mu\text{s}$	14 12Ka36 ET	2012	IT $\approx$ 100
$^{124}\text{Ag}$	-66230	250					
$^{124}\text{Ag}^m$	-66180#	260#	50# 50#	*	177.9 ms 2.6	(2 <sup>-</sup> )	15 14Ba18 J 1984 $\beta^-=100;\beta^-n=1.3.9$
$^{124}\text{Ag}^n$	-66070	250	155.6 0.5	*	144 ms 20	9 <sup>-</sup> #	15 14Ba18 TDJ 1995 $\beta^-=100;\beta^-n ?$
$^{124}\text{Ag}^p$	-66000	250	231.1 0.7		140 ns 50	(1 <sup>+</sup> )	15 13La11 TJ 2012 IT=100
$^{124}\text{Cd}$	-76699.4	2.6			1.48 $\mu\text{s}$ 0.15	(1 <sup>-</sup> )	15 13La11 TJ 2012 IT=100
$^{124}\text{In}$	-80870	30		*	1.25 s 0.02	0 <sup>+</sup>	08 1974 $\beta^-=100$
$^{124}\text{In}^m$	-80890	50	-20 60	BD*	3.12 s 0.09	3 <sup>+</sup> *	08 1964 $\beta^-=100$
$^{124}\text{In}^n$	-80890	50			3.67 s 0.03	8 <sup>-</sup> *	08 14Le20 T 1974 $\beta^- \approx 100; IT ?$
$^{124}\text{Sn}$	-88231.5	1.3			STABLE >100Py	0 <sup>+</sup>	08 52Ka41 T 1922 IS=5.79 5;2 $\beta^- ?$
$^{124}\text{Sn}^m$	-86026.9	1.3	2204.620 0.023		270 ns 60	5 <sup>-</sup>	08 FGK127 J 1979 IT=100
$^{124}\text{Sn}^n$	-85906.5	1.3	2324.96 0.04		3.1 $\mu\text{s}$ 0.5	7 <sup>-</sup>	08 14Is04 EJ 1979 IT=100
$^{124}\text{Sn}^p$	-85574.9	1.3	2656.6 0.3		51 $\mu\text{s}$ 3	10 <sup>+</sup>	08 14Is04 EJ 1992 IT=100
$^{124}\text{Sn}^q$	-83679.1	1.3	4552.4 0.3		260 ns 25	15 <sup>-</sup>	14Is04 EJ 2012 IT=100
$^{124}\text{Sb}$	-87619.1	1.4			60.20 d 0.03	3 <sup>-</sup> *	08 1939 $\beta^-=100$
$^{124}\text{Sb}^m$	-87608.2	1.4	10.8627 0.0008		93 s 5	5 <sup>+</sup>	08 1947 IT=75 5; $\beta^-$ =25 5
$^{124}\text{Sb}^n$	-87582.3	1.4	36.8440 0.0014		20.2 m 0.2	(8) <sup>-</sup>	08 1947 IT=100[gs=0,m=100]
$^{124}\text{Sb}^p$	-87578.3	1.4	40.8038 0.0007		3.2 $\mu\text{s}$ 0.3	(3 <sup>+</sup> )	08 FGK208 J 1989 IT=100
$^{124}\text{Te}$	-90524.1	1.4			STABLE	0 <sup>+</sup>	08 1932 IS=4.74 14
$^{124}\text{I}$	-87364.6	2.3			4.1760 d 0.0003	2 <sup>-</sup> *	08 92Wo03 T 1938 $\beta^+=100$
$^{124}\text{Xe}$	-87667.4	1.4			STABLE >200Ty	0 <sup>+</sup>	08 89Ba22 T 1922 IS=0.095 5;2 $\beta^+ ?$
$^{124}\text{Cs}$	-81741	9			30.9 s 0.4	1 <sup>+</sup> *	08 1969 $\beta^+=100$
$^{124}\text{Cs}^m$	-81278	9	462.63 0.14		6.41 s 0.07	(7) <sup>+</sup>	08 17Ra20 D 1983 IT=99.89 2; $\beta^+$ =0.11 2
$^{124}\text{Cs}^n$	-81711	22	30 20		R=?	<i>spmix</i>	
$^{124}\text{Ba}$	-79090	12			11.0 m 0.5	0 <sup>+</sup>	08 1967 $\beta^+=100$
$^{124}\text{La}$	-70260	60		*&	29.21 s 0.17	(7,8 <sup>-</sup> )	08 92Id01 J 1978 $\beta^+=100$
$^{124}\text{La}^m$	-70160#	120#	100# 100#	*&	21 s 4	2 <sup>-</sup> #	08 1992 $\beta^+=100$
$^{124}\text{Ce}$	-64920#	300#			9.1 s 1.2	0 <sup>+</sup>	08 97As05 T 1978 $\beta^+=100$
$^{124}\text{Pr}$	-53150#	400#			1.2 s 0.2		08 1986 $\beta^+=100;\beta^+p=?$
$^{124}\text{Nd}$	-44830#	500#			500# ms	0 <sup>+</sup>	$\beta^+ ?;\beta^+p ?$
* $^{124}\text{Pd}$	T : from 15Lo04; others 06Mo07=38(+38-19) 14SmZZ=0.144(+25-24)						
* $^{124}\text{Ag}$	T : average 15Lo04=180(3) 95Fe12=172(5);others 14Ba18=191(28) 84Hi03=170(30)						
* $^{124}\text{Ag}$	D : % $\beta^-$ -n from 06Mo07, probably includes gs and isomer						
* $^{124}\text{Ag}^m$	J : $\beta^-$ feeding to 8+ and 10+ levels in $^{124}\text{Cd}$ in 14Ba18 would be						
* $^{124}\text{Ag}^n$	J : consistent with J=9; 14Ba18 assumes J=(8-)						
* $^{124}\text{Ag}^p$	E : 12Ka36=75.5(0.5) and 155.6(0.5) gamma rays in a cascade to gs						

Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>124</sup> Ag <sup>p</sup>	T : average 13La11=1.46(0.20) 12Ka36=1.62(+0.29-0.24) 05WaZY=1.3(0.3)						**
* <sup>124</sup> In <sup>m</sup>	T : from 14Le20; others 86Go10=3.7(0.2) 74Fo23=2.4(0.3)						**
* <sup>124</sup> Sn <sup>m</sup>	J : E1 to 4+; L(p,p)=5 for <sup>124</sup> Sn <sup>m</sup>						**
* <sup>124</sup> Sn <sup>n</sup>	T : from 17Ki09=2.83(0.12); other 79Fo10=3.1(0.5)						**
* <sup>124</sup> Sn <sup>p</sup>	T : average 17Ki09=55.0(+4.7-4.1) 92Br06=45(5)						**
* <sup>124</sup> Sn <sup>q</sup>	T : average 14Is04=260(30) 12As05=260(25)						**
* <sup>124</sup> Sb <sup>p</sup>	J : E2 to 5+						**
* <sup>124</sup> I	T : other (similar precision) 16Lu16=4.1758(14) 21Pi01=1.179(0.006)						**
* <sup>124</sup> Xe	T : 2nu-εε 16Ab03(XMASS)>4.7 Zy (at 90% C.L.); 17Ap02(XENON)>0.65 Zy;						**
* <sup>124</sup> Xe	T : 2nu-ε(K)ε(K) 19Ap03=18(5,stat)(1,syst) Zy						**
* <sup>124</sup> Cs <sup>m</sup>	T : from 14Le20; other 83We07=6.3(0.2)						**
* <sup>124</sup> Ce	T : average 97As05=10.8(1.5) 78Bo32=6(2)						**
<sup>125</sup> Ru	-28370#	300#		12# ms >550ns	3/2 <sup>+</sup> #	18Sh11 I 2018	β <sup>-</sup> ?; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>125</sup> Rh	-41830#	500#		26.5 ms 2.0	7/2 <sup>+</sup> #	15 2010	β <sup>-</sup> =100; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>125</sup> Pd	-53960#	400#		60 ms 6	3/2 <sup>+</sup> #	15 14SmZZ TD 2008	β <sup>-</sup> =100; β <sup>-</sup> n=12 4 *
<sup>125</sup> Pd <sup>m</sup>	-53860#	400#	100# 50#	50# ms	11/2 <sup>-</sup> #	19Ch24 ID 2019	β <sup>-</sup> ≈100; IT ?
<sup>125</sup> Pd <sup>n</sup>	-52160#	400#	1805.23 0.18	144 ns 4	(23/2 <sup>+</sup> )	15 19Wa14 ETJ 2019	IT=100
<sup>125</sup> Ag	-64520	430		160 ms 5	(9/2 <sup>+</sup> )*	15 14SmZZ TD 1994	β <sup>-</sup> =100; β <sup>-</sup> n=11.8 10 *
<sup>125</sup> Ag <sup>m</sup>	-64420	430	97.1 0.5	50# ms	(1/2 <sup>-</sup> )*	19Ch24 E 2019	β <sup>-</sup> ?; IT ?; β <sup>-</sup> n ? *
<sup>125</sup> Ag <sup>n</sup>	-63020	430	1501.2 0.6	491 ns 20	(17/2 <sup>-</sup> )	15 2009	IT=100
<sup>125</sup> Cd	-73348.1	2.9		680 ms 40	3/2 <sup>+</sup> *	11 13Yo02 J 1986	β <sup>-</sup> =100
<sup>125</sup> Cd <sup>m</sup>	-73162	3	186 4 MD	480 ms 30	11/2 <sup>-</sup> *	11 13Yo02 J 1986	β <sup>-</sup> =100
<sup>125</sup> Cd <sup>n</sup>	-71700	5	1648 4	19 μs 3	(19/2 <sup>+</sup> )	11Si32 EJT 2011	IT=100 *
<sup>125</sup> In	-80412.3	1.8		2.36 s 0.04	9/2 <sup>+</sup> *	11 1967	β <sup>-</sup> =100
<sup>125</sup> In <sup>m</sup>	-80060	12	352 12	12.2 s 0.2	1/2 <sup>-</sup> *	11 1974	β <sup>-</sup> =100
<sup>125</sup> In <sup>n</sup>	-78402.9	1.9	2009.4 0.7	9.4 μs 0.6	(19/2 <sup>+</sup> )	11 1998	IT=100
<sup>125</sup> In <sup>p</sup>	-78251.1	2.0	2161.2 0.9	5.0 ms 1.5	(23/2 <sup>-</sup> )	11 1998	IT=100
<sup>125</sup> Sn	-85893.7	1.3		9.634 d 0.015	11/2 <sup>-</sup> *	11 20Yo.A J 1939	β <sup>-</sup> =100 *
<sup>125</sup> Sn <sup>m</sup>	-85866.2	1.3	27.50 0.14	9.77 m 0.25	3/2 <sup>+</sup> *	11 20Yo.A J 1939	β <sup>-</sup> =100 *
<sup>125</sup> Sn <sup>n</sup>	-84000.9	1.3	1892.8 0.3	6.2 μs 0.2	19/2 <sup>+</sup>	11 08Lo07 J 2000	IT=100
<sup>125</sup> Sn <sup>p</sup>	-83834.2	1.4	2059.5 0.4	650 ns 60	23/2 <sup>+</sup>	11 16Is03 T 2008	IT=100 *
<sup>125</sup> Sn <sup>q</sup>	-83270.2	1.4	2623.5 0.5	230 ns 17	27/2 <sup>-</sup>	11 08Lo07 T 2000	IT=100 *
<sup>125</sup> Sb	-88255.1	2.5		2.7576 y 0.0011	7/2 <sup>+</sup> *	11 FGK209 T 1951	β <sup>-</sup> =100 *
<sup>125</sup> Sb <sup>m</sup>	-86283.9	2.5	1971.25 0.20	4.1 μs 0.2	15/2 <sup>-</sup>	11 2007	IT=100
<sup>125</sup> Sb <sup>n</sup>	-86143.0	2.5	2112.1 0.3	28.5 μs 0.5	19/2 <sup>-</sup>	11 FGK128 J 2007	IT=100 *
<sup>125</sup> Sb <sup>q</sup>	-85784.1	2.5	2471.0 0.4	277.0 ns 6.4	(23/2 <sup>+</sup> ) <sup>+</sup>	11 19Bi04 T 2007	IT=100 *
<sup>125</sup> Te	-89021.8	1.4		STABLE	1/2 <sup>+</sup> *	11 1931	IS=7.07 15
<sup>125</sup> Te <sup>m</sup>	-88877.0	1.4	144.775 0.008	57.40 d 0.15	11/2 <sup>-</sup>	11 1949	IT=100
<sup>125</sup> I	-88836.0	1.4		59.392 d 0.008	5/2 <sup>+</sup> *	11 FGK209 T 1947	ε=100
<sup>125</sup> Xe	-87199.4	1.4		16.87 h 0.08	1/2 <sup>+</sup> *	11 19S201 T 1950	β <sup>+</sup> =100
<sup>125</sup> Xe <sup>m</sup>	-86946.8	1.4	252.61 0.14	56.9 s 0.9	9/2 <sup>-</sup> *	11 1954	IT=100
<sup>125</sup> Xe <sup>n</sup>	-86903.5	1.4	295.89 0.15	140 ns 30	7/2 <sup>+</sup>	11 1979	IT=100
<sup>125</sup> Cs	-84090	8		44.35 m 0.29	1/2 <sup>+</sup> *	11 19S201 T 1954	β <sup>+</sup> =100
<sup>125</sup> Cs <sup>m</sup>	-83824	8	266.1 1.1	900 μs 30	(11/2 <sup>-</sup> )	11 98Su16 J 1998	IT=100
<sup>125</sup> Ba	-79669	11		3.3 m 0.3	1/2 <sup>+</sup> *	11 1968	β <sup>+</sup> =100
<sup>125</sup> Ba <sup>m</sup>	-79549#	23#	120# 20#	2.76 μs 0.14	(7/2 <sup>-</sup> )	11 FGK128 J 1989	IT=100 *
<sup>125</sup> La	-73759	26		64.8 s 1.2	11/2 <sup>-</sup> #	11 1973	β <sup>+</sup> =100
<sup>125</sup> La <sup>m</sup>	-73652	26	107.00 0.10	390 ms 40	(3/2 <sup>+</sup> )	11 99Ca21 J 1998	IT=100
<sup>125</sup> Ce	-66660#	200#		9.7 s 0.3	(7/2 <sup>-</sup> )	11 02Pe15 J 1978	β <sup>+</sup> =100; β <sup>+</sup> p=?
<sup>125</sup> Ce <sup>m</sup>	-66570#	200#	93.6 0.4	13 s 10	(1/2 <sup>+</sup> )	11 07Su07 ETJ 2007	IT=100 *
<sup>125</sup> Pr	-58070#	300#		3.3 s 0.7	3/2 <sup>+</sup> #	11 2002	β <sup>+</sup> =100; β <sup>+</sup> p ?
<sup>125</sup> Nd	-48070#	400#		650 ms 150	(5/2)( <sup>+</sup> #)	11 1999	β <sup>+</sup> =100; β <sup>+</sup> p>0
* <sup>125</sup> Pd	T : average 15Lo04=57(10) 14SmZZ=61(+8-7)						**
* <sup>125</sup> Ag	J : 00Kr18=(9/2) g <sub>9/2</sub> ; hfs comparison to <sup>107,107m</sup> Ag						**
* <sup>125</sup> Ag	T : average 15Lo04=150(8) 14SmZZ=163(+11-9) 95Fe12=166(7)						**
* <sup>125</sup> Ag <sup>m</sup>	J : 00Kr18=(1/2) p <sub>1/2</sub> ; hfs comparison to <sup>107,107m</sup> Ag						**
* <sup>125</sup> Cd <sup>n</sup>	E : 11Si32=1461.8(0.7) keV above <sup>125</sup> Cd <sup>m</sup>						**
* <sup>125</sup> Sn	T : average 20Gu04=9.63(0.02) 68Er03=9.67(0.04) 66La13=9.625(0.025)						**
* <sup>125</sup> Sn <sup>m</sup>	T : unweighted average 20Gu04=10.01(0.08) 68Er03=9.52(0.05);						**
* <sup>125</sup> Sn <sup>n</sup>	T : Birge ratio=5.19						**
* <sup>125</sup> Sn <sup>p</sup>	J : E2 to 19/2+						**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>125</sup> Sn <sup>g</sup>	J : E2 to 23/2-						**
* <sup>125</sup> Sn <sup>g</sup>	T : average 08Lo07=230(20) 00Zh47=230(30)						**
* <sup>125</sup> Sb	T : evaluated by FGK209=1007.2(0.4) d using the world data; other						**
* <sup>125</sup> Sb	T : Ensdf2011=2.75856(0.00025) based on old data, but the small						**
* <sup>125</sup> Sb	T : uncertainty is not justified						**
* <sup>125</sup> Sb <sup>n</sup>	J : E2 to 15/2-						**
* <sup>125</sup> Sb <sup>n</sup>	T : average 19Bi04=28.8(0.6) 07Ju06=28.0(0.7)						**
* <sup>125</sup> Sb <sup>g</sup>	T : average 19Bi04=278(7) 07Ju06=272(16)						**
* <sup>125</sup> Ba <sup>m</sup>	E : 67.7(0.4) keV above 5/2+ level, estimated at 50#(20#) keV by Nubase						**
* <sup>125</sup> Ce <sup>m</sup>	T : symmetrized from 07Su07=134(+641-61) s for a fully ionized ion and						**
* <sup>125</sup> Ce <sup>m</sup>	T : $\alpha_T=38.1$ for a 93.6(0.4) keV, E3 transition;						**
* <sup>125</sup> Ce <sup>m</sup>	T : Ensdf2011=3.4(2.7) s						**
<sup>126</sup> Rh	-37200# 500#		19 ms 3	1-#	15	2010	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$
<sup>126</sup> Pd	-51790# 400#		48.6 ms 0.8	0+	15 14SmZZ D	2008	$\beta^- = 100; \beta^- n = 22.9$ *
<sup>126</sup> Pd <sup>m</sup>	-49770# 400#	2023.5 0.7	330 ns 40	(5-)	15 13Wa24 ETJ	2013	IT=100
<sup>126</sup> Pd <sup>n</sup>	-49680# 400#	2109.7 0.9	440 ns 30	(7-)	15 13Wa24 ETJ	2013	IT=100
<sup>126</sup> Pd <sup>p</sup>	-49380# 400#	2406.0 1.0	23.0 ms 0.8	(10+)	15 13Wa24 ETJ	2014	B=72.8; IT=28.8
<sup>126</sup> Ag	-60720# 200#		52 ms 10	3+#	15 14Ba18 TD	1994	$\beta^- = 100; \beta^- n = 13.7. 11$ *
<sup>126</sup> Ag <sup>m</sup>	-60620# 220#	100# 100#	108.4 ms 2.4	9-#	15 14Ba18 TD	1995	$\beta^- = 100; IT ?; \beta^- n ?$ *
<sup>126</sup> Ag <sup>n</sup>	-60470# 200#	254.8 0.5	27 $\mu$ s 6	1-#	15 13La11 JTD	2012	IT=100
<sup>126</sup> Cd	-72255.7 2.3		512 ms 5	0+	03 18Ha30 T	1978	$\beta^- = 100$ *
<sup>126</sup> In	-77809 4		1.53 s 0.01	3+*	03	1974	$\beta^- = 100$
<sup>126</sup> In <sup>m</sup>	-77719 5	90 7 MD	1.64 s 0.05	8-*	03	1970	$\beta^- = 100$
<sup>126</sup> In <sup>n</sup>	-77566 4	243.3 0.2	22 $\mu$ s 2	1-	04Sc42 ETJ	2003	IT=100
<sup>126</sup> Sn	-86015 11		230 ky 14	0+	03	1962	$\beta^- = 100$
<sup>126</sup> Sn <sup>m</sup>	-83796 11	2218.99 0.08	6.1 $\mu$ s 0.7	7-	03 12As05 T	1979	IT=100 *
<sup>126</sup> Sn <sup>n</sup>	-83451 11	2564.5 0.5	7.6 $\mu$ s 0.3	10+	03 12As05 TJ	2000	IT=100 *
<sup>126</sup> Sn <sup>p</sup>	-81668 11	4347.4 0.4	114 ns 2	15-	14Is04 EJT	2012	IT=100 *
<sup>126</sup> Sb	-86390 30		12.35 d 0.06	8-	03 76Sm01 J	1956	$\beta^- = 100$
<sup>126</sup> Sb <sup>m</sup>	-86370 30	17.7 0.3	19.15 m 0.08	5+	03 76Sm01 JD	1956	$\beta^- = 86.4; IT = 14.4$
<sup>126</sup> Sb <sup>n</sup>	-86350 30	40.4 0.3	$\sim 11$ s	3-	03 76Sm01 JD	1976	IT=100[gs=0,m=100]
<sup>126</sup> Sb <sup>p</sup>	-86290 30	104.6 0.3	553 ns 5	3+	03 76Sm01 JD	1976	IT=100
<sup>126</sup> Sb <sup>g</sup>	-84580 30	1810.7 1.7	90 ns 16	(13+)	19Bi04 EJT	2019	IT=100
<sup>126</sup> Te	-90064.2 1.4		STABLE	0+	03	1924	IS=8.84 25
<sup>126</sup> I	-87910 4		12.93 d 0.05	2-*	03	1938	$\beta^+ = 52.7. 5; \beta^- = 47.3. 5$
<sup>126</sup> I <sup>m</sup>	-87799 4	111.00 0.23	128 ns	3+	12Mo.A EJT	2012	IT=100
<sup>126</sup> Xe	-89146.387 0.006		STABLE	0+	03	1922	IS=0.089 3; $\beta^+ ?$ *
<sup>126</sup> Cs	-84351 10		1.64 m 0.02	1+*	03	1954	$\beta^+ = 100$
<sup>126</sup> Cs <sup>m</sup>	-84078 10	273.0 0.7	$\sim 1 \mu$ s	(4-)	03 91TaZX TD	1993	IT=100 *
<sup>126</sup> Cs <sup>n</sup>	-83755 10	596.1 1.1	171 $\mu$ s 14	8-#	03 07Wa09 J	1993	IT=100 *
<sup>126</sup> Ba	-82670 12		100 m 2	0+	03	1954	$\beta^+ = 100$
<sup>126</sup> La	-74970 90		54 s 2	5-#	03	1961	$\beta^+ = 100$
<sup>126</sup> La <sup>m</sup>	-74760 400	210 410 BD*	20 s 20	1-#	03	1997	$\beta^+ = 100$ *
<sup>126</sup> Ce	-70821 28		51.0 s 0.3	0+	03	1978	$\beta^+ = 100$
<sup>126</sup> Pr	-60320# 200#		3.12 s 0.18	(4,5)	03 88Ba42 T	1983	$\beta^+ = 100; \beta^+ p ?$ *
<sup>126</sup> Nd	-53380# 300#		1# s >200ns	0+	03 00So11 I	2000	$\beta^+ ?; \beta^+ p ?$
<sup>126</sup> Pm	-39750# 500#		500# ms				$\beta^+ ?; \beta^+ p ?$
* <sup>126</sup> Pd	T : from 15Lo04, 14Wa26=48.6(0.8); other 14SmZZ=56(+11-9)						**
* <sup>126</sup> Ag	D : $\% \beta^- n$ from 14SmZZ, probably includes gs and isomer						**
* <sup>126</sup> Ag <sup>m</sup>	T : average 14Ba18=92(9) 15Lo04=98(5) 14SmZZ=114(3) 95Fe12=107(12)						**
* <sup>126</sup> Cd	T : average 18Ha30=515(17) 15Lo04=513(6) 78Ga18=506(15); other 86Go10=600(30)						**
* <sup>126</sup> Sn <sup>m</sup>	T : average 12As05=6.6(1.4) 10Il01=5.9(0.8)						**
* <sup>126</sup> Sn <sup>n</sup>	T : average 12As05=7.7(0.5) 10Il01=7.5(0.3)						**
* <sup>126</sup> Sn <sup>p</sup>	T : other 12As05=160(20) not used (at variance)						**
* <sup>126</sup> Xe	T : 2nu- $\epsilon\epsilon$ 16Ab03>4.7Zy (90% CL); 2nu- $\epsilon$ (K) $\epsilon$ (K) 19Ab04>19 Zy						**
* <sup>126</sup> Cs <sup>m</sup>	T : 91TaZX $\leq$ 1 us; 218-keV and 241-keV gamma-ray transition below						**
* <sup>126</sup> Cs <sup>m</sup>	T : the (4-) isomer show the same lifetime, 166(15) us and 176(15) us,						**
* <sup>126</sup> Cs <sup>m</sup>	T : respectively, which is identical to <sup>126</sup> Cs <sup>n</sup>						**
* <sup>126</sup> Cs <sup>n</sup>	D : 112-keV and 223-keV gamma rays to 5- and 6- members of K=4- band in						**
* <sup>126</sup> Cs <sup>n</sup>	D : 93Ko25 and 91TaZX						**
* <sup>126</sup> La <sup>m</sup>	T : 97As05: "by far shorter than 50 s"						**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>126</sup> Pr	T : average 95Os03=3.14(0.22) 88Ba42=3.0(0.4) 83Ni05=3.2(0.6)						**
<sup>127</sup> Rh	-33730#	600#	28 ms 14	7/2 <sup>+</sup> #	15 15Lo04	TD 2015	$\beta^- = 100; \beta^- n ?; \beta^- 2n ?$
<sup>127</sup> Pd	-47220#	500#	38 ms 2	11/2 <sup>-</sup> #	15 14SmZZ	TD 2010	$\beta^- = 100; \beta^- n < 19; \beta^- 2n ?$
<sup>127</sup> Pd <sup>m</sup>	-45500#	500#	1717.91 0.23	(19/2 <sup>+</sup> )	15 19Wa14	ETJ 2019	IT=100
<sup>127</sup> Ag	-58650#	200#	89 ms 2	(9/2 <sup>+</sup> )*	11 14SmZZ	D 1995	$\beta^- = 100; \beta^- n = 14.6 15$
<sup>127</sup> Ag <sup>m</sup>	-58630#	200#	20# ms	(1/2 <sup>-</sup> )*	00Kr18	J	$\beta^- ?; IT ?$
<sup>127</sup> Ag <sup>n</sup>	-56710#	200#	1938 17	(27/2 <sup>+</sup> )	21Wa.A	DJT 2021	$\beta^- = 91.2 8; IT = 8.8 8$
<sup>127</sup> Cd	-68741	6	480 ms 100	3/2 <sup>+</sup> *	11 13Yo02	J 1986	$\beta^- = 100; \beta^- n ?$
<sup>127</sup> Cd <sup>m</sup>	-68456	4	285 8 MD	11/2 <sup>-</sup> *	13Yo02	J 2019	$\beta^- = 100$
<sup>127</sup> Cd <sup>n</sup>	-66896	10	1845 8	(19/2 <sup>+</sup> )	10Na17	ETJ 2010	IT=100
<sup>127</sup> In	-76880	10	1.086 s 0.007	9/2 <sup>+</sup> *	11 93Ru01	TD 1975	$\beta^- = 100; \beta^- n < 0.03$
<sup>127</sup> In <sup>m</sup>	-76486	15	394 18	1/2 <sup>-</sup> #	11 93Ru01	TD 1974	$\beta^- = 100; \beta^- n = 0.70 4$
<sup>127</sup> In <sup>n</sup>	-75110	40	1770 40 MD	(21/2 <sup>-</sup> )	11 18Ba08	E 2004	$\beta^- = 100; \beta^- n ?$
<sup>127</sup> In <sup>p</sup>	-74515	10	2364.7 0.9	(29/2 <sup>+</sup> )	11 04Sc42	ETJ 2004	IT=100
<sup>127</sup> Sn	-83470	9	2.10 h 0.04	11/2 <sup>-</sup> *	11 20Yo.A	J 1951	$\beta^- = 100$
<sup>127</sup> Sn <sup>m</sup>	-83465	9	5.07 0.06	3/2 <sup>+</sup> *	11 20Yo.A	J 1962	$\beta^- = 100$
<sup>127</sup> Sn <sup>n</sup>	-81643	9	1826.67 0.16	19/2 <sup>+</sup>	11 08Lo07	J 2000	IT=100
<sup>127</sup> Sn <sup>p</sup>	-81539	9	1930.97 0.17	(23/2 <sup>+</sup> )	11	2004	IT=100
<sup>127</sup> Sn <sup>q</sup>	-80918	9	2552.4 1.0	(27/2 <sup>-</sup> )	11 08Lo07	J 2008	IT=100
<sup>127</sup> Sb	-86698	5	3.85 d 0.05	7/2 <sup>+</sup>	11	1939	$\beta^- = 100$
<sup>127</sup> Sb <sup>m</sup>	-84778	5	1920.19 0.21	15/2 <sup>-</sup>	11 19Bi04	T 1974	IT=100
<sup>127</sup> Sb <sup>n</sup>	-84373	5	2324.7 0.4	23/2 <sup>+</sup>	11 19Bi04	T 2005	IT=100
<sup>127</sup> Te	-88280.5	1.4	9.35 h 0.07	3/2 <sup>+</sup>	11	1938	$\beta^- = 100$
<sup>127</sup> Te <sup>m</sup>	-88192.3	1.4	88.23 0.07	11/2 <sup>-</sup>	11 17Ni03	D 1940	IT=97.86 3; $\beta^- = 2.14 3$
<sup>127</sup> I	-88983	4	STABLE	5/2 <sup>+</sup> *	11	1920	IS=100
<sup>127</sup> Xe	-88321	4	36.342 d 0.003	1/2 <sup>+</sup> *	11 FGK209	T 1950	$\epsilon = 100$
<sup>127</sup> Xe <sup>m</sup>	-88024	4	297.10 0.08	9/2 <sup>-</sup> *	11 90NeZY	J 1940	IT=100
<sup>127</sup> Cs	-86240	6	6.25 h 0.10	1/2 <sup>+</sup> *	11	1950	$\beta^+ = 100$
<sup>127</sup> Cs <sup>m</sup>	-85788	6	452.23 0.21	(11/2 <sup>-</sup> )	11	1980	IT=100
<sup>127</sup> Ba	-82818	11	12.7 m 0.4	1/2 <sup>+</sup> *	11	1952	$\beta^+ = 100$
<sup>127</sup> Ba <sup>m</sup>	-82738	11	80.32 0.11	7/2 <sup>-</sup>	11	1992	IT=100
<sup>127</sup> La	-77896	26	5.1 m 0.1	(11/2 <sup>-</sup> )	11	1963	$\beta^+ = 100$
<sup>127</sup> La <sup>m</sup>	-77882	26	14.2 0.4	(3/2 <sup>+</sup> )	11	1963	$\beta^+ \approx 100$
<sup>127</sup> Ce	-71979	29	34 s 2	(1/2 <sup>+</sup> )	11	1978	$\beta^+ = 100$
<sup>127</sup> Ce <sup>m</sup>	-71972	29	7.3 1.1	(5/2 <sup>+</sup> )	11	1978	$\beta^+ = 100$
<sup>127</sup> Ce <sup>n</sup>	-71942	29	36.9 1.1	(7/2 <sup>-</sup> )	11	1995	IT=100
<sup>127</sup> Pr	-64540#	200#	> 10 $\mu$ s	3/2 <sup>+</sup> #	11	1995	$\beta^+ = 100$
<sup>127</sup> Pr <sup>m</sup>	-63940#	280#	600# 200#	(11/2 <sup>-</sup> )	11 98Mo30	J 1998	IT ?
<sup>127</sup> Nd	-55910#	300#	1.8 s 0.4	5/2 <sup>+</sup> #	11	1983	$\beta^+ = 100; \beta^+ p = ?$
<sup>127</sup> Pm	-45310#	400#	1# s	3/2 <sup>+</sup> #			$\beta^+ ?; p ?$
* <sup>127</sup> Rh	T : symmetrized from 15Lo04=20(+20-7)						**
* <sup>127</sup> Pd	T : from 15Lo04; other 14SmZZ=73(+24-23)						**
* <sup>127</sup> Ag	T : from 15Lo04; other 96Wo.A=79(3), supersedes 95Fe12=109(25), same group						**
* <sup>127</sup> Ag	J : 00Kr18=(9/2) g <sub>9/2</sub> ; hfs comparison to <sup>107,107m</sup> Ag						**
* <sup>127</sup> Ag <sup>m</sup>	J : 00Kr18=(1/2) p <sub>1/2</sub> ; hfs comparison to <sup>107,107m</sup> Ag						**
* <sup>127</sup> Ag <sup>n</sup>	E : from 21Wa.A=150(+14-20) keV above the 1792.2(0.9) keV						**
* <sup>127</sup> Cd	T : symmetrized 19Lo04=450(+120-80); other 15Lo04=330(20) mixed states						**
* <sup>127</sup> Cd <sup>m</sup>	T : from 19Lo04						**
* <sup>127</sup> Cd <sup>n</sup>	E : 1560.1(0.5) keV above <sup>127</sup> Cd <sup>m</sup>						**
* <sup>127</sup> Cd <sup>n</sup>	T : others 21Wa.A=18(1) 12Ka36=11.0(+9.2-3.5)						**
* <sup>127</sup> In	T : average 93Ru01=1.083(0.007) 86Go10=1.22(0.05) 83Sh07=0.99(0.10)						**
* <sup>127</sup> In	T : 81En05=1.10(0.04); others 80Lu04=1.12(0.02), superseded by 93Ru01,						**
* <sup>127</sup> In	T : 02Pf04=1.090(0.010), compilation						**
* <sup>127</sup> In <sup>m</sup>	T : average 93Ru01=3.580(0.025) 86ReZU=3.70(0.04) 83Sh07=3.76(0.31)						**
* <sup>127</sup> In <sup>m</sup>	T : 80De35=3.7(0.1); other 02Pf04=3.67(4), compilation						**
* <sup>127</sup> In <sup>m</sup>	D : % $\beta^- n$ average 93Ru01=0.72(0.04) 86ReZU=0.54(0.11)						**
* <sup>127</sup> Sb <sup>m</sup>	T: other 74Ap01=11(1)						**
* <sup>127</sup> Sb <sup>n</sup>	T : others 09Wa24=234(12) 05Po03=165(20) outlier, not used						**
* <sup>127</sup> Xe	J : 90NeZY=1/2						**
* <sup>127</sup> Xe <sup>m</sup>	J : 90NeZY=9/2						**
* <sup>127</sup> Ce <sup>n</sup>	E : 95Os03=29.56(0.05) keV above <sup>127</sup> Ce <sup>m</sup>						**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
* <sup>127</sup> Pr <sup>m</sup>	T : estimated by FGK208 from BE3(11/2- to 5/2+)=41.2(2.0) [W.u.] in							
* <sup>127</sup> Pr <sup>m</sup>	T : <sup>131</sup> Pr							
<sup>128</sup> Rh	-27340#	300#	8# ms >550ns		18Sh11 I	2018	$\beta^- ?; \beta^- n ?; \beta^- 2n ?$	
<sup>128</sup> Pd	-44390#	500#	35 ms 3	0 <sup>+</sup>	16 15Lo04 T	2010	$\beta^- =100; \beta^- n ?$	
<sup>128</sup> Pd <sup>m</sup>	-42240#	500#	2151.0 1.0	(8 <sup>+</sup> )	16	2013	IT=100	
<sup>128</sup> Ag	-54710#	300#	60 ms 3	3 <sup>+</sup> #	15 14SmZZ TD	2000	$\beta^- =100; \beta^- n =20 5; \beta^- 2n ?$	
<sup>128</sup> Cd	-67238	6	246 ms 2	0 <sup>+</sup>	15 16Du13 T	1986	$\beta^- =100; \beta^- n ?$	
<sup>128</sup> Cd <sup>m</sup>	-65368	6	1870.5 0.3	(5 <sup>-</sup> )	15	2009	IT=100	
<sup>128</sup> Cd <sup>n</sup>	-64523	6	2714.6 0.4	(10 <sup>+</sup> )	15	2009	IT=100	
<sup>128</sup> Cd <sup>p</sup>	-62951	6	4286.6 1.5	(15 <sup>-</sup> )	17Ju02 ETJ	2016	IT=100	
<sup>128</sup> In	-74190.1	1.3	816 ms 27	(3 <sup>+</sup> )	15 93Ru01 D	1975	$\beta^- =100; \beta^- n =0.038 3$	
<sup>128</sup> In <sup>m</sup>	-73942.2	1.3	247.87 0.10	(1 <sup>-</sup> )	15 04Sc42 J	1988	IT=100	
<sup>128</sup> In <sup>n</sup>	-73905.0	2.0	285.1 2.2 MD	(8 <sup>-</sup> )	15 20Ne06 E	1986	$\beta^- =100; IT ?; \beta^- n ?$	
<sup>128</sup> In <sup>p</sup>	-72392.5	1.4	1797.6 1.6 MD	(16 <sup>+</sup> )	20Ne06 EJT	2020	$\beta^- \approx 100; IT ?; \beta^- n ?$	
<sup>128</sup> Sn	-83361	18	59.07 m 0.14	0 <sup>+</sup>	15	1956	$\beta^- =100$	
<sup>128</sup> Sn <sup>m</sup>	-81270	18	2091.50 0.11	6.5 s 0.5	7 <sup>-</sup>	15 1979	IT=100	
<sup>128</sup> Sn <sup>n</sup>	-80869	18	2491.91 0.17	2.91 $\mu$ s 0.14	10 <sup>+</sup>	15 1981	IT=100	
<sup>128</sup> Sn <sup>p</sup>	-79262	18	4099.5 0.4	220 ns 30	(15 <sup>-</sup> )	15 2011	IT=100	
<sup>128</sup> Sb	-84630	19	*	9.05 h 0.04	8 <sup>-*</sup>	15 1956	$\beta^- =100$	
<sup>128</sup> Sb <sup>m</sup>	-84620	18	6	10.41 m 0.18	5 <sup>+</sup>	15 1955	$\beta^- =96.4 10; IT =3.6 10$	
<sup>128</sup> Sb <sup>n</sup>	-83013	19	1617.3 0.7	500 ns 20	(11 <sup>+</sup> )	19Bi04 ETJ	2019	IT=100
<sup>128</sup> Sb <sup>p</sup>	-82860	19	1769.9 1.2	217 ns 7	(13 <sup>+</sup> )	19Bi04 ETJ	2019	IT=100
<sup>128</sup> Te	-88993.8	0.7	2.25 Yy 0.09	0 <sup>+</sup>	15 20Ba.A T	1924	IS=31.74 8; 2 $\beta^- =100$	
<sup>128</sup> Te <sup>m</sup>	-86203.0	0.8	2790.8 0.3	363 ns 27	(10 <sup>+</sup> )	15 04Va03 T	1998	IT=100
<sup>128</sup> I	-87738	4	24.99 m 0.02	1 <sup>+</sup> *	15	1934	$\beta^- =93.1 8; \beta^+ =6.9 8$	
<sup>128</sup> I <sup>m</sup>	-87600	4	137.851 0.003	845 ns 20	4 <sup>-</sup>	15 1982	IT=100	
<sup>128</sup> I <sup>n</sup>	-87571	4	167.368 0.004	175 ns 15	(6 <sup>-</sup> )	15 1991	IT=100	
<sup>128</sup> Xe	-89860.534	0.005	STABLE	0 <sup>+</sup>	15	1922	IS=1.910 13	
<sup>128</sup> Xe <sup>m</sup>	-87073.3	0.3	2787.2 0.3	83 ns 2	8 <sup>-</sup>	15 1981	IT=100	
<sup>128</sup> Cs	-85932	5	3.640 m 0.014	1 <sup>+</sup> *	15 93Al03 T	1951	$\beta^+ =100$	
<sup>128</sup> Ba	-85369.2	1.6	2.43 d 0.05	0 <sup>+</sup>	15	1950	$\epsilon =100$	
<sup>128</sup> La	-78630	50	*	5.18 m 0.14	(5 <sup>+</sup> )	15 97Ha30 T	1961	$\beta^+ =100$
<sup>128</sup> La <sup>m</sup>	-78530#	110#	100# 100# *	< 1.4 m	(1 <sup>+</sup> , 2 <sup>-</sup> )	15 97Ha30 T	1995	$\beta^+ =100; IT ?$
<sup>128</sup> Ce	-75534	28	3.93 m 0.02	0 <sup>+</sup>	15 00Li08 T	1968	$\beta^+ =100$	
<sup>128</sup> Pr	-66331	30	2.85 s 0.09	(3 <sup>+</sup> )	15 99Xi03 J	1985	$\beta^+ =100; \beta^+ p =?$	
<sup>128</sup> Nd	-60530#	200#	5# s	0 <sup>+</sup>	15	1985	$\beta^+ ?$	
<sup>128</sup> Pm	-48220#	300#	1.0 s 0.3	4 <sup>+</sup> #	15 93Li40 D	1999	$\beta^+ \approx 100; \beta^+ p = ?; p =0$	
<sup>128</sup> Sm	-39150#	500#	500# ms	0 <sup>+</sup>			$\beta^+ ?; \beta^+ p ?$	
* <sup>128</sup> Pd	T : other 14SmZZ < 262 ms							
* <sup>128</sup> Ag	T : average 15Lo04=59(5) 14SmZZ=73(+10-9) 96Wo.A=58(5) 95Kr.A=60(10)							
* <sup>128</sup> Cd	T : average 16Du13=246.2(2.1) 15Lo04=245(5)							
* <sup>128</sup> In	T : average 15Lo04=810(30) 86Go10=840(60)							
* <sup>128</sup> In	D : % $\beta^- n$ from 93Ru01, probably includes gs and isomer							
* <sup>128</sup> Sn <sup>m</sup>	J : E3 to 4+							
* <sup>128</sup> Sn <sup>n</sup>	J : E2 to 8+							
* <sup>128</sup> Sb <sup>m</sup>	E : less than 20 keV above the ground state; see Ensdf2015 for details							
* <sup>128</sup> Te <sup>m</sup>	T : average 04Va03=337(59) 98Zh09=370(30)							
* <sup>128</sup> Cs	T : average 93Al03=3.66(0.02) 76He04=3.62(0.02)							
* <sup>128</sup> La	T : average 97Ha30=5.4(0.2) 77Zo02=5.2(0.4) 66Pa06=4.9(0.4) 66Li04=4.9(0.4)							
* <sup>128</sup> Ce	T : average 00Li08=4.0(0.1) 97Ha30=4.1(0.3) 97As05=3.925(0.021)							
* <sup>128</sup> Pr	T : average 99Xi03=2.8(0.1) 88Ba42=3.1(0.3) 85Wi07=3.2(+0.5-0.4)							
* <sup>128</sup> Pr	D : $\beta^+ p$ observed in 85Wi07, but was not quantified							
* <sup>128</sup> Nd	T : 83Ni05=4(2)s, but 85Wi07 associated it with decay of <sup>128</sup> Pr							
* <sup>128</sup> Pm	D : %p 93Li40=0							
<sup>129</sup> Pd	-37880#	600#	31 ms 7	7/2 <sup>-</sup> #	15	2015	$\beta^- =100; \beta^- n ?; \beta^- 2n ?$	
<sup>129</sup> Ag	-51870#	400#	49.9 ms 3.5	9/2 <sup>+</sup> #	14 14SmZZ D	2000	$\beta^- =100; \beta^- n < 20$	
<sup>129</sup> Ag <sup>m</sup>	-51850#	400#	20# 20# *	1/2 <sup>-</sup> #	14		$\beta^- ?; \beta^- n ?$	
<sup>129</sup> Cd	-63122	5	147 ms 3	11/2 <sup>-*</sup>	14 16Du13 T	2003	$\beta^- =100; \beta^- n =?$	
<sup>129</sup> Cd <sup>m</sup>	-62779	6	343 8 MD	3/2 <sup>+</sup> *	14 20Ma09 E	1986	$\beta^- =100; \beta^- n =?$	

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{129}\text{Cd}^n$	-60839	9	2283	8	3.6 ms 0.2	(21/2 <sup>+</sup> )	14 14Ta29 TJE 2014	IT=100 *
$^{129}\text{In}$	-72834.9	2.0			570 ms 10	9/2 <sup>+</sup>	14 15Lo04 T 1975	$\beta^- = 100; \beta^-_n = 0.23\ 7$ *
$^{129}\text{In}^m$	-72384.2	2.0	450.73	0.16	1.23 s 0.03	1/2 <sup>-</sup>	14 04Ga24 J 1976	$\beta^- \approx 100; IT \ ?; \beta^-_n = 3.6\ 4$ *
$^{129}\text{In}^n$	-71146.9	2.0	1687.97	0.25	11.2 $\mu$ s 0.2	(17/2 <sup>-</sup> )	14 14Ta.A T 2003	IT=100
$^{129}\text{In}^p$	-71180	50	1650	50	670 ms 100	(23/2 <sup>-</sup> )	14 04Ga24 ETJ 2004	$\beta^- \approx 100; IT \ ?$
$^{129}\text{In}^q$	-70890	50	1941	50	110 ms 15	(29/2 <sup>+</sup> )	14	IT $\approx$ 100; $\beta^- \ ?$ *
$^{129}\text{Sn}$	-80591	17			2.23 m 0.04	3/2 <sup>+</sup> *	14 20Yo.A J 1962	$\beta^- = 100$
$^{129}\text{Sn}^m$	-80556	17	35.15	0.05	6.9 m 0.1	11/2 <sup>-*</sup>	14 20Yo.A J 1962	$\beta^- = 100$
$^{129}\text{Sn}^n$	-78829	17	1761.6	1.0	3.49 $\mu$ s 0.11	(19/2 <sup>+</sup> )	14 08Lo07 T 2000	IT=100 *
$^{129}\text{Sn}^p$	-78788	17	1802.6	1.0	2.22 $\mu$ s 0.13	23/2 <sup>+</sup>	14 08Lo07 TJ 2000	IT=100 *
$^{129}\text{Sn}^q$	-78038	17	2552.9	1.1	221 ns 18	(27/2 <sup>-</sup> )	14 08Lo07 J 2008	IT=100
$^{129}\text{Sb}$	-84629	21			4.366 h 0.026	7/2 <sup>+</sup>	14	1939 $\beta^- = 100$
$^{129}\text{Sb}^m$	-82778	21	1851.31	0.06	17.7 m 0.1	19/2 <sup>-</sup>	14 19Bi04 J 1982	$\beta^- = 85; IT = 15$
$^{129}\text{Sb}^n$	-82768	21	1861.06	0.05	2.23 $\mu$ s 0.17	15/2 <sup>-</sup>	14 19Bi04 TJ 1987	IT=100 *
$^{129}\text{Sb}^p$	-82490	21	2139.4	0.3	0.89 $\mu$ s 0.03	23/2 <sup>+</sup>	14 19Bi04 TJ 2003	IT=100
$^{129}\text{Te}$	-87004.9	0.7			69.6 m 0.3	3/2 <sup>+</sup>	14	1939 $\beta^- = 100$
$^{129}\text{Te}^m$	-86899.4	0.7	105.51	0.03	33.6 d 0.1	11/2 <sup>-</sup>	14	1940 IT=64 7; $\beta^- = 36\ 7$
$^{129}\text{I}$	-88507	3			16.14 My 0.12	7/2 <sup>+</sup> *	14 18Ga37 T 1951	$\beta^- = 100$
$^{129}\text{Xe}$	-88696.070	0.005			STABLE	1/2 <sup>+</sup> *	14	1920 IS=26.401 138
$^{129}\text{Xe}^m$	-88459.93	0.03	236.14	0.03	8.88 d 0.02	11/2 <sup>-*</sup>	14 90NeZY J 1951	IT=100 *
$^{129}\text{Cs}$	-87499	5			32.06 h 0.06	1/2 <sup>+</sup> *	14	1950 $\beta^+ = 100$
$^{129}\text{Cs}^m$	-86924	5	575.40	0.14	718 ns 21	(11/2 <sup>-</sup> )	14	1977 IT=100
$^{129}\text{Ba}$	-85061	11			2.23 h 0.11	1/2 <sup>+</sup>	14	1950 $\beta^+ = 100$
$^{129}\text{Ba}^m$	-85053	11	8.42	0.06	2.135 h 0.010	7/2 <sup>+</sup>	14	1950 $\beta^+ \approx 100; IT = ?$
$^{129}\text{La}$	-81324	21			11.6 m 0.2	(3/2 <sup>+</sup> )	14	1963 $\beta^+ = 100$
$^{129}\text{La}^m$	-81152	21	172.33	0.20	560 ms 50	(11/2 <sup>-</sup> )	14	1969 IT=100
$^{129}\text{Ce}$	-76288	28			3.5 m 0.3	(5/2 <sup>+</sup> )	14	1977 $\beta^+ = 100$
$^{129}\text{Pr}$	-69774	30			30 s 4	(3/2 <sup>+</sup> )	14 96Gi08 J 1977	$\beta^+ = 100$
$^{129}\text{Pr}^m$	-69390	30	382.57	0.24	26# $\mu$ s 11#	(11/2 <sup>-</sup> )	14 FGK208 T 1997	IT=100 *
$^{129}\text{Nd}$	-62380#	200#			6.8 s 0.6	7/2 <sup>-#</sup>	14 10Xu12 T 1977	$\beta^+ = 100; \beta^+_p = ?$ *
$^{129}\text{Nd}^m$	-62330#	220#	50#	100#	2.6 s 0.4	1/2 <sup>+#</sup>	14	2010 $\beta^+ = 100; \beta^+_p = ?$
$^{129}\text{Pm}$	-53180#	300#			2.4 s 0.9	5/2 <sup>+#</sup>	14	2004 $\beta^+ = 100; \beta^+_p \ ?; p \ ?$
$^{129}\text{Sm}$	-42330#	500#			550 ms 100	(1/2 <sup>+</sup> , 3/2 <sup>+</sup> )	14	1999 $\beta^+ = 100; \beta^+_p = ?$
* $^{129}\text{Ag}$	T : average 15Lo04=52(4) 00Kr18=46(+5-9) **							
* $^{129}\text{Ag}$	D : $\beta^-_n$ was observed by 00Kr18, but it was not quantified **							
* $^{129}\text{Ag}^m$	T : 00Kr18=158(60) ms is not convincing; not adopted in 02Pf04 (same group) **							
* $^{129}\text{Cd}$	T : from 16Du13=147(3) gated by gammas's in the high-spin decay; others **							
* $^{129}\text{Cd}$	T : 15Ta13=155(3) 09Ar04=104(6) **							
* $^{129}\text{Cd}$	D : $\beta^-_n$ was observed by 05Kr20, but it was not quantified **							
* $^{129}\text{Cd}$	J : 13Yo02=11/2 **							
* $^{129}\text{Cd}^m$	D : $\beta^-_n$ was observed by 05Kr20, but it was not quantified **							
* $^{129}\text{Cd}^m$	T : from 16Du13=157(8) gated by gammas's in the low-spin decay; others **							
* $^{129}\text{Cd}^m$	T : 15Ta13=146(8) 15Lo04=154.5(2.0), mixture of two states, 09Ar04=242(8) **							
* $^{129}\text{Cd}^m$	J : 13Yo02=3/2 **							
* $^{129}\text{Cd}^n$	E : 14Ta29=1940.0(1.4) keV above $^{129}\text{Cd}^m$ **							
* $^{129}\text{In}$	J : from 04Ga24 **							
* $^{129}\text{In}^m$	E : from a least-squares fit to the level scheme in 15Ta13; others **							
* $^{129}\text{In}^m$	E : 13Ka08=459(5) 18Ba08=444(15), direct mass measurements **							
* $^{129}\text{In}^q$	E : 281.0(0.2) keV gamma above $^{129}\text{In}^p$ **							
* $^{129}\text{Sn}^n$	T : average 08Lo07=3.4(0.4) 04Ga24=3.2(0.2) 00Pi03=3.7(0.2) **							
* $^{129}\text{Sn}^n$	T : 00Ge07=3.6(0.2) **							
* $^{129}\text{Sn}^p$	T : average 08Lo07=2.4(4) 04Ga24=2.0(2) 00Ge07=2.4(2) **							
* $^{129}\text{Sb}^n$	T : average 19Bi04=2.3(0.3) 03Ge04=2.2(0.2) **							
* $^{129}\text{Xe}^m$	J : 90NeZY=11/2 **							
* $^{129}\text{Pr}^m$	T : estimated by FGK208 from BE3(11/2- to 5/2+)=41.2(2.0) [W.u.] in **							
* $^{129}\text{Pr}^m$	T : $^{131}\text{Pr}$ using the known branching intensities and $\alpha_T$ **							
* $^{129}\text{Nd}$	T : average 10Xu12=6.7(0.7) 97Gi07=7(1); other 85Wi07=4.9(0.2) is **							
* $^{129}\text{Nd}$	T : for a mixture between the gs and the isomer **							
$^{130}\text{Pd}$	-32730#	300#			27# ms >550ns	0 <sup>+</sup>	18Sh11 I 2018	$\beta^- \ ?; \beta^-_n \ ?; \beta^-_{2n} \ ?$
$^{130}\text{Ag}$	-45900#	420#			40.6 ms 4.5	1 <sup>-#</sup>	15 15Lo04 T 2000	$\beta^- = 100; \beta^-_n \ ?; \beta^-_{2n} \ ?$ *
$^{130}\text{Cd}$	-61118	22			126.8 ms 1.8	0 <sup>+</sup>	08 16Du13 T 1986	$\beta^- = 100; \beta^-_n = 3.5\ 10$ *



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)		
$^{130}\text{Cd}^m$	-58988	22	2129.6	1.0	240 ns 16	(8 <sup>+</sup> )	08 12Ka36 ET	2007	IT=100	*	
$^{130}\text{In}$	-69906.5	1.8		*	273 ms 5	1(⁻)	08 15Lo04 T	1973	$\beta^- = 100; \beta^-_n = 0.93$	13 *	
$^{130}\text{In}^m$	-69840.0	2.0	66.5	2.7	540 ms 10	(10 <sup>-</sup> )	08 20Ne06 E	1973	$\beta^- = 100; \beta^-_n = 1.80$	7 *	
$^{130}\text{In}^n$	-69521.1	2.1	385.4	2.6	540 ms 10	(5 <sup>+</sup> )	08 20Ne06 E	1986	$\beta^- = 100; \beta^-_n = 1.80$	7 *	
$^{130}\text{In}^p$	-69518.2	1.8	388.3	0.2	4.6 $\mu$ s 0.2	(3 <sup>+</sup> )	08 16Ju03 T	2003	IT=100	*	
$^{130}\text{Sn}$	-80132.2	1.9			3.72 m 0.07	0 <sup>+</sup>	01	1972	$\beta^- = 100$		
$^{130}\text{Sn}^m$	-78185.3	1.9	1946.88	0.10	1.7 m 0.1	7 <sup>-</sup>	01 05Le34 J	1974	$\beta^- = 100$		
$^{130}\text{Sn}^n$	-77697.4	1.9	2434.79	0.12	1.501 $\mu$ s 0.017	(10 <sup>+</sup> )	01 11Pi05 T	1981	IT=100		
$^{130}\text{Sb}$	-82286	14			39.5 m 0.8	8 <sup>-</sup>	01 02Ge07 J	1962	$\beta^- = 100$		
$^{130}\text{Sb}^m$	-82281	14	4.80	0.20	6.3 m 0.2	4 <sup>+</sup>	01 94WaZU J	1962	$\beta^- = 100$		
$^{130}\text{Sb}^n$	-82201	14	84.67	0.04	800 ns 100	6 <sup>-</sup>	01 02Ge07 TJ	2002	IT=100		
$^{130}\text{Sb}^p$	-80778	14	1508	1	600 ns 15	(11 <sup>+</sup> )	19Bi04 ETJ	2019	IT=100		
$^{130}\text{Sb}^q$	-80741	14	1544.7	0.5	1.25 $\mu$ s 0.01	(13 <sup>+</sup> )	02Ge07 ETJ	2002	IT=100	*	
$^{130}\text{Te}$	-87352.960	0.011			791 Ey 21	0 <sup>+</sup>	01 20Ba.A T	1924	IS=34.08 62; $2\beta^- = 100$		
$^{130}\text{Te}^m$	-85206.55	0.04	2146.41	0.04	186 ns 11	7 <sup>-</sup>	01 04Va03 T	1972	IT=100	*	
$^{130}\text{Te}^n$	-84685.8	0.8	2667.2	0.8	1.90 $\mu$ s 0.08	(10 <sup>+</sup> )	01 04Br19 E	1998	IT=100	*	
$^{130}\text{Te}^p$	-82979.1	0.9	4373.9	0.9	53 ns 8	(15 <sup>-</sup> )	01 14As01 JTE	1998	IT=100	*	
$^{130}\text{I}$	-86936	3			12.36 h 0.01	5 <sup>+</sup> *	01	1938	$\beta^- = 100$		
$^{130}\text{I}^m$	-86896	3	39.9525	0.0013	8.84 m 0.06	2 <sup>+</sup>	01	1966	IT=84 2; $\beta^- = 16$	2 *	
$^{130}\text{I}^n$	-86866	3	69.5865	0.0007	133 ns 7	6 <sup>-</sup>	01	1989	IT=100	*	
$^{130}\text{I}^p$	-86854	3	82.3960	0.0019	315 ns 15	(8 <sup>-</sup> )	01	1989	IT=100		
$^{130}\text{I}^q$	-86851	3	85.1099	0.0010	254 ns 4	6 <sup>-</sup>	01	1975	IT=100	*	
$^{130}\text{Xe}$	-89880.474	0.009			STABLE	0 <sup>+</sup>	01	1922	IS=4.071 22		
$^{130}\text{Cs}$	-86900	8			29.21 m 0.04	1 <sup>+</sup> *	01	1952	$\beta^+ = 98.4; \beta^- = 1.6$		
$^{130}\text{Cs}^m$	-86737	8	163.25	0.11	3.46 m 0.06	5 <sup>-</sup> *	01	1977	IT $\approx$ 100; $\beta^+ = 0.16$	2 *	
$^{130}\text{Cs}^x$	-86873	17	27	15	R = 0.2 0.1	<i>fsmix</i>					
$^{130}\text{Ba}$	-87256.78	0.29			STABLE	$\sim 1Zy$	0 <sup>+</sup>	01 15Ba11 T	1936	IS=0.11 1; $2\beta^+ ?$	
$^{130}\text{Ba}^m$	-84781.7	0.3	2475.12	0.18	9.54 ms 0.14	8 <sup>-</sup>	01 02Mo31 T	1969	IT=100	*	
$^{130}\text{La}$	-81627	26			8.7 m 0.1	3(⁺)	01	1961	$\beta^+ = 100$		
$^{130}\text{La}^m$	-81413	26	214.0	0.5	742 ns 28	(5 <sup>+</sup> )	14Io01 ETJ	2012	IT=100	*	
$^{130}\text{Ce}$	-79423	28			22.9 m 0.5	0 <sup>+</sup>	01	1965	$\beta^+ = 100$		
$^{130}\text{Ce}^m$	-76969	28	2453.6	0.3	100 ns 8	7 <sup>-</sup>	01 99Io02 J	1999	IT=100		
$^{130}\text{Pr}$	-71180	60			40.0 s 0.4	(6,7)(⁺#)	01 88Ba42 J	1977	$\beta^+ = 100$		
$^{130}\text{Pr}^m$	-71080#	120#	100#	100#	10# s	2 <sup>+</sup> #	01 88Ba42 J	1988	$\beta^+ ?$	*	
$^{130}\text{Nd}$	-66596	28			21 s 3	0 <sup>+</sup>	01 01Gi17 T	1977	$\beta^+ = 100$	*	
$^{130}\text{Pm}$	-55470#	200#			2.6 s 0.2	(5 <sup>+</sup> , 6 <sup>+</sup> , 4 <sup>+</sup> )	01 99Xi03 J	1985	$\beta^+ = 100; \beta^+ p = ?$		
$^{130}\text{Sm}$	-47700#	400#			1# s	0 <sup>+</sup>	01	1999	$\beta^+ ?$		
$^{130}\text{Eu}$	-33510#	540#			1.0 ms 0.4	(1 <sup>+</sup> )	08	2004	$p \approx 100; \beta^+ ?; \beta^+ p ?$	*	
* $^{130}\text{Ag}$	T : average 15Lo04=42(5) 05Kr20=35(10)										
* $^{130}\text{Cd}$	T : average 16Du13=126(4) 15Lo04=127(2)										
* $^{130}\text{Cd}^m$	T : average 12Ka36=248(+21-19) 07Ju05=220(30)										
* $^{130}\text{Cd}^n$	E : 12Ka36=128.0(0.5), 138.0(0.5), 538.2(0.5) and 1325.4(0.5)-keV										
* $^{130}\text{Cd}^m$	E : gamma rays in a cascade to gs										
* $^{130}\text{In}$	T : average 15Lo04=284(10) 93Ru01=256(9) 86ReZU=278(7)										
* $^{130}\text{In}$	D : % $\beta^-_n$ average 93Ru01=1.49(0.22) 86Wa17=0.90(0.05), Birge ratio=2.615;										
* $^{130}\text{In}$	D : other 86ReZU=0.91(10), same as 86Wa17										
* $^{130}\text{In}^m$	D : % $\beta^-_n$ average 93Ru01=2.03(0.12) 86Wa17=1.67(0.09), includes both										
* $^{130}\text{In}^m$	D : $^{130}\text{In}^m$ and $^{130}\text{In}^n$ isomers										
* $^{130}\text{In}^m$	J : 81Fo02=(10 <sup>-</sup> ); direct $\beta^-$ decay to 9- agrees with 8-, 9- or 10-										
* $^{130}\text{In}^n$	D : % $\beta^-_n$ average 93Ru01=2.03(0.12) 86Wa17=1.67(0.09), includes both										
* $^{130}\text{In}^n$	D : $^{130}\text{In}^m$ and $^{130}\text{In}^n$ isomers										
* $^{130}\text{In}^p$	T : average 16Ju03=4.4(0.2) 12Ka36=5.25(+0.40-0.35); other 04Sc42=3.1(0.3)										
* $^{130}\text{Sb}^q$	T : from 19Bi04; other 02Ge07=1.8(0.2)										
* $^{130}\text{Te}^m$	T : other 72Ke28=115(11)										
* $^{130}\text{Te}^n$	T : other 98Zh09=4.2(0.9), conflicting data, not used										
* $^{130}\text{Te}^p$	T : others 04Br19=45 98HoZP=261(33)										
* $^{130}\text{I}^n$	J : E1 to 5+										
* $^{130}\text{I}^q$	J : E1 to 5+										
* $^{130}\text{Ba}^m$	T : others 66Br14=8.8(0.2) 69Wa.A=13.5(1.0)										
* $^{130}\text{La}^m$	T : average 14Io01=760(90) 12Ta18=740(30)										
* $^{130}\text{Pr}^m$	J : evidence for a low-spin component in $^{130}\text{Pr}$ activity in 88Ba42										
* $^{130}\text{Nd}$	T : others 00Xu08=13(3) 77Bo02=28(3) conflicting, not used										
* $^{130}\text{Eu}$	T : symmetrized from 0.90(+0.49-0.29)										

Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{131}\text{Pd}$	-25740#	300#	20# ms >550ns	$7/2^-$ #	18Sh11 I	2018	$\beta^-$ ?; $\beta^-n$ ?; $\beta^-2n$ ?
$^{131}\text{Ag}$	-40750#	500#	35 ms 8	$9/2^+$ #	15	2013	$\beta^-$ =100; $\beta^-n$ ?; $\beta^-2n$ =10
$^{131}\text{Cd}$	-55212	19	98 ms 2	$7/2^-$ #	06 15Lo04 T	2000	$\beta^-$ =100; $\beta^-n$ =3.5 10; $\beta^-2n$ ? *
$^{131}\text{In}$	-68024.4	2.2	261.5 ms 2.8	$9/2^+$	06 19Du12 TDJ	1976	$\beta^-$ =100; $\beta^-n$ =2.25 21 *
$^{131}\text{In}^m$	-67648.4	2.5	376 3 MD	$1/2^-$	06 19Or.A E	1984	$\beta^-$ =100; $\beta^-n$ =2.25 21; IT? *
$^{131}\text{In}^n$	-64280	90	322 ms 41	$(21/2^+)$	06 19Du12 TD	1984	$\beta^-$ =100; $\beta^-n$ =12 7; IT? *
$^{131}\text{In}^p$	-64240.8	2.3	669 ns 34	$(17/2^+)$	09Go40 TJ	2009	IT=100 *
$^{131}\text{Sn}$	-77265	4	56.0 s 0.5	$3/2^+$ *	06 20Yo.A J	1963	$\beta^-$ =100
$^{131}\text{Sn}^m$	-77200	4	65.1 0.3	$11/2^-$ *	06 04Fo06 E	1977	$\beta^-$ =100; IT? *
$^{131}\text{Sn}^n$	-72595	4	4670.0 0.4	$(23/2^-)$	06 19Du12 T	2001	IT=100 *
$^{131}\text{Sb}$	-81981.4	2.1	23.03 m 0.04	$7/2^+$	06	1956	$\beta^-$ =100
$^{131}\text{Sb}^m$	-80305.3	2.1	1676.06 0.06	$15/2^-$	06 19Bi04 T	1969	IT=100 *
$^{131}\text{Sb}^n$	-80294.2	2.3	1687.2 0.9	$4.3 \mu\text{s}$ 0.8	06 00Ge18 TJ	2000	IT=100
$^{131}\text{Sb}^p$	-79815.8	2.6	2165.6 1.5	$0.97 \mu\text{s}$ 0.03	06 19Bi04 T	2000	IT=100 *
$^{131}\text{Te}$	-85211.02	0.06	25.0 m 0.1	$3/2^+$	06	1939	$\beta^-$ =100
$^{131}\text{Te}^m$	-85028.76	0.06	182.258 0.018	$11/2^-$	06 08Ea01 T	1940	$\beta^-$ =74.1 5; IT=25.9 5
$^{131}\text{Te}^n$	-83271.0	0.4	1940.0 0.4	$(23/2^+)$	06	1998	IT=100
$^{131}\text{I}$	-87442.7	0.6	8.0249 d 0.0006	$7/2^+$ *	06 FGK209 T	1939	$\beta^-$ =100
$^{131}\text{I}^m$	-85524.3	0.7	24 $\mu\text{s}$ 1	$19/2^-$	09Wa11 EJT	2009	IT=100
$^{131}\text{Xe}$	-88413.575	0.005	STABLE	$3/2^+$ *	06	1920	IS=21.232 51
$^{131}\text{Xe}^m$	-88249.645	0.009	11.948 d 0.012	$11/2^-$ *	06 FGK209 T	1966	IT=100 *
$^{131}\text{Cs}$	-88055.57	0.18	9.689 d 0.016	$5/2^+$ *	06	1947	$\epsilon$ =100
$^{131}\text{Ba}$	-86679.0	0.4	11.52 d 0.01	$1/2^+$ *	06 12Da04 T	1947	$\beta^+$ =100
$^{131}\text{Ba}^m$	-86491.0	0.4	14.26 m 0.09	$9/2^-$ *	06 12Da04 T	1963	IT=100
$^{131}\text{La}$	-83769	28	59 m 2	$3/2^+$ *	06	1951	$\beta^+$ =100
$^{131}\text{La}^m$	-83464	28	170 $\mu\text{s}$ 7	$11/2^-$	06	1966	IT=100
$^{131}\text{Ce}$	-79710	30	10.3 m 0.3	$7/2^+$	06	1966	$\beta^+$ =100
$^{131}\text{Ce}^m$	-79650	30	63.09 0.09	$(1/2^+)$	06 96Gi08 E	1966	$\beta^+$ =100
$^{131}\text{Pr}$	-74300	50	1.50 m 0.03	$(3/2^+)$	06 96Gi08 TJ	1977	$\beta^+$ =100 *
$^{131}\text{Pr}^m$	-74150	50	5.73 s 0.20	$(11/2^-)$	06	1996	IT=96.4 12; $\beta^+$ =3.6 12
$^{131}\text{Nd}$	-67768	28	25.4 s 0.9	$(5/2^+)$	06	1977	$\beta^+$ =100; $\beta^+p$ =?
$^{131}\text{Pm}$	-59770#	200#	6.3 s 0.8	$(11/2^-)$	06 99Ga41 T	1998	$\beta^+$ =100
$^{131}\text{Sm}$	-50280#	400#	1.2 s 0.2	$5/2^+$ #	06	1986	$\beta^+$ =100; $\beta^+p$ =?
$^{131}\text{Eu}$	-39460#	400#	17.8 ms 1.9	$3/2^+$	06	1998	$p$ =89 9; $\beta^+$ ?; $\beta^+p$ ?
* $^{131}\text{Cd}$	T: 15Lo04=98.0(0.2) is a typo; other 00Ha55=68(3) **						
* $^{131}\text{In}$	D: % $\beta^-n$ average 93Ru01=2.2(0.3) 19Du12 $\beta^-n$ =2.3(0.3), value includes **						
* $^{131}\text{In}$	D: gs and $^{131}\text{In}^m$ **						
* $^{131}\text{In}$	T: average 19Du12=265(8) 15Lo04=261(3) **						
* $^{131}\text{In}$	J: from $\beta^-$ decay properties in 04Fo06 and 19Du12 **						
* $^{131}\text{In}^m$	T: from 19Du12 **						
* $^{131}\text{In}^m$	J: from $\beta^-$ decay properties in 04Fo06 and 19Du12 **						
* $^{131}\text{In}^m$	D: % $\beta^-n$ average 93Ru01=2.2(0.3) 19Du12 $\beta^-n$ =2.3(0.3), value includes **						
* $^{131}\text{In}^m$	D: gs and $^{131}\text{In}^m$ **						
* $^{131}\text{In}^n$	T: average 19Du12=323(55) 84Fo19=320(60) **						
* $^{131}\text{In}^p$	T: average 12Ka36=685(+42-39) 09Go40=630(60) **						
* $^{131}\text{Sn}^m$	J: 20Yo.A=11/2 **						
* $^{131}\text{Sn}^n$	E: 4605.02(0.21) keV above $^{131}\text{Sn}^m$ **						
* $^{131}\text{Sn}^n$	T: others 12Ka36=309(+24-23) 84Fo19=300(20) **						
* $^{131}\text{Sb}^m$	T: average 19Bi04=64(3) 00Ge18=65(5) **						
* $^{131}\text{Sb}^p$	J: from 00Ge18 **						
* $^{131}\text{Xe}^m$	J: 90NeZY=11/2 **						
* $^{131}\text{Pr}$	T: average 96Gi08=1.57(0.07) 93Al03=1.48(0.02) 83Ga.A=1.58(0.05) **						
$^{132}\text{Ag}$	-34400#	500#	30 ms 14	$6^-$ #	15 15Lo04 TD	2015	$\beta^-$ =100; $\beta^-n$ ?; $\beta^-2n$ ?
$^{132}\text{Cd}$	-50470	60	84 ms 5	$0^+$	18 00Ha55 TD	2000	$\beta^-$ =100; $\beta^-n$ =60 15; $\beta^-2n$ ?
$^{132}\text{In}$	-62410	60	202.2 ms 0.2	$(7^-)$	18 20Be16 TD	1973	$\beta^-$ =100; $\beta^-n$ =12.3 4; $\beta^-2n$ ?
$^{132}\text{Sn}$	-76546.6	2.0	39.7 s 0.8	$0^+$	18	1963	$\beta^-$ =100
$^{132}\text{Sn}^m$	-71698.1	2.0	2.080 $\mu\text{s}$ 0.016	$8^+$	18 17Ch51 T	1986	IT=100 *
$^{132}\text{Sb}$	-79635.3	2.5	2.79 m 0.07	$(4)^+$	18	1956	$\beta^-$ =100
$^{132}\text{Sb}^m$	-79490#	50#	4.10 m 0.05	$(8^-)$	18 89St06 E	1956	$\beta^-$ =100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{132}\text{Sb}^n$	-79380.8	2.5	254.5 0.3	102 ns 4	(6 <sup>-</sup> )	18	1974 IT=100
$^{132}\text{Te}$	-85188	3		3.204 d 0.013	0 <sup>+</sup>	05	1948 $\beta^-$ =100
$^{132}\text{Te}^m$	-83413	3	1774.80 0.09	145 ns 8	6 <sup>+</sup>	05	1973 IT=100
$^{132}\text{Te}^n$	-83263	3	1925.47 0.09	28.5 $\mu\text{s}$ 0.9	7 <sup>-</sup>	05 17Ki09 T	1979 IT=100 *
$^{132}\text{Te}^p$	-82465	3	2723.3 0.8	3.62 $\mu\text{s}$ 0.06	(10 <sup>+</sup> )	05 17Ki09 T	1979 IT=100 *
$^{132}\text{I}$	-85704	4		2.295 h 0.013	4 <sup>+</sup> *	05	1948 $\beta^-$ =100
$^{132}\text{I}^m$	-85594	10	110 11 BD	1.387 h 0.015	(8 <sup>-</sup> )	05	1973 IT=86 2; $\beta^-$ =14 2
$^{132}\text{Xe}$	-89278.975	0.005		STABLE	0 <sup>+</sup>	05	1920 IS=26.909 55
$^{132}\text{Xe}^m$	-86526.77	0.17	2752.21 0.17	8.39 ms 0.11	(10 <sup>+</sup> )	05	1976 IT=100
$^{132}\text{Cs}$	-87152.7	1.0		6.480 d 0.006	2 <sup>+</sup> *	05	1953 $\beta^+$ =98.13 9; $\beta^-$ =1.87 9
$^{132}\text{Ba}$	-88434.9	1.1		STABLE >300Ey	0 <sup>+</sup>	05 96Ba24 T	1936 IS=0.10 1; $2\beta^+$ ?
$^{132}\text{La}$	-83720	40		4.59 h 0.04	2 <sup>-</sup> *	05 18Ab02 T	1951 $\beta^+$ =100
$^{132}\text{La}^m$	-83530	40	188.20 0.11	24.3 m 0.5	6 <sup>-</sup> *	05	1969 IT=76; $\beta^+$ =24
$^{132}\text{Ce}$	-82469	20		3.51 h 0.11	0 <sup>+</sup>	05	1960 $\beta^+$ =100
$^{132}\text{Ce}^m$	-80128	20	2341.15 0.21	9.4 ms 0.3	8 <sup>-</sup>	05 09Pe31 J	1969 IT=100
$^{132}\text{Pr}$	-75227	29		1.49 m 0.11	(2 <sup>+</sup> )	05 94Bu18 TJ	1974 $\beta^+$ =100 *
$^{132}\text{Pr}^m$	-75200#	40#	30# 30# *	1# s	(5 <sup>+</sup> )	05 90Ko25 J	1990 $\beta^+$ ?;IT ?
$^{132}\text{Pr}^n$	-74980#	40#	250# 30#	2.46 $\mu\text{s}$ 0.04	(8 <sup>+</sup> )	12Ta18 TJD	2012 IT=100 *
$^{132}\text{Pr}^p$	-74930#	40#	300# 30#	486 ns 70	(8 <sup>-</sup> )	12Ta18 TJD	2012 IT=100 *
$^{132}\text{Nd}$	-71426	24		1.56 m 0.10	0 <sup>+</sup>	05 95Bu11 T	1977 $\beta^+$ =100 *
$^{132}\text{Pm}$	-61630#	150#		6.2 s 0.6	(3 <sup>+</sup> )	05	1977 $\beta^+$ =100; $\beta^+$ p $\approx$ 5e-5
$^{132}\text{Sm}$	-55140#	300#		4.0 s 0.3	0 <sup>+</sup>	05	1989 $\beta^+$ =100; $\beta^+$ p ?
$^{132}\text{Eu}$	-42200#	400#		100# ms	1 <sup>+</sup> #	05 93Li40 D	$\beta^+$ ?; $\beta^+$ p ?;p=0
* $^{132}\text{Ag}$	T : symmetrized from $^{15}\text{Lo}04=28(+15-12)$ **						
* $^{132}\text{Cd}$	T : average $^{15}\text{Lo}04=82(4)$ $^{00}\text{Ha}55,^{01}\text{Ha}39=97(10)$ **						
* $^{132}\text{In}$	T : others: $^{20}\text{Wh}02=194(5)$ $^{15}\text{Lo}04=198(2)$ $^{02}\text{Di}12=206(6)$ $^{93}\text{Ru}01=221(11)$ **						
* $^{132}\text{In}$	T : $^{86}\text{Wa}17=204(6)$ , $^{85}\text{Re}.A=204(6)$ (same as $^{86}\text{Wa}17$ ) $^{86}\text{Bj}01=186(22)$ **						
* $^{132}\text{In}$	D : % $\beta^-$ -n average $^{20}\text{Wh}02=12.3(0.4)$ $^{20}\text{Be}16=12(2)$ $^{93}\text{Ru}01=10.7(3.3)$ ; others: **						
* $^{132}\text{In}$	D : $^{86}\text{Re}ZU=6.8(1.4)$ $^{80}\text{Lu}04=4.2(0.9)$ **						
* $^{132}\text{Sn}^m$	T : average $^{17}\text{Ch}51=2.15(0.16)$ $^{12}\text{Ka}36=2.088(0.017)$ $^{94}\text{Fo}14=2.03(4)$ ; **						
* $^{132}\text{Sn}^m$	T : other $^{82}\text{Ka}25=1.7(2)$ **						
* $^{132}\text{Te}^n$	T : average $^{17}\text{Ki}09=28.6(+1.2-1.1)$ $^{79}\text{Si}08=28.1(1.5)$ **						
* $^{132}\text{Te}^n$	J : E1 to 6+ **						
* $^{132}\text{Te}^p$	T : average $^{17}\text{Ki}09=3.52(0.09)$ $^{01}\text{Ge}07=3.70(0.09)$ $^{79}\text{Si}08=3.9(0.3)$ **						
* $^{132}\text{Pr}$	T : average $^{94}\text{Bu}18=1.47(0.12)$ $^{74}\text{Ar}27=1.6(0.3)$ **						
* $^{132}\text{Pr}^n$	E : $^{12}\text{Ta}18=219.9(0.2)$ keV above $^{132}\text{Pr}^m$ **						
* $^{132}\text{Pr}^p$	E : $^{12}\text{Ta}18=273.0(0.6)$ keV above $^{132}\text{Pr}^m$ **						
* $^{132}\text{Nd}$	T : average $^{95}\text{Bu}11=1.47(0.12)$ $^{77}\text{Bo}02=1.75(0.17)$ **						
$^{133}\text{Ag}$	-29080#	500#			9/2 <sup>+</sup> #		$\beta^-$ ?; $\beta^-$ n ?
$^{133}\text{Cd}$	-44140#	200#		61 ms 6	7/2 <sup>-</sup> #	11 $^{15}\text{Lo}04$ T	2010 $\beta^-$ =100; $\beta^-$ n=?; $\beta^-$ 2n ? *
$^{133}\text{In}$	-57690#	200#		163.0 ms 1.6	(9/2 <sup>+</sup> )	11 $^{20}\text{Be}16$ TD	1996 $\beta^-$ =100; $\beta^-$ n=?; $\beta^-$ 2n ? *
$^{133}\text{In}^m$	-57360#	200#	330# 40#	167 ms 11	(1/2 <sup>-</sup> )	11 $^{20}\text{Be}16$ TD	1996 $\beta^-$ =100; $\beta^-$ n=93 3 *
$^{133}\text{Sn}$	-70873.9	1.9		1.37 s 0.07	7/2 <sup>-</sup> *	11 $^{93}\text{Ru}01$ TD	1973 $\beta^-$ =100; $\beta^-$ n=0.0294 24 *
$^{133}\text{Sb}$	-78924	3		2.34 m 0.05	(7/2 <sup>+</sup> )	11	1966 $\beta^-$ =100
$^{133}\text{Sb}^m$	-74383	9	4541 9	16.54 $\mu\text{s}$ 0.19	(21/2 <sup>+</sup> )	11 $^{16}\text{Bo}19$ E	1978 IT=100 *
$^{133}\text{Te}$	-82937.1	2.1		12.5 m 0.3	3/2 <sup>+</sup> #	11	1940 $\beta^-$ =100
$^{133}\text{Te}^m$	-82602.8	2.1	334.26 0.04	55.4 m 0.4	(11/2 <sup>-</sup> )	11	1957 $\beta^-$ =83.5 20;IT=16.5 20
$^{133}\text{Te}^n$	-81326.7	2.2	1610.4 0.5	100 ns 5	(19/2 <sup>-</sup> )	11	2001 IT=100
$^{133}\text{I}$	-85857	6		20.83 h 0.08	7/2 <sup>+</sup> *	11	1940 $\beta^-$ =100
$^{133}\text{I}^m$	-84223	6	1634.148 0.010	9 s 2	(19/2 <sup>-</sup> )	11	1970 IT=100
$^{133}\text{I}^n$	-84128	6	1729.137 0.010	$\sim$ 170 ns	(15/2 <sup>-</sup> )	11	1984 IT=100
$^{133}\text{I}^p$	-83363	6	2493.7 0.4	469 ns 15	(23/2 <sup>+</sup> )	11	2009 IT=100
$^{133}\text{Xe}$	-87643.6	2.4		5.2474 d 0.0005	3/2 <sup>+</sup> *	11 FGK209 T	1940 $\beta^-$ =100
$^{133}\text{Xe}^m$	-87410.4	2.4	233.221 0.015	2.198 d 0.013	11/2 <sup>-</sup> *	11 $^{90}\text{Ne}Z\text{Y}$ J	1951 IT=100 *
$^{133}\text{Xe}^n$	-85497#	20#	2147# 20#	8.64 ms 0.13	(23/2 <sup>+</sup> )	11 $^{17}\text{Vo}06$ EJ	2017 IT=100 *
$^{133}\text{Cs}$	-88070.943	0.008		STABLE	7/2 <sup>+</sup> *	11	1921 IS=100
$^{133}\text{Ba}$	-87553.5	1.0		10.5379 y 0.0016	1/2 <sup>+</sup> *	11 FGK209 T	1941 $\epsilon$ =100
$^{133}\text{Ba}^m$	-87265.2	1.0	288.252 0.009	38.90 h 0.06	11/2 <sup>-</sup> *	11 $^{12}\text{Da}04$ T	1941 IT=99.9896 4; $\epsilon$ =0.0104 5 *
$^{133}\text{La}$	-85494	28		3.912 h 0.008	5/2 <sup>+</sup> *	11	1950 $\beta^+$ =100
$^{133}\text{Ce}$	-82418	16		97 m 4	1/2 <sup>+</sup> *	11	1951 $\beta^+$ =100
$^{133}\text{Ce}^m$	-82381	16	37.2 0.7	5.1 h 0.3	9/2 <sup>-</sup> *	11	1951 $\beta^+$ =100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{133}\text{Pr}$	-77938	12		6.5 m 0.3	$5/2^+*$	11	1970	$\beta^+=100$	
$^{133}\text{Pr}^m$	-77746	12	192.12 0.14	1.1 s 0.2	$(11/2^-)$	11	1995	IT=100	
$^{133}\text{Nd}$	-72330	50		70 s 10	$(7/2^+)$	11	1977	$\beta^+=100$	
$^{133}\text{Nd}^m$	-72200	50	127.97 0.12	~70 s	$(1/2)^+$	11 95Br24 DT	1993	$\beta^+=?;IT=?$	
$^{133}\text{Nd}^n$	-72150	50	176.10 0.10	301 ns 18	$(9/2^-)$	11	1993	IT=100	
$^{133}\text{Pm}$	-65410	50		13.5 s 2.1	$(3/2^+)$	11	1977	$\beta^+=100$	
$^{133}\text{Pm}^m$	-65280	50	129.7 0.7	8# s	$(11/2^-)$	11	1996	$\beta^+ ?;IT ?$	
$^{133}\text{Sm}$	-57230#	300#		2.89 s 0.16	$(5/2^+)$	11	1977	$\beta^+=100;\beta^+p=?$	
$^{133}\text{Sm}^m$	-57110#	310#	120# 60#	3.5 s 0.4	$(1/2^-)$	11	1993	$\beta^+=?;IT ?;\beta^+p ?$	
$^{133}\text{Eu}$	-47240#	300#		200# ms	$5/2^+ \#$			$\beta^+ ?;\beta^+p ?$	
$^{133}\text{Gd}$	-36060#	500#		10# ms	$5/2^+ \#$			$\beta^+ ?;\beta^+p ?$	
$^{133}\text{Cd}$	T : average 15Lo04=64(8) 05Kr20=57(10)								
$^{133}\text{Cd}$	D : $\beta^-n$ was observed in 05Kr20, but it was not quantified								
$^{133}\text{In}$	T : average 19Pi04,20Be16=162(2) 15Lo04=163(7) 02Di12=165(3);								
$^{133}\text{In}$	T : other: 96Ho16=180(15)								
$^{133}\text{In}$	D : % $\beta^-n$ from 20Be16, supersedes 19Pi04=74(5) (same experiment);								
$^{133}\text{In}$	D : other: 96Ho16=85(10)								
$^{133}\text{In}^m$	T : from 19Pi04,20Be16=167(11)								
$^{133}\text{In}^n$	D : % $\beta^-n$ from 20Be16, supersedes 19Pi04=80(5) (same experiment);								
$^{133}\text{Sn}$	T : average 06KeZZ=1.57(0.14) 93Ru01=1.20(0.05) 78Si05=1.37(0.07)								
$^{133}\text{Sn}$	T : 73Bo42=1.47(0.04), Birge ratio=2.57								
$^{133}\text{Sn}$	J : 20Ro19=7/2								
$^{133}\text{Sb}^m$	E : from 4526+x keV above gs, with x<30 keV in 16Bo19								
$^{133}\text{Xe}^m$	J : 90NeZY=11/2								
$^{133}\text{Xe}^n$	E : from 2107+x keV in 17Vo06; x=40#(20#) estimated by Nubase								
$^{133}\text{Xe}^n$	T : from 18Ka47								
$^{133}\text{Ba}^m$	T : average 12Da04=38.88(0.08) 11Gr01=38.92(0.09)								
$^{134}\text{Cd}$	-39460#	300#		65 ms 15	$0^+$	15	2015	$\beta^-=100;\beta^-n ?;\beta^-2n ?$	
$^{134}\text{In}$	-51970#	200#		136 ms 4	$7^- \#$	04 15Lo04 T	1996	$\beta^-=100;\beta^-n \approx 65;\beta^-2n < 4$	*
$^{134}\text{In}^m$	-51910#	200#	56.7 0.1	3.5 $\mu\text{s}$ 0.4	$(5^-)$	19Ph02 ETJ	2019	IT=100	
$^{134}\text{Sn}$	-66434	3		0.93 s 0.08	$0^+$	04 75As04 TD	1974	$\beta^-=100;\beta^-n=17$ 13	*
$^{134}\text{Sn}^m$	-65187	3	1247.4 0.5	87 ns 8	$6^+$	04 12Ka36 T	2000	IT=100	*
$^{134}\text{Sb}$	-74019	3		674 ms 4	$(0^-)$	11 18Si28 T	1967	$\beta^-=100;\beta^-n ?$	
$^{134}\text{Sb}^m$	-73740	3	279 1	10.01 s 0.04	$(7^-)$	11 18Si28 T	1968	$\beta^-=100;\beta^-n=0.088$ 4	*
$^{134}\text{Te}$	-82533.8	2.7		41.8 m 0.8	$0^+$	04	1948	$\beta^-=100$	
$^{134}\text{Te}^m$	-80842.5	2.7	1691.34 0.16	164.5 ns 0.7	$6^+$	04 17Ur03 T	1970	IT=100	*
$^{134}\text{I}$	-84043	5		52.5 m 0.2	$(4)^+$	04	1948	$\beta^+=100$	
$^{134}\text{I}^m$	-83727	5	316.49 0.22	3.52 m 0.04	$(8)^-$	04	1970	IT=97.7 10; $\beta^-=2.3$ 10	
$^{134}\text{Xe}$	-88125.834	0.006		STABLE >11Py	$0^+$	04 89Ba22 T	1920	IS=10.436 35; $2\beta^- ?$	*
$^{134}\text{Xe}^m$	-86160.3	0.5	1965.5 0.5	290 ms 17	$7^-$	04	1968	IT=100	
$^{134}\text{Xe}^n$	-85100.6	1.5	3025.2 1.5	5 $\mu\text{s}$ 1	$(10)^+$	04	2001	IT=100	
$^{134}\text{Cs}$	-86891.165	0.016		2.0650 y 0.0004	$4^+*$	04 FGK209 T	1940	$\beta^-=100;\epsilon=0.00030$ 12	*
$^{134}\text{Cs}^m$	-86752.421	0.016	138.7441 0.0026	2.912 h 0.002	$8^-*$	04	1975	IT=100	
$^{134}\text{Ba}$	-88950.00	0.25		STABLE	$0^+$	04	1936	IS=2.42 15	
$^{134}\text{Ba}^m$	-85992.8	0.6	2957.2 0.5	2.61 $\mu\text{s}$ 0.13	$10^+$	04 19Ka36 JT	1982	IT=100	*
$^{134}\text{La}$	-85219	20		6.45 m 0.16	$1^+$	04	1951	$\beta^+=100$	
$^{134}\text{La}^m$	-84780#	100#	440# 100#	29 $\mu\text{s}$ 4	$(6^-)$	04	1985	IT=100	*
$^{134}\text{Ce}$	-84833	20		3.16 d 0.04	$0^+$	04	1951	$\epsilon=100$	
$^{134}\text{Ce}^m$	-81624	20	3208.6 0.4	308 ns 5	$10^+$	04	1980	IT=100	
$^{134}\text{Pr}$	-78528	20		17 m 2	$2^-*$	04	1967	$\beta^+=100$	
$^{134}\text{Pr}^m$	-78460	20	67.7 0.4	~11 m	$6^-$	04 11Ti10 EJ	1973	$\beta^+=100$	*
$^{134}\text{Nd}$	-75646	12		8.5 m 1.5	$0^+$	04	1970	$\beta^+=100$	
$^{134}\text{Nd}^m$	-73353	12	2293.0 0.4	389 $\mu\text{s}$ 17	$8^-$	04 17Pe03 TJ	1969	IT=100	*
$^{134}\text{Pm}$	-66760	40		22 s 1	$(5^+)$	04	1977	$\beta^+=100$	
$^{134}\text{Pm}^m$	-66710#	60#	50# 50#	~5 s	$(2^+)$	04	1988	$\beta^+=100$	
$^{134}\text{Pm}^n$	-66640#	60#	120# 50#	20 $\mu\text{s}$ 1	$(7^-)$	09Cu02 TJ	2009	IT=100	*
$^{134}\text{Sm}$	-61380#	200#		9.5 s 0.8	$0^+$	04	1977	$\beta^+=100$	
$^{134}\text{Eu}$	-49800#	300#		500 ms 200		04	1989	$\beta^+=100;\beta^+p=?$	
$^{134}\text{Gd}$	-41530#	400#		400# ms	$0^+$	04		$\beta^+ ?;\beta^+p ?$	
$^{134}\text{In}$	T : average 15Lo04=126(7) 02Di12=141(5) 96Ho16=138(8)								
$^{134}\text{In}$	D : % $\beta^-n$ from 96Ho16; % $\beta^-2n$ intensity limit from 95Jo.A								

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
* <sup>134</sup> Sn	T : unweighted average 93Ru01=1.050(0.011) 75As04=0.7(0.2) 15Lo04=0.89(0.2)						**	
* <sup>134</sup> Sn	T : 20Wu04=1.07(0.27); Birge ratio=4.15; others 90Fo03=1.2(0.1)						**	
* <sup>134</sup> Sn	T : 76Lu02=1.04(0.02) 02Pf04=1.12(0.08), compilation						**	
* <sup>134</sup> Sn <sup>m</sup>	T : symmetrized from 12Ka36=86(+8-7); other 00Ko15=80(15)						**	
* <sup>134</sup> Sb <sup>m</sup>	T : average 18Si28=9.87(0.08) 93Ru01=10.07(0.05)						**	
* <sup>134</sup> Te <sup>m</sup>	T : average 17Ur03=165.1(1.0) 01Mi22=165(6) 95Om01=164(1) 70Jo20=163(7)						**	
* <sup>134</sup> Te <sup>m</sup>	T : 74CIZX=163(4) 74Su04=170(4) 76ChZD=161(4); others 17Ki09=190(10)						**	
* <sup>134</sup> Te <sup>m</sup>	T : 74B103=196(7) 04Hw02=197(20)						**	
* <sup>134</sup> Xe	T : others 0nu-BB 89Ba22>58Zy and >26Zy for 0+>0+ and 0+>2+ respectively						**	
* <sup>134</sup> Cs	D : %E from 75Va12						**	
* <sup>134</sup> Ba <sup>m</sup>	T : average 19Ka36=2.51(0.30) 82BeZY=2.63(0.14)						**	
* <sup>134</sup> La <sup>m</sup>	E : from 336.44(17)+x keV; x=100#(100#) keV estimated by Nubase						**	
* <sup>134</sup> Pr <sup>m</sup>	E : from a least-squares fit to the level scheme of 11Ti10						**	
* <sup>134</sup> Nd <sup>m</sup>	T : average 17Pe03=380(20) 72Pa26=410(30)						**	
* <sup>134</sup> Pm <sup>n</sup>	E : 70.7(0.2) keV above a 6+ state that decays via a low-energy gamma to 5+						**	
<sup>135</sup> Cd	-32820#	400#		5/2 <sup>-</sup> #			$\beta^- ?; \beta^- n ?$	
<sup>135</sup> In	-47110#	300#	103 ms 3	9/2 <sup>+</sup> #	16 20PhZZ TD	2002	$\beta^- =100; \beta^- n = ?; \beta^- 2n ?$	
<sup>135</sup> Sn	-60632	3	515 ms 5	7/2 <sup>-</sup> #	16 15Lo04 T	1994	$\beta^- =100; \beta^- n =21 3; \beta^- 2n ?$	
<sup>135</sup> Sb	-69690.3	2.6	1.668 s 0.009	(7/2 <sup>+</sup> )	16 20Wa04 D	1964	$\beta^- =100; \beta^- n =19.1 5$	
<sup>135</sup> Te	-77728.8	1.7	19.0 s 0.2	(7/2 <sup>-</sup> )	08	1969	$\beta^- =100$	
<sup>135</sup> Te <sup>m</sup>	-76173.9	1.7	1554.89 0.16	511 ns 20	(19/2 <sup>-</sup> )	08	IT=100	
<sup>135</sup> I	-83779.2	2.1	6.58 h 0.03	7/2 <sup>+</sup> *	08	1940	$\beta^- =100$	
<sup>135</sup> Xe	-86413	4	9.14 h 0.02	3/2 <sup>+</sup>	08	1940	$\beta^- =100$	
<sup>135</sup> Xe <sup>m</sup>	-85886	4	526.551 0.013	15.29 m 0.05	11/2 <sup>-</sup> *	08 90NeZY J	1960	IT≈100; $\beta^- =0.30 17$
<sup>135</sup> Cs	-87582.0	0.4		1.33 My0.19	7/2 <sup>+</sup> *	08 16Ma05 T	1949	$\beta^- =100$
<sup>135</sup> Cs <sup>m</sup>	-85949.1	1.6	1632.9 1.5	53 m 2	19/2 <sup>-</sup> *	08	IT=100	
<sup>135</sup> Ba	-87850.65	0.25		STABLE	3/2 <sup>+</sup> *	08	IS=6.59 10	
<sup>135</sup> Ba <sup>m</sup>	-87582.43	0.25	268.218 0.020	28.11 h 0.02	11/2 <sup>-</sup>	08 12Da04 T	1948	IT=100
<sup>135</sup> Ba <sup>n</sup>	-85462.7	0.6	2388.0 0.5	1.06 ms 0.04	(23/2 <sup>+</sup> )	18Ka47 ETJ	2018	IT=100
<sup>135</sup> La	-86643	9	18.91 h 0.02	5/2 <sup>+</sup> *	08 18Ab02 T	1948	$\beta^+ =100$	
<sup>135</sup> Ce	-84616	10	17.7 h 0.3	1/2 <sup>+</sup> *	08	1948	$\beta^+ =100$	
<sup>135</sup> Ce <sup>m</sup>	-84170	10	445.81 0.21	20 s 1	(11/2 <sup>-</sup> )	08	IT=100	
<sup>135</sup> Pr	-80936	12	24 m 1	3/2 <sup>+</sup> *	08	1954	$\beta^+ =100$	
<sup>135</sup> Pr <sup>m</sup>	-80578	12	358.06 0.06	105 μs 10	(11/2 <sup>-</sup> )	08	IT=100	
<sup>135</sup> Nd	-76214	19	12.4 m 0.6	9/2 <sup>-</sup> *	08	1970	$\beta^+ =100$	
<sup>135</sup> Nd <sup>m</sup>	-76149	19	64.95 0.24	5.5 m 0.5	(1/2 <sup>+</sup> )	08	$\beta^+ \approx 100; IT ?$	
<sup>135</sup> Pm	-70060	80	49 s 3	(3/2 <sup>+</sup> , 5/2 <sup>+</sup> )	08	1975	$\beta^+ =100$	
<sup>135</sup> Pm <sup>m</sup>	-69830#	50#	240# 100#	40 s 3	(11/2 <sup>-</sup> )	08 89Ko07 TJ	1989	$\beta^+ =100$
<sup>135</sup> Sm	-62860	150	10.3 s 0.5	(7/2 <sup>+</sup> )	08 77Bo02 J	1977	$\beta^+ =100; \beta^+ p =0.02 1$	
<sup>135</sup> Eu	-54150#	200#	1.5 s 0.2	5/2 <sup>+</sup> #	16	1989	$\beta^+ =100; \beta^+ p ?$	
<sup>135</sup> Gd	-44250#	400#	1.1 s 0.2	(5/2 <sup>+</sup> )	16	1996	$\beta^+ =100; \beta^+ p \approx 2$	
<sup>135</sup> Tb	-33050#	400#	1.01 ms 0.28	(7/2 <sup>-</sup> )	16	2004	$p \approx 100; \beta^+ ?$	
* <sup>135</sup> In	T : average 20PhZZ=104(4) 15Lo04=103(5) 02Di12=92(10)						**	
* <sup>135</sup> In	D : $\beta^- n$ was observed in 02Di12 and 20PhZZ, but was not quantified						**	
* <sup>135</sup> Sb	D : % $\beta^- n$ average 20Wa04=14.6(0.4,stat)(1.2,syst) 17AgZZ=24.5(1.0)						**	
* <sup>135</sup> Sb	D : 93Ru01=21.0(1.1) 02Sh08=22(3) 78Cr03=14(1) 93Ru01=22(4),						**	
* <sup>135</sup> Sb	D : supersedes 77Ru04=19.9(2.1); Birge ratio=3.46						**	
* <sup>135</sup> Sb	T : average 93Ru01=1.662(0.010) 68To18=1.696(0.021); other (recent)						**	
* <sup>135</sup> Sb	T : 20Wu04=1.57(0.23), outweighed						**	
* <sup>135</sup> Xe <sup>m</sup>	D : % $\beta^-$ ranging from 0.004% to 0.6%						**	
* <sup>135</sup> Xe <sup>m</sup>	J : 90NeZY=11/2						**	
* <sup>135</sup> Cs	T : average 16Ma05=1.6(0.6) by AMS and 1.3(0.2) ICPMS						**	
* <sup>135</sup> Pr	J : 19Fr08,72Ek04=3/2						**	
* <sup>135</sup> Pm <sup>m</sup>	E : from TNN of 11/2 <sup>-</sup> level in Pm isotopes: <sup>133</sup> Pm: 130 keV						**	
* <sup>135</sup> Pm <sup>m</sup>	E : <sup>137</sup> Pm: 150(50) keV <sup>139</sup> Pm: 189 keV <sup>141</sup> Pm: 629 keV						**	
* <sup>135</sup> Tb	T : symmetrized from 04Wo07=940(+330-220) us						**	
<sup>136</sup> In	-40970#	300#	86 ms 9	7 <sup>-</sup> #	18 15Lo04 TD	2015	$\beta^- =100; \beta^- n ?; \beta^- 2n ?$	
<sup>136</sup> Sn	-56170#	200#	355 ms 18	0 <sup>+</sup>	18 20Ju02 T	1994	$\beta^- =100; \beta^- n =28 3; \beta^- 2n ?$	
<sup>136</sup> Sb	-64507	6	923 ms 14	(1 <sup>-</sup> )	18 20Wa04 D	1976	$\beta^- =100; \beta^- n =24.7 5;$	

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{136}\text{Sb}^m$	-64238	6	269.3	0.5	570 ns	5	$\beta^-$ 2n=0.14 3
$^{136}\text{Te}$	-74425.3	2.3			17.63 s	0.09	IT=100
$^{136}\text{I}$	-79545	14			83.4 s	0.4	$\beta^-$ =100; $\beta^-$ n=1.37 4
$^{136}\text{I}^m$	-79339	5	206	15	BD		$\beta^-$ =100
$^{136}\text{Xe}$	-86429.170	0.007			46.6 s	1.1	$\beta^-$ =100
$^{136}\text{Xe}^m$	-84537.43	0.07	1891.74	0.07	2.18 Zy	0.05	IS=8.857 72; $2\beta^-$ =100
$^{136}\text{Cs}$	-86338.9	1.9			2.92 $\mu$ s	0.03	IT=100
$^{136}\text{Cs}^m$	-85821.0	1.9	517.9	0.1	13.01 d	0.05	$\beta^-$ =100
$^{136}\text{Ba}$	-88887.08	0.24			17.5 s	0.2	IT=?; $\beta^-$ ?
$^{136}\text{Ba}^m$	-86856.55	0.24	2030.535	0.018	STABLE		IS=7.85 24
$^{136}\text{Ba}^m$	-85529.9	0.3	3357.19	0.25	308.4 ms	1.9	IT=100
$^{136}\text{La}$	-86040	50			91 ns	2	IT=100
$^{136}\text{La}^m$	-85780	50	259.5	0.3	9.87 m	0.03	$\beta^+$ =100
$^{136}\text{La}^n$	-83520	50	2520.6	0.4	114 ms	5	IT=100
$^{136}\text{Ce}$	-86508.5	0.3			187 ns	27	IT=100
$^{136}\text{Ce}^m$	-83413.5	0.7	3095.0	0.6	STABLE	>32Py	IS=0.186 2; $2\beta^+$ ?
$^{136}\text{Pr}$	-81340	11			1.96 $\mu$ s	0.09	IT=100
$^{136}\text{Nd}$	-79199	12			13.1 m	0.1	$\beta^+$ =100
$^{136}\text{Pm}$	-71170	70		*	50.65 m	0.33	$\beta^+$ =100
$^{136}\text{Pm}^m$	-71070	90	100	120	MD*		$\beta^+$ =100
$^{136}\text{Pm}^n$	-71130	70	42.7	0.2	90 s	35	$\beta^+$ =100
$^{136}\text{Sm}$	-66811	12			1.5 $\mu$ s	0.1	IT=100
$^{136}\text{Sm}^m$	-64546	12	2264.7	1.1	47 s	2	$\beta^+$ =100
$^{136}\text{Eu}$	-56240#	200#		*	15 $\mu$ s	1	IT=100
$^{136}\text{Eu}^m$	-56140#	220#	100#	100#	*		$\beta^+$ =100; $\beta^+$ p $\approx$ 0.09
$^{136}\text{Gd}$	-49090#	300#			3.3 s	0.3	$\beta^+$ =100; $\beta^+$ p $\approx$ 0.09
$^{136}\text{Tb}$	-35900#	500#			1# s	>200ns	$\beta^+$ ?; $\beta^+$ p ?
$^{136}\text{In}$	T : symmetrized from 15Lo04=85(+10-8)						
$^{136}\text{Sn}$	D : % $\beta^-$ -n average 11Ar18=27(4) 02Sh08=30(5)						
$^{136}\text{Sn}$	T : average 20Ju02=361(5) (supersedes 15Lo04=350(5)) 11Ar18=300(15);						
$^{136}\text{Sb}$	T : Birge ratio=3.86						
$^{136}\text{Sb}$	D : % $\beta^-$ -n average 20Wa04=17.6(1.0,stat)(2.7,syst) 18Ca22=32.2(1.5)						
$^{136}\text{Sb}$	D : 15CaZM=19.2(1.8) 93Ru01=16.3(3.2); Birge ratio=3.97; % $\beta^-$ 2n from 18Ca22						
$^{136}\text{Sb}^m$	T : others 15Lo08=489(40) 07Si27=480(100) 01Mi22=570(50)						
$^{136}\text{Te}$	D : % $\beta^-$ -n average 18Ca22=1.47(0.06) 93Ru01=1.31(0.05) 12Ma63=1.34(0.13)						
$^{136}\text{Xe}$	T : value for 2v- $\beta\beta$ ; other 19Ga11=2.23(0.08) 14Al03=2.165(0.061)						
$^{136}\text{Xe}$	T : 12Ga17=2.38(0.14) 15Ba11=2.19(0.06) (evaluation); Onu-BB: 18Al05>18 Yy						
$^{136}\text{Xe}$	T : 16As01>2.5 Yy 13Ga07>19 Yy 12Au03>16 Yy 02Be74>10Zy (all at 90% C.L.)						
$^{136}\text{Xe}^m$	T : average 17Ki09=2.92(0.03) 70Jo20=2.80(0.37), 3.40(0.35)						
$^{136}\text{Xe}^m$	T : 74ClZX=2.78(0.17), 3.35(0.47) 70Ca25=2.8(0.2),3.0(0.3)						
$^{136}\text{Xe}^m$	T : 70Gr38=3.10(0.38) 69Wa29=3.4(0.4)						
$^{136}\text{Cs}$	J : 81Th06,76Fu06,71Da01=5						
$^{136}\text{Cs}^m$	J : 81Th06=8						
$^{136}\text{Cs}^m$	E : also 83We07=518(5)						
$^{136}\text{La}$	J : 76Fu06,73In04=1						
$^{136}\text{Ce}$	T : for 2K cature-2nu; see also 17Be21						
$^{136}\text{Ce}^m$	T : average 13Va10=1.9(0.1) 75Yo01=2.2(0.2)						
$^{136}\text{Pr}$	J : 19Fr08,76Fu06,72Ek04=2						
$^{136}\text{Pm}$	J : expected conf=p5/2[532] n9/2[514],K=7+ (prolate shape); supported by the						
$^{136}\text{Pm}$	J : observed direct feeding to I=6,7 levels following $^{136}\text{Pm}$ $\beta^+$ decay						
$^{136}\text{Pm}^n$	T : from 30 s < T1/2 <150 s in 89Vi04; other 88Ke03=300(50)s, but according						
$^{136}\text{Pm}^m$	T : to the authors the value is poorly defined						
$^{136}\text{Pm}^m$	J : expected p5/2[532] n9/2[514],K=2+ (prolate shape); supported by the						
$^{136}\text{Pm}^m$	J : observed direct feeding to I=2,3 levels following $^{136}\text{Pm}$ $\beta^+$ decay						
$^{136}\text{Pm}^n$	E : 08Ri05=42.7(0.2) keV above a long lived state that could be either the						
$^{136}\text{Pm}^n$	E : gs or the isomer						
$^{136}\text{Pm}^n$	J : expected conf=p5/2[413] n9/2[514],K=7- (prolate shape); 42.7g E1,						
$^{136}\text{Pm}^n$	J : consistent with the measured half-life						
$^{136}\text{Eu}$	J : expected conf=p5/2[413] n7/2[404],K=6+ (prolate shape)						
$^{136}\text{Eu}$	J : expected conf=p5/2[413] n7/2[404],K=1+ (prolate shape)						

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{137}\text{In}$	-35830#	400#	70 ms	40	9/2 <sup>+</sup> #	15 15Lo04 TD	2015 $\beta^- = 100; \beta^- n ?; \beta^- 2n ?$ *	
$^{137}\text{Sn}$	-50150#	300#	249 ms	15	5/2 <sup>-</sup> #	07 11Ar18 TD	1994 $\beta^- = 100; \beta^- n = 48.6; \beta^- 2n ?$ *	
$^{137}\text{Sb}$	-60060	50	497 ms	21	7/2 <sup>+</sup> #	07 20Wu04 T	1994 $\beta^- = 100; \beta^- n = 49.6; \beta^- 2n ?$ *	
$^{137}\text{Te}$	-69303.8	2.1	2.49 s	0.05	3/2 <sup>-</sup> #	07 17AgZZ D	1975 $\beta^- = 100; \beta^- n = 2.94.14$ *	
$^{137}\text{I}$	-76356	8	24.13 s	0.12	7/2 <sup>+</sup> #	07 20Cz01 D	1943 $\beta^- = 100; \beta^- n = 7.51.11$ *	
$^{137}\text{Xe}$	-82383.41	0.10	3.818 m	0.013	7/2 <sup>-</sup> *	07 89Bo03 J	1943 $\beta^- = 100$ *	
$^{137}\text{Cs}$	-86545.8	0.3	30.04 y	0.04	7/2 <sup>+</sup> *	07 FGK204 T	1951 $\beta^- = 100$ *	
$^{137}\text{Ba}$	-87721.40	0.25	STABLE		3/2 <sup>+</sup> *	07	1932 IS=11.23.23	
$^{137}\text{Ba}^m$	-87059.74	0.25	661.659	0.003	2.552 m	0.001	11/2 <sup>-</sup> *	07 1965 IT=100
$^{137}\text{Ba}^n$	-85372.3	0.6	2349.1	0.5	589 ns	20	(19/2 <sup>-</sup> )	07 17Vo01 T 1973 IT=100 *
$^{137}\text{La}$	-87140.9	1.6	60 ky	20	7/2 <sup>+</sup> *	07	1948 $\epsilon = 100$	
$^{137}\text{La}^m$	-85271.4	1.6	1869.50	0.21	342 ns	25	19/2 <sup>-</sup>	07 1982 IT=100
$^{137}\text{Ce}$	-85918.8	0.4	9.0 h	0.3	3/2 <sup>+</sup> *	07	1948 $\beta^+ = 100$	
$^{137}\text{Ce}^m$	-85664.5	0.4	254.29	0.05	34.4 h	0.3	11/2 <sup>-</sup> *	07 1958 IT=99.21.4; $\beta^+ = 0.79.4$
$^{137}\text{Pr}$	-83202	8	1.28 h	0.03	5/2 <sup>+</sup> *	07	1958 $\beta^+ = 100$ *	
$^{137}\text{Pr}^m$	-82641	8	561.22	0.23	2.66 $\mu\text{s}$	0.07	11/2 <sup>-</sup>	07 1987 IT=100
$^{137}\text{Nd}$	-79584	12	38.5 m	1.5	1/2 <sup>+</sup> *	07	1970 $\beta^+ = 100$	
$^{137}\text{Nd}^m$	-79065	12	519.43	0.20	1.60 s	0.15	11/2 <sup>-</sup>	07 1970 IT=100
$^{137}\text{Pm}$	-74073	13	&		2# m		5/2 <sup>-</sup> #	1975 $\beta^+ ?$
$^{137}\text{Pm}^m$	-73910	40	160	50	BD &		2.4 m	0.1 11/2 <sup>-</sup> 07 1973 $\beta^+ = 100$
$^{137}\text{Sm}$	-67992	29	45 s	1	(9/2 <sup>-</sup> )	07	1986 $\beta^+ = 100$	
$^{137}\text{Sm}^m$	-67890#	60#	100#	50#	20# s		1/2 <sup>+</sup> #	$\beta^+ ?$
$^{137}\text{Eu}$	-60146	4	8.4 s	0.5	5/2 <sup>+</sup> #	07 88Be.A T	1982 $\beta^+ = 100$	
$^{137}\text{Gd}$	-51210#	300#	2.2 s	0.2	(7/2)( <sup>+</sup> #)	07	1999 $\beta^+ = 100; \beta^+ p = ?$	
$^{137}\text{Tb}$	-40970#	400#	600# ms		3/2 <sup>+</sup> #		$p ?; \beta^+ ?$	
* $^{137}\text{In}$	T : symmetrized from 15Lo04=65(+40-30)							**
* $^{137}\text{Sn}$	T : average 11Ar18=273(7) 20Wu04=204(12) 20Ju02t=238(8), supersedes							**
* $^{137}\text{Sn}$	T : 15Lo04=230(30), 02Sh08=190(60); Birge ratio=3.09							**
* $^{137}\text{Sn}$	D : % $\beta^- n$ average 11Ar18=50(8) 02Sh08=58(15) 20Ju02d=32(15)							**
* $^{137}\text{Sb}$	T : average 20Wu04=566(52) 11Ar18=492(25) 02Sh08=450(50)							**
* $^{137}\text{Sb}$	D : % $\beta^- n$ average 11Ar18=49(8)% 02Sh08=49(10)%							**
* $^{137}\text{Te}$	J : TNN of N=85 isotones. Ensdf2007=(7/2-) from shell-model prediction							**
* $^{137}\text{Te}$	D : % $\beta^- n$ average 17AgZZ=2.6(0.3) 93Ru01=3.04(0.16)							**
* $^{137}\text{I}$	T : from 93Ru01, supersedes 74Ru08=24.5(0.2) (same group)							**
* $^{137}\text{I}$	D : % $\beta^- n$ average 20Cz01=6.66(0.34) 17Ra10=7.9(0.5) 16Ag03=7.76(0.14)							**
* $^{137}\text{I}$	D : 93Ru01=7.14(0.23)							**
* $^{137}\text{Xe}$	J : 89Bo03,90NeZY=7/2							**
* $^{137}\text{Ba}^n$	J : from 19Ka04							**
* $^{137}\text{Pr}$	J : 19Fr08,72Ek04=3/2							**
$^{138}\text{Sn}$	-45510#	400#	148 ms	9	0 <sup>+</sup>	17 20Wu04 T	2010 $\beta^- = 100; \beta^- n = 36.12; \beta^- 2n ?$ *	
$^{138}\text{Sn}^m$	-44170#	400#	210 ns	45	(6 <sup>+</sup> )	17	2014 IT=100	
$^{138}\text{Sb}$	-54650#	300#	333 ms	7	(3 <sup>-</sup> )	17 20Wu04 T	1994 $\beta^- = 100; \beta^- n = 72.8; \beta^- 2n ?$ *	
$^{138}\text{Te}$	-65696	4	1.46 s	0.25	0 <sup>+</sup>	17 20Wu04 T	1975 $\beta^- = 100; \beta^- n = 4.80.23$ *	
$^{138}\text{I}$	-71980	6	6.26 s	0.03	(1 <sup>-</sup> )	17 20Cz01 D	1949 $\beta^- = 100; \beta^- n = 5.33.11$ *	
$^{138}\text{I}^m$	-71912	6	1.26 $\mu\text{s}$	0.16	(3 <sup>-</sup> )	17	2007 IT=100	
$^{138}\text{Xe}$	-79972.2	2.8	14.14 m	0.07	0 <sup>+</sup>	17	1943 $\beta^- = 100$	
$^{138}\text{Cs}$	-82887	9	33.5 m	0.2	3 <sup>-</sup> *	17	1943 $\beta^- = 100$ *	
$^{138}\text{Cs}^m$	-82807	9	79.9	0.3	2.91 m	0.10	6 <sup>-</sup> *	17 1971 IT=81.3; $\beta^- = 19.3$ *
$^{138}\text{Cs}^x$	-82847	25	40	23	R = ?	<i>fsmix</i>		
$^{138}\text{Ba}$	-88261.81	0.25	STABLE		0 <sup>+</sup>	17	1925 IS=71.70.29	
$^{138}\text{Ba}^m$	-86171.27	0.25	2090.536	0.021	850 ns	100	6 <sup>+</sup>	17 1971 IT=100
$^{138}\text{La}$	-86513.4	0.4	103 Gy	1	5 <sup>+</sup> *	17	1947 IS=0.0.8881; $\beta^+ = 65.5.4$ ; $\beta^- = 34.5.4$ *	
$^{138}\text{La}^m$	-86440.8	0.4	72.57	0.03	116 ns	5	(3 <sup>+</sup> )	17 1975 IT=100
$^{138}\text{La}^n$	-85774.6	0.4	738.80	0.20	2.0 $\mu\text{s}$	0.3	7 <sup>-</sup>	17 14As02 TJ 2014 IT=100
$^{138}\text{Ce}$	-87565.9	0.5	STABLE		>44Py		0 <sup>+</sup>	17 1936 IS=0.251.2; $2\beta^+ ?$ *
$^{138}\text{Ce}^m$	-85436.6	0.5	2129.28	0.12	8.73 ms	0.20	7 <sup>-</sup>	17 1960 IT=100
$^{138}\text{Pr}$	-83129	10	1.45 m	0.05	1 <sup>+</sup>	17	1951 $\beta^+ = 100$	
$^{138}\text{Pr}^m$	-82779	16	350	19	BD		2.12 h	0.04 7 <sup>-</sup> 17 1958 $\beta^+ = 100$ *
$^{138}\text{Nd}$	-82017	12	5.04 h	0.09	0 <sup>+</sup>	17	1965 $\beta^+ = 100$	
$^{138}\text{Nd}^m$	-78843	12	3174.5	0.4	370 ns	5	10 <sup>+</sup>	17 1975 IT=100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)		
$^{138}\text{Pm}$	-74914	12		3.24 m 0.05	$3^-$ #	17	1973	$\beta^+=100$	*	
$^{138}\text{Pm}^m$		<i>non-exist</i>		10 s 2	$1^+$ #	17 83A106	IT	$\beta^+=100$	*	
$^{138}\text{Sm}$	-71498	12		3.1 m 0.2	$0^+$	17	1982	$\beta^+=100$	*	
$^{138}\text{Eu}$	-61750	28		5# s	$2^-$ #	17 FGK205	EJ	$\beta^+?$	*	
$^{138}\text{Eu}^m$	-61650#	60#	100# 50#	12.1 s 0.6	$7^-$ #	17 FGK205	EJ	$\beta^+=100$	*	
$^{138}\text{Gd}$	-55660#	200#		4.7 s 0.9	$0^+$	17	1985	$\beta^+=100$	*	
$^{138}\text{Gd}^m$	-53430#	200#	2232.6 1.1	6.2 $\mu$ s 0.2	$(8^-)$	17	1997	IT=100	*	
$^{138}\text{Tb}$	-43600#	300#		800# ms >200ms	$2^+$ #	17	1993	$\beta^+?; \beta^+p?; p=0$	*	
$^{138}\text{Dy}$	-34930#	500#		200# ms	$0^+$			$\beta^+?; \beta^+p?$	*	
$^{138}\text{Sn}$	T: average 20Wu04=158(15) 20Ju02t=142(12), supersedes 15Lo04=140(+30-20)									
$^{138}\text{Sb}$	T: average 20Wu04=326(8) 15Le14=346(19) 11Ar18=350(15)									
$^{138}\text{Te}$	D: % $\beta^-$ -n from 17AgZZ; other 75As04=6.3(2.1)									
$^{138}\text{Te}$	T: average 20Wu04=1.50(0.32) 75As04=1.4(0.4); other 06KeZZ=1.151(0.028)									
$^{138}\text{I}$	D: % $\beta^-$ -n average 20Cz01=6.07(0.34) 17AgZZ=4.98(0.18) 11Go37=5.32(0.20)									
$^{138}\text{I}$	D: $^{93}\text{Ru}01=5.56(0.22)$									
$^{138}\text{Cs}$	J: 67St22,78Sc27,79Bo01=3									
$^{138}\text{Cs}$	J: 81Th06=6									
$^{138}\text{La}$	J: 55So31,72Fi14=5									
$^{138}\text{Ce}$	T: see also 17Be21t									
$^{138}\text{Pr}^m$	J: 72Ek04=7									
$^{138}\text{Pm}$	J: expected conf=p5/2[532]n1/2[400],K=3- (deformed shape)									
$^{138}\text{Pm}^m$	I: not confirmed in 00Be42									
$^{138}\text{Eu}^m$	J: from the expected conf=p5/2[413] n9/2[514],K=2- (deformed shape);									
$^{138}\text{Eu}^m$	J: from systematics (p5/2[413] at Z=63 and n9/2[514] at N=75);									
$^{138}\text{Eu}^m$	J: Ensdf2017=(1-)									
$^{138}\text{Eu}^m$	J: from the expected conf=p5/2[413] n9/2[514],K=7- (deformed shape);									
$^{138}\text{Eu}^m$	J: from systematics (p5/2[413] at Z=63 and n9/2[514] at N=75);									
$^{138}\text{Eu}^m$	J: Ensdf17 J=(6-) as a ground state, but the proposed									
$^{138}\text{Eu}^m$	J: conf=p5/2[532] n7/2[404] violates the GM rule									
$^{139}\text{Sn}$	-39310#	400#		120 ms 38	$5/2^-$ #	16 20Wu04	T	2015	$\beta^-=100; \beta^-n?; \beta^-2n?$	*
$^{139}\text{Sb}$	-50050#	400#		182 ms 9	$7/2^+$ #	16 20Wu04	T	1994	$\beta^-=100; \beta^-n=90\ 10; \beta^-2n?$	*
$^{139}\text{Te}$	-60205	4		724 ms 81	$5/2^-$ #	16 20Wu04	TD	1994	$\beta^-=100; \beta^-n?$	*
$^{139}\text{I}$	-68471	4		2.280 s 0.011	$7/2^+$ #	16 17AgZZ	D	1949	$\beta^-=100; \beta^-n=9.74\ 24$	*
$^{139}\text{Xe}$	-75644.6	2.1		39.68 s 0.14	$3/2^-*$	16	1951	$\beta^-=100$	*	
$^{139}\text{Cs}$	-80701	3		9.27 m 0.05	$7/2^+*$	16	1939	$\beta^-=100$	*	
$^{139}\text{Ba}$	-84913.92	0.25		82.93 m 0.09	$7/2^-*$	16	1937	$\beta^-=100$	*	
$^{139}\text{La}$	-87222.4	0.6		STABLE	$7/2^+*$	16	1924	IS=99.9 1119	*	
$^{139}\text{La}^m$	-85422.0	0.7	1800.4 0.4	315 ns 35	$(17/2^+)$		2012	IT=100	*	
$^{139}\text{Ce}$	-86957.7	2.1		137.642 d 0.020	$3/2^+*$	16 FGK204	T	1948	$\epsilon=100$	*
$^{139}\text{Ce}^m$	-86203.5	2.1	754.24 0.08	57.58 s 0.32	$11/2^-$	16	1967	IT=100	*	
$^{139}\text{Pr}$	-84829	4		4.41 h 0.04	$5/2^+*$	16	1951	$\beta^+=100$	*	
$^{139}\text{Nd}$	-82017	28		29.7 m 0.5	$3/2^+*$	16	1951	$\beta^+=100$	*	
$^{139}\text{Nd}^m$	-81786	28	231.16 0.05	5.50 h 0.20	$11/2^-*$	16	1951	$\beta^+=87.0\ 10; IT=13.0\ 10$	*	
$^{139}\text{Nd}^m$	-79400	28	2616.9 0.6	276.8 ns 1.8	$23/2^+$	16	1980	IT=100	*	
$^{139}\text{Pm}$	-77501	14		4.15 m 0.05	$(5/2^+)$	16	1967	$\beta^+=100$	*	
$^{139}\text{Pm}^m$	-77312	14	188.7 0.3	180 ms 20	$(11/2)^-$	16	1975	IT $\approx$ 100; $\beta^+?$	*	
$^{139}\text{Sm}$	-72380	11		2.57 m 0.10	$1/2^+*$	16	1971	$\beta^+=100$	*	
$^{139}\text{Sm}^m$	-71923	11	457.38 0.23	10.7 s 0.6	$11/2^-$	16	1973	IT=93.7 5; $\beta^+=6.3\ 5$	*	
$^{139}\text{Eu}$	-65398	13		17.9 s 0.6	$(11/2)^-$	16	1975	$\beta^+=100$	*	
$^{139}\text{Eu}^m$	-65250	13	148.3 0.3	10 $\mu$ s 2	$(7/2^+)$	16	2011	IT=100	*	
$^{139}\text{Gd}$	-57630#	200#		5.7 s 0.3	$9/2^-$ #	16 99Xi04	T	1983	$\beta^+=100; \beta^+p=?$	*
$^{139}\text{Gd}^m$	-57380#	250#	250# 150#	4.8 s 0.9	$1/2^+$ #	16	1983	$\beta^+=100; \beta^+p=?$	*	
$^{139}\text{Tb}$	-48130#	300#		1.6 s 0.2	$5/2^-$ #	16	1999	$\beta^+=100; \beta^+p?$	*	
$^{139}\text{Dy}$	-37700#	500#		600 ms 200	$(7/2^+)$	01	1999	$\beta^+=100; \beta^+p\approx 11$	*	
$^{139}\text{Sn}$	T: average 20Wu04=114(49) 15Lo04=130(60)									
$^{139}\text{Sb}$	T: other 11Ar18=93(+14-3)									
$^{139}\text{Te}$	T: others (not trusted) 11Ar18=1600(300) 06KeZZ=598(20)									
$^{139}\text{I}$	D: % $\beta^-$ -n average 17AgZZ=9.27(0.33) 93Ru01=10.3(0.4) 81Ho07=10.0(1.1)									
$^{139}\text{I}$	D: $^{75}\text{As}04=10.2(0.9)$									
$^{139}\text{Xe}$	J: also 90NeZY=3/2									
$^{139}\text{Cs}$	J: 79Bo01,79Ek02,81Th06,87Co19=7/2									



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>139</sup> Ba	J : 83Mu12=7/2							**
* <sup>139</sup> La	J : 71Ch02=7/2							**
* <sup>139</sup> Ce	J : 73In04=3/2							**
* <sup>139</sup> Pr	J : 72Ek04=5/2							**
* <sup>139</sup> Nd	J : 72Ek04=3/2							**
* <sup>139</sup> Nd <sup>m</sup>	J : 72Ek04=11/2							**
* <sup>139</sup> Nd <sup>n</sup>	T : average 13Va10=278(2) 08Fe02=272(4)							**
* <sup>139</sup> Sm	J : 92Le09=1/2							**
* <sup>139</sup> Gd	T : average 99Xi04=5.8(0.9) 88Be.A=5.8(0.4); other 83Ni05=4.9(1.0)							**
* <sup>139</sup> Gd	T : not used since it corresponds to a mixture of gs and the isomer							**
* <sup>139</sup> Gd	D : β <sup>+</sup> p were observed in 83Ni05 and it is assumed that they are							**
* <sup>139</sup> Gd	D : associated with both the ground state and the isomer							**
* <sup>139</sup> Gd <sup>m</sup>	D : β <sup>+</sup> p were observed in 83Ni05 and it is assumed that they are							**
* <sup>139</sup> Gd <sup>n</sup>	D : associated with both the ground state and the isomer							**
<sup>140</sup> Sn	-34490# 300#			50# ms >550ns	0 <sup>+</sup>	18Sh11 I	2018	β <sup>-</sup> ?; β <sup>-</sup> n ?; β <sup>-</sup> 2n ?
<sup>140</sup> Sb	-44390# 600#			170 ms 6	(3 <sup>-</sup> )	18 20Wu04 T	2010	β <sup>-</sup> =100; β <sup>-</sup> n=23 4; β <sup>-</sup> 2n=7.6 25
<sup>140</sup> Sb <sup>m</sup>	-44060# 600#	330#	30#	41 μs 8	(6 <sup>-</sup> , 7 <sup>-</sup> )	18 16Lo01 ETJ	2016	IT=100
<sup>140</sup> Te	-56367 14			351 ms 5	0 <sup>+</sup>	18 20Wu04 TD	1994	β <sup>-</sup> =100; β <sup>-</sup> n=?
<sup>140</sup> I	-63606 12			588 ms 10	(2 <sup>-</sup> )	18 17Mo19 J	1972	β <sup>-</sup> =100; β <sup>-</sup> n=7.60 28; β <sup>-</sup> 2n ?
<sup>140</sup> Xe	-72986.5 2.3			13.60 s 0.10	0 <sup>+</sup>	18	1951	β <sup>-</sup> =100
<sup>140</sup> Cs	-77050 8			63.7 s 0.3	1 <sup>-*</sup>	18	1950	β <sup>-</sup> =100
<sup>140</sup> Cs <sup>m</sup>	-77036 8	13.931	0.021	471 ns 51	(2 <sup>-</sup> )	18	1974	IT=100
<sup>140</sup> Ba	-83268 8			12.7534 d 0.0021	0 <sup>+</sup>	18 FGK204 T	1939	β <sup>-</sup> =100
<sup>140</sup> La	-84312.1 0.6			40.289 h 0.004	3 <sup>-*</sup>	18 FGK209 T	1935	β <sup>+</sup> =100
<sup>140</sup> Ce	-88074.2 1.3			STABLE	0 <sup>+</sup>	18	1925	IS=88.449 51
<sup>140</sup> Ce <sup>m</sup>	-85966.3 1.3	2107.854	0.024	7.3 μs 1.5	6 <sup>+</sup>	18	1966	IT=100
<sup>140</sup> Pr	-84686 6			3.39 m 0.01	1 <sup>+</sup> *	18	1938	β <sup>+</sup> =100; e <sup>+</sup> =48.7 22; ε=51.3 22
<sup>140</sup> Pr <sup>m</sup>	-84558 6	127.8	0.3	350 ns 20	5 <sup>+</sup>	18	1964	IT=100
<sup>140</sup> Pr <sup>n</sup>	-83922 6	763.7	0.5	3.05 μs 0.20	(7 <sup>-</sup> )	18	1964	IT=100
<sup>140</sup> Nd	-84257 3			3.37 d 0.02	0 <sup>+</sup>	18	1949	ε=100
<sup>140</sup> Nd <sup>m</sup>	-82035 3	2221.65	0.09	600 μs 50	7 <sup>-</sup>	18	1962	IT=100
<sup>140</sup> Nd <sup>n</sup>	-76822 3	7435.1	0.4	1.22 μs 0.06	20 <sup>+</sup>	08Fe02 TJ	2008	IT=100
<sup>140</sup> Pm	-78212 24			9.2 s 0.2	1 <sup>+</sup>	18	1966	β <sup>+</sup> =100
<sup>140</sup> Pm <sup>m</sup>	-77783 13	429	28	5.95 m 0.05	8 <sup>-</sup>	18	1966	β <sup>+</sup> =100
<sup>140</sup> Sm	-75456 12			14.82 m 0.12	0 <sup>+</sup>	18	1967	β <sup>+</sup> =100
<sup>140</sup> Eu	-66990 50			1.51 s 0.02	1 <sup>+</sup>	18 91Fi03 TD	1982	β <sup>+</sup> =100; e <sup>+</sup> =95.1 7; ε=4.9 7
<sup>140</sup> Eu <sup>m</sup>	-66780 50	210	14	125 ms 2	(5 <sup>-</sup> )	18 91Fi03 TDE	1988	IT≈100; β <sup>+</sup> <1
<sup>140</sup> Eu <sup>n</sup>	-66320 50	669	14	299.8 ns 2.1	(8 <sup>+</sup> )	18	2002	IT=100
<sup>140</sup> Gd	-61782 28			15.8 s 0.4	0 <sup>+</sup>	18 91Fi03 TD	1985	β <sup>+</sup> =100; e <sup>+</sup> =67 8; ε=33 8
<sup>140</sup> Tb	-50480 800			2.29 s 0.15	(7 <sup>+</sup> )	18 91Fi03 D	1986	β <sup>+</sup> =100; ε<3; β <sup>+</sup> p=0.26 13
<sup>140</sup> Dy	-42830# 400#			700# ms	0 <sup>+</sup>	18	2002	β <sup>+</sup> ?; β <sup>+</sup> p ?
<sup>140</sup> Dy <sup>m</sup>	-40660# 400#	2166.1	0.5	7.0 μs 0.5	8 <sup>-</sup>	18 15Ko14 JE	2002	IT=100
<sup>140</sup> Ho	-29320# 500#			6 ms 3	8 <sup>+</sup> #	18	1999	p=?; β <sup>+</sup> ?; β <sup>+</sup> p ?
* <sup>140</sup> Sb	T : average 20Wu04=169(7) 17Mo12=173(12)							**
* <sup>140</sup> Sb <sup>m</sup>	E : 16Lo01=298.2+x keV; x=30#(30#) keV estimated by the authors							**
* <sup>140</sup> Te	T : average 20Wu04=360(21) 17Mo19=350(5); other 06KeZZ=334(14)							**
* <sup>140</sup> I	T : average 20Wu04=553(46) 76Lu02=590(10); others 76Ah01=860(40)							**
* <sup>140</sup> I	T : 75Kr17=870(40) 74Kr21=880(130) 70HeZH=860(140) 70WiZn=880(120)							**
* <sup>140</sup> I	D : %β <sup>-</sup> n from 17AgZZ=7.60(0.28); other (recent)20Wa04=7.6(0.9,sta)(2.7,syst)							**
* <sup>140</sup> Cs	J : 79Ek02,79Bo01=1							**
* <sup>140</sup> La	J : 76Fu06=3							**
* <sup>140</sup> Pr	T : other: 07Li71=7.3(0.4) for q=59+ (bare ion) 3.04(0.10) for q=58+							**
* <sup>140</sup> Pr	T : (H-like ion) and 3.84(0.15) for q=57+ (He-like ion)							**
* <sup>140</sup> Pr	D : %e <sup>+</sup> =42.4(2.3); %ε=57.6(2.3) for q=58+ (H-like ion) and							**
* <sup>140</sup> Pr	D : %e <sup>+</sup> =51.2(3.1); %ε=48.8(3.1) for q=57+ (He-like ion)							**
* <sup>140</sup> Pr	D : e <sup>+</sup> decay for the ground state (neutral atom) from 72Ev01							**
* <sup>140</sup> Nd <sup>n</sup>	T : average 13Va10=1.2(0.1) 08Fe02=1.23(0.07)							**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
* <sup>140</sup> Eu <sup>m</sup>	E : from 185.3+x keV and x<50 keV in 91Fi03									**	
* <sup>140</sup> Eu <sup>n</sup>	E : from 459.5(0.3) keV above <sup>140</sup> Eu <sup>m</sup>									**	
* <sup>140</sup> Ho	D : %p observed in 99Ry04									**	
<sup>141</sup> Sb	-39540#	500#			103 ms 29	7/2 <sup>+</sup> #	18	20Wu04	TD 2018	$\beta^- = 100; \beta^-_n ?; \beta^- 2n ?$	
<sup>141</sup> Te	-50670#	400#			193 ms 16	5/2 <sup>-</sup> #	14	20Wu04	TD 1994	$\beta^- = 100; \beta^-_n ?; \beta^- 2n ?$	
<sup>141</sup> I	-59927	16			420 ms 7	7/2 <sup>+</sup> #	14	20Wu04	TD 1974	$\beta^- = 100; \beta^-_n = 21.2$ 30	
<sup>141</sup> Xe	-68197.3	2.9			1.73 s 0.01	5/2 <sup>-</sup> *	14	90NeZY	J 1951	$\beta^- = 100; \beta^-_n = 0.044$ 5	
<sup>141</sup> Cs	-74477	9			24.84 s 0.16	7/2 <sup>+</sup> *	14	93Ru01	TD 1962	$\beta^- = 100; \beta^-_n = 0.0342$ 14	
<sup>141</sup> Ba	-79732	5			18.27 m 0.07	3/2 <sup>-</sup> *	14		1945	$\beta^- = 100$	
<sup>141</sup> La	-82930	4			3.92 h 0.03	(7/2 <sup>+</sup> )	14		1951	$\beta^- = 100$	
<sup>141</sup> Ce	-85431.1	1.3			32.505 d 0.010	7/2 <sup>-</sup> *	14	FGK204	T 1948	$\beta^- = 100$	
<sup>141</sup> Pr	-86014.5	1.5			STABLE	5/2 <sup>+</sup> *	14		1924	IS=100	
<sup>141</sup> Nd	-84192	3			2.49 h 0.03	3/2 <sup>+</sup> *	14		1949	$\beta^+ = 100; \epsilon = 97.28$ 16; e <sup>+</sup> =2.72 16	
<sup>141</sup> Nd <sup>m</sup>	-83435	3	756.51	0.05	62.0 s 0.8	11/2 <sup>-</sup>	14	70Ab05	D 1960	IT≈100; $\beta^+ = 0.032$ 8	
<sup>141</sup> Pm	-80523	14			20.90 m 0.05	5/2 <sup>+</sup> *	14		1952	$\beta^+ = 100$	
<sup>141</sup> Pm <sup>m</sup>	-79894	14	628.62	0.07	630 ns 20	11/2 <sup>-</sup>	14		1970	IT=100	
<sup>141</sup> Pm <sup>n</sup>	-77992	14	2530.75	0.17	> 2 $\mu$ s	(23/2 <sup>+</sup> )	14	85Ar19	TDJ 1985	IT=100	
<sup>141</sup> Sm	-75934	9			10.2 m 0.2	1/2 <sup>+</sup> *	14		1967	$\beta^+ = 100$	
<sup>141</sup> Sm <sup>m</sup>	-75758	9	175.9	0.3	22.6 m 0.2	11/2 <sup>-</sup> *	14		1967	$\beta^+ = 99.69$ 3; IT=0.31 3	
<sup>141</sup> Eu	-69926	13			40.7 s 0.7	5/2 <sup>+</sup>	14		1977	$\beta^+ = 100$	
<sup>141</sup> Eu <sup>m</sup>	-69830	13	96.45	0.07	2.7 s 0.3	11/2 <sup>-</sup>	14		1973	IT=86 3; $\beta^+ = 14$ 3	
<sup>141</sup> Gd	-63224	20			14 s 4	(1/2 <sup>+</sup> )	14		1986	$\beta^+ = 100; \beta^+_p = 0.03$ 1	
<sup>141</sup> Gd <sup>m</sup>	-62846	20	377.76	0.09	24.5 s 0.5	(11/2 <sup>-</sup> )	14		1986	$\beta^+ = 89$ 2; IT=11 2	
<sup>141</sup> Tb	-54540	110			3.5 s 0.2	(5/2 <sup>-</sup> )	14		1986	$\beta^+ = 100$	
<sup>141</sup> Tb <sup>m</sup>	-54540#	230#	0#	200#	7.9 s 0.6	11/2 <sup>-</sup> #	14	88Be.A	I 1988	$\beta^+ = 100$	
<sup>141</sup> Dy	-45380#	300#			900 ms 140	(9/2 <sup>-</sup> )	14		1984	$\beta^+ = 100; \beta^+_p = ?$	
<sup>141</sup> Ho	-34360#	400#			4.1 ms 0.1	(7/2 <sup>-</sup> )	14		1998	p≈100; $\beta^+ ?; \beta^+_p ?$	
<sup>141</sup> Ho <sup>m</sup>	-34290#	400#	66	2	7.3 $\mu$ s 0.3	(1/2 <sup>+</sup> )	14		1998	p=100	
* <sup>141</sup> I	D : % $\beta^-_n$ from 80A115									**	
* <sup>141</sup> I	T : average 20Wu04=418(8) 80A115=430(20); others 74Kr21=410(80)									**	
* <sup>141</sup> I	T : 76Lu02=480(30) 70HeZX=430(80)									**	
* <sup>141</sup> Xe	J : also 90NeZY=5/2									**	
* <sup>141</sup> Cs	T : average 93Ru01=24.34(0.12) 86Ok03=24.98(0.13) 76Ot03=24.94(0.06),									**	
* <sup>141</sup> Cs	T : Birge ratio=3.26									**	
* <sup>141</sup> Nd	D : %e <sup>+</sup> average 72Ev01=2.72(0.20) 66Gr05=2.73(0.27)									**	
* <sup>141</sup> Eu <sup>m</sup>	D : symmetrized from %IT=87(+2-4) and % $\beta^+ = 13(+4-2)$									**	
* <sup>141</sup> Gd	J : weak J $^\pi$ arguments in Ensdf2001									**	
* <sup>141</sup> Gd <sup>m</sup>	J : weak J $^\pi$ arguments in Ensdf2001									**	
<sup>142</sup> Sb	-33610#	300#			80 ms 50			20Wu04	T 2018	$\beta^- = 100; \beta^-_n ?; \beta^- 2n ?$	
<sup>142</sup> Te	-46550#	500#			147 ms 8	0 <sup>+</sup>	11	20Wu04	TD 1994	$\beta^- = 100; \beta^-_n ?; \beta^- 2n ?$	
<sup>142</sup> I	-54803	5			235 ms 11	2 <sup>-</sup> #	11	20Wu04	TD 1975	$\beta^- = 100; \beta^-_n ?; \beta^- 2n ?$	
<sup>142</sup> Xe	-65229.6	2.7			1.23 s 0.02	0 <sup>+</sup>	11		1960	$\beta^- = 100; \beta^-_n = 0.37$ 3	
<sup>142</sup> Cs	-70515	7			1.687 s 0.010	0 <sup>-</sup> *	11	93Ru01	TD 1962	$\beta^- = 100; \beta^-_n = 0.089$ 3	
<sup>142</sup> Ba	-77842	6			10.6 m 0.2	0 <sup>+</sup>	11		1959	$\beta^- = 100$	
<sup>142</sup> La	-80024	6			91.1 m 0.5	2 <sup>-</sup>	11	19Kr10	T 1953	$\beta^- = 100$	
<sup>142</sup> La <sup>m</sup>	-79878	6	145.82	0.08	870 ns 170	(4) <sup>-</sup>	11		1983	IT=100	
<sup>142</sup> Ce	-84532.9	2.4			STABLE	>2.9Ey	0 <sup>+</sup>	11	19Be29	T 1925	IS=11.114 51; $\alpha ?; 2\beta^- ?$
<sup>142</sup> Pr	-83786.4	1.5			19.12 h 0.04	2 <sup>-</sup> *	11		1935	$\beta^- \approx 100; \epsilon = 0.0164$ 8	
<sup>142</sup> Pr <sup>m</sup>	-83782.7	1.5	3.694	0.003	14.6 m 0.5	5 <sup>-</sup> *	11		1967	IT=100	
<sup>142</sup> Nd	-85950.1	1.3			STABLE		0 <sup>+</sup>	11	1924	IS=27.153 40	
<sup>142</sup> Nd <sup>m</sup>			non-exist	EU	16.5 $\mu$ s	6 <sup>+</sup>	14	87Pr09	I 1964	IT=100	
<sup>142</sup> Pm	-81142	24			40.5 s 0.5	1 <sup>+</sup>	11	91Fi03	TD 1959	$\beta^+ = 100; \epsilon^+ = 77.1$ 27; $\epsilon = 22.9$ 27	
<sup>142</sup> Pm <sup>m</sup>	-80259	24	883.17	0.16	2.0 ms 0.2	(8) <sup>-</sup>	11		1971	IT=100	
<sup>142</sup> Pm <sup>n</sup>	-78313	24	2828.7	0.6	67 $\mu$ s 5	(13 <sup>-</sup> )	11		1974	IT=100	
<sup>142</sup> Sm	-78981.9	1.9			72.49 m 0.05	0 <sup>+</sup>	11	91Fi03	TD 1959	$\beta^+ = 100; \epsilon^+ < 5$	
<sup>142</sup> Sm <sup>m</sup>	-76609.8	1.9	2372.1	0.4	170 ns 2	7 <sup>-</sup>	11		1975	IT=100	
<sup>142</sup> Sm <sup>n</sup>	-75319.7	2.0	3662.2	0.7	480 ns 60	10 <sup>+</sup>	11		1979	IT=100	



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
* <sup>143</sup> Gd	D : 39 particles detected						**	
* <sup>143</sup> Gd <sup>m</sup>	J : from 05Ba64						**	
* <sup>143</sup> Dy	T : 03Xu04=5.6(1.0); 84Ni03=3.2(0.6) 83Ni05=4.1(0.3) in diff. experiments						**	
* <sup>143</sup> Dy <sup>m</sup>	E : 95.6(0.5) keV above <sup>143</sup> Dy <sup>m</sup>						**	
* <sup>143</sup> Dy <sup>n</sup>	J : from depopulating 95.6-keV gamma being most likely E2 in 05Ri17						**	
<sup>144</sup> Te	-36220#	300#	93 ms 60	0 <sup>+</sup>	18 20Wu04 TD	2015	$\beta^- = 100; \beta^-_n ?; \beta^-_{2n} ?$	
<sup>144</sup> I	-45330#	400#	94 ms 8	1 <sup>-</sup> #	01 20Wu04 TD	1994	$\beta^- = 100; \beta^-_n ?; \beta^-_{2n} ?$	
<sup>144</sup> Xe	-56872	5	388 ms 7	0 <sup>+</sup>	01 03Be05 TD	2003	$\beta^- = 100; \beta^-_n = 3.0 3$	
<sup>144</sup> Cs	-63271	20	994 ms 6	1 <sup>-</sup> *	10 20Cz01 D	1967	$\beta^- = 100; \beta^-_n = 2.98 6$	
<sup>144</sup> Cs <sup>m</sup>	-63179	20	92.2 0.5	1.1 $\mu$ s 0.1	(4 <sup>-</sup> )	10	IT=100	
<sup>144</sup> Cs <sup>n</sup>		<i>non-exist</i>	EU	< 1 s	(> 3 <sup>-</sup> )	10 78MoZQ IJT	1978	$\beta^- = ?; IT ?; \beta^-_n ?$
<sup>144</sup> Ba	-71767	7	11.73 s 0.08	0 <sup>+</sup>	01 19KoZX T	1967	$\beta^- = 100$	
<sup>144</sup> La	-74850	13	44.0 s 0.7	(3 <sup>-</sup> )	01 FGK205 T	1967	$\beta^- = 100$	
<sup>144</sup> Ce	-80431.9	2.8	284.886 d 0.025	0 <sup>+</sup>	01 FGK209 T	1945	$\beta^- = 100$	
<sup>144</sup> Pr	-80750.6	2.7	17.28 m 0.05	0 <sup>-</sup>	01	1951	$\beta^- = 100$	
<sup>144</sup> Pr <sup>m</sup>	-80691.6	2.7	59.03 0.03	7.2 m 0.3	3 <sup>-</sup>	01	IT $\approx$ 100; $\beta^- \approx 0.07$	
<sup>144</sup> Nd	-83748.0	1.3	2.29 Py 0.16	0 <sup>+</sup>	01	1924	IS=23.798 19; $\alpha=100$	
<sup>144</sup> Pm	-81416.1	2.9	363 d 14	5 <sup>-</sup>	01 94Hi05 D	1952	$\epsilon=100; e^+ < 8e-5$	
<sup>144</sup> Pm <sup>m</sup>	-80575.2	2.9	840.90 0.05	780 ns 200	(9 <sup>+</sup> )	01	IT=100	
<sup>144</sup> Pm <sup>n</sup>	-72820	4	8595.8 2.2	$\sim 2.7 \mu$ s	(27 <sup>+</sup> )	01	IT=100	
<sup>144</sup> Sm	-81965.6	1.5	STABLE	0 <sup>+</sup>	01	1933	IS=3.08 4; $2\beta^+ ?$	
<sup>144</sup> Sm <sup>m</sup>	-79642.0	1.5	2323.60 0.08	880 ns 25	6 <sup>+</sup>	01	IT=100	
<sup>144</sup> Eu	-75619	11	10.2 s 0.1	1 <sup>+</sup>	01	1965	$\beta^+ = 100$	
<sup>144</sup> Eu <sup>m</sup>	-74491	11	1127.6 0.6	1.0 $\mu$ s 0.1	8 <sup>-</sup>	01 FGK127 J	1976	IT=100
<sup>144</sup> Gd	-71760	28	4.47 m 0.06	0 <sup>+</sup>	01	1968	$\beta^+ = 100$	
<sup>144</sup> Gd <sup>m</sup>	-68327	28	3433.1 0.5	145 ns 30	(10 <sup>+</sup> )	01	IT=100	
<sup>144</sup> Tb	-62368	28	$\sim 1$ s	1 <sup>+</sup>	01	1982	$\beta^+ = 100$	
<sup>144</sup> Tb <sup>m</sup>	-61971	28	396.9 0.5	4.25 s 0.15	6 <sup>-</sup>	01	IT=66; $\beta^+ = 34$	
<sup>144</sup> Tb <sup>n</sup>	-61892	28	476.2 0.5	2.8 $\mu$ s 0.3	(8 <sup>-</sup> )	01	IT=100	
<sup>144</sup> Tb <sup>p</sup>	-61851	28	517.1 0.5	670 ns 60	(9 <sup>+</sup> )	01	IT=100	
<sup>144</sup> Tb <sup>q</sup>	-61824	28	544.5 0.6	< 300 ns	(10 <sup>+</sup> )	01	IT=100	
<sup>144</sup> Dy	-56570	7	9.1 s 0.4	0 <sup>+</sup>	01	1986	$\beta^+ = 100; \beta^+ p = ?$	
<sup>144</sup> Ho	-44610	8	700 ms 100	(5 <sup>-</sup> )	08	1986	$\beta^+ = 100; \beta^+ p = ?$	
<sup>144</sup> Ho <sup>m</sup>	-44345	8	265.3 0.3	519 ns 5	(8 <sup>+</sup> )	08 10Ma08 T	2001	IT=100
<sup>144</sup> Er	-36610#	200#	400# ms >200ns	0 <sup>+</sup>	06	2003	$\beta^+ ?$	
<sup>144</sup> Tm	-22160#	400#	2.3 $\mu$ s 0.9	(10 <sup>+</sup> )	08	2005	$p = ?; \beta^+ ?$	
* <sup>144</sup> Cs	D : % $\beta^-_n$ average 20Cz01=2.95(0.24) 93Ru01=3.17(0.13) 79Ri09=2.95(0.25)						**	
* <sup>144</sup> Cs	D : 80ReZQ=3.12(0.11), 2.67(0.12)						**	
* <sup>144</sup> Cs	T : other (recent) 17Wu04=932(76)						**	
* <sup>144</sup> Cs <sup>n</sup>	I : introduced in 78MoZQ, but no $\beta^-$ decaying isomer was observed in later						**	
* <sup>144</sup> Cs <sup>m</sup>	I : studies; most likely this is <sup>144</sup> Cs <sup>m</sup>						**	
* <sup>144</sup> Ba	T : average 19KoZX=11.6(0.1) 82Ch22=11.5(0.2) 79En02=12.0(0.4)						**	
* <sup>144</sup> Ba	T : 76AmZW=11.9(0.6) 78Wo09=12.3(0.4) 74Gr29=11.1(0.5)						**	
* <sup>144</sup> Ba	T : 69WiZX=12.3(0.2) 69Ru14=11.9(0.3)						**	
* <sup>144</sup> La	T : other Ensdf2001=40.8(0.4) is likely affected by <sup>144</sup> Ba impurities						**	
* <sup>144</sup> Sm	T : 0nu-BB 18No01 > 1 Py						**	
* <sup>144</sup> Eu <sup>m</sup>	J : E2 to 6-						**	
* <sup>144</sup> Tb <sup>m</sup>	T : other 03Li42=12(2) s for q=65+ (bare ion)						**	
* <sup>144</sup> Tb <sup>m</sup>	J : E3 to 3+						**	
* <sup>144</sup> Tm	T : symmetrized from 05Gr32,05Bi24=1.9(+1.2-0.5) us						**	
<sup>145</sup> Te	-30010#	300#	75# ms >550ns	0 <sup>+</sup>	18 18Sh11 I	2018	$\beta^- ?; \beta^-_n ?; \beta^-_{2n} ?$	
<sup>145</sup> I	-41130#	500#	89.7 ms 9.3	7/2 <sup>+</sup> #	10 20Wu04 TD	2010	$\beta^- = 100; \beta^-_n ?; \beta^-_{2n} ?$	
<sup>145</sup> Xe	-51493	11	188 ms 4	3/2 <sup>-</sup> #	09	2003	$\beta^- = 100; \beta^-_n = 5.0 6; \beta^-_{2n} ?$	
<sup>145</sup> Cs	-60054	9	582 ms 4	3/2 <sup>+</sup> *	09 20Cz01 D	1971	$\beta^- = 100; \beta^-_n = 12.8 3$	
<sup>145</sup> Cs <sup>m</sup>	-59291	9	762.9 0.4	500 ns 100	13/2#	15YaZW TD	2015	IT=100
<sup>145</sup> Ba	-67516	8	4.31 s 0.16	5/2 <sup>-</sup> *	09	1974	$\beta^- = 100$	
<sup>145</sup> La	-72835	12	24.8 s 2.0	(5/2 <sup>+</sup> )	09	1974	$\beta^- = 100$	
<sup>145</sup> Ce	-77070	30	3.01 m 0.06	5/2 <sup>-</sup> #	09	1954	$\beta^- = 100$	
<sup>145</sup> Pr	-79626	7	5.984 h 0.010	7/2 <sup>+</sup>	09	1954	$\beta^- = 100$	



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
* <sup>146</sup> La	T : average FGK205=9.9(0.1) 78MoYW=10.0(0.4) 74Ar17=11(1) 79En02=9.0(0.6)						**	
* <sup>146</sup> La	J : region of deformation p5/2[413] n5/2[523],K=5-;						**	
* <sup>146</sup> La	J : Ensdf2016=(6-) based on shell model						**	
* <sup>146</sup> La <sup>m</sup>	T : average FGK205=6.1(0.3) 78MoYW=6.2(0.6) 81GoZN=6.0(0.4)						**	
* <sup>146</sup> La <sup>m</sup>	J : region of deformation p5/2[413] n1/2[530] or n5/2[523], K=1- or 2-;						**	
* <sup>146</sup> La <sup>m</sup>	J : Ensdf2016=(2-) based on shell model						**	
* <sup>146</sup> Nd	T : partial $\alpha$ half-life 15St09>1.6 Ey; other 0nu-BB 18No01>45 Ey						**	
* <sup>146</sup> Sm	T : from 12Ki16t=68(7); others 87Me08=103.1(4.5) 66Fr11=102.6(4.8)						**	
* <sup>146</sup> Tb <sup>n</sup>	E : 779.57 keV above <sup>146</sup> Tb <sup>m</sup> from Ensdf2016						**	
* <sup>146</sup> Tm	T : other 05Bb02=190(80) ms						**	
* <sup>146</sup> Tm <sup>m</sup>	T : average 06Ta08=68(3), supersedes 05Bb02=75(3), 05Ro40=82(4);						**	
* <sup>146</sup> Tm <sup>m</sup>	T : Birge ratio=2.8						**	
* <sup>146</sup> Tm <sup>n</sup>	T : average 07DaZU=213(9) 06Ta08=198(3)						**	
<sup>147</sup> I	-31200#	300#	60# ms >550ns	3/2 <sup>-</sup> #	18Sh11 I	2018	$\beta^-$ ?; $\beta^-_n$ ?; $\beta^-_{2n}$ ?	
<sup>147</sup> Xe	-42400#	200#	88 ms 14	3/2 <sup>-</sup> #	09 20Wu04 T	1994	$\beta^-$ =100; $\beta^-_n$ <8; $\beta^-_{2n}$ ?	
<sup>147</sup> Cs	-51920	8	230.5 ms 0.9	(3/2 <sup>+</sup> )	09 93Ru01 TD	1978	$\beta^-$ =100; $\beta^-_n$ =28.5 15	
<sup>147</sup> Cs <sup>m</sup>	-51219	8	190 ns 20	13/2#	15YaZW TD	2015	IT=100	
<sup>147</sup> Ba	-60264	20	893 ms 1	5/2 <sup>-</sup>	09 13Rz01 J	1978	$\beta^-$ =100; $\beta^-_n$ =0.07 5	
<sup>147</sup> La	-66678	11	4.026 s 0.020	(5/2 <sup>+</sup> )	09 96Ur02 J	1979	$\beta^-$ =100; $\beta^-_n$ =0.041 3	
<sup>147</sup> Ce	-72014	9	56.4 s 1.0	(5/2 <sup>-</sup> )	09	1964	$\beta^-$ =100	
<sup>147</sup> Pr	-75444	16	13.39 m 0.04	3/2 <sup>+</sup>	09 15Ru09 T	1964	$\beta^-$ =100	
<sup>147</sup> Nd	-78146.8	1.3	10.98 d 0.01	5/2 <sup>-</sup> *	09	1947	$\beta^-$ =100	
<sup>147</sup> Pm	-79042.0	1.3	2.6234 y 0.0002	7/2 <sup>+</sup> *	09	1947	$\beta^-$ =100	
<sup>147</sup> Sm	-79266.0	1.3	106.6 Gy 0.5	7/2 <sup>-</sup> *	09 FGK204 T	1933	IS=15.00 14; $\alpha$ =100	
<sup>147</sup> Eu	-77544.6	2.6	24.1 d 0.6	5/2 <sup>+</sup> *	09	1951	$\beta^+$ $\approx$ 100; $\alpha$ =0.0022 6	
<sup>147</sup> Eu <sup>m</sup>	-76919.3	2.6	625.27 0.05	765 ns 15	09	1970	IT=100	
<sup>147</sup> Gd	-75356.9	1.9	38.06 h 0.12	7/2 <sup>-</sup> *	09	1957	$\beta^+$ =100	
<sup>147</sup> Gd <sup>m</sup>	-66769.1	2.0	8587.8 0.5	510 ns 20	09 20Br06 J	1982	IT=100	
<sup>147</sup> Tb	-70743	8	1.64 h 0.03	(1/2 <sup>+</sup> )	09	1969	$\beta^+$ =100	
<sup>147</sup> Tb <sup>m</sup>	-70692	8	50.6 0.9	1.87 m 0.05	(11/2 <sup>-</sup> )	09 93Al03 T	1987	$\beta^+$ =100
<sup>147</sup> Dy	-64196	9	67 s 7	(1/2 <sup>+</sup> )	09	1975	$\beta^+$ =100; $\beta^+$ p $\approx$ 0.05	
<sup>147</sup> Dy <sup>m</sup>	-63446	9	750.5 0.4	55.2 s 0.5	(11/2 <sup>-</sup> )	09	$\beta^+$ =68.9 23; IT=31.1 23	
<sup>147</sup> Dy <sup>n</sup>	-60789	9	3407.2 0.8	400 ns 10	(27/2 <sup>-</sup> )	09	IT=100	
<sup>147</sup> Ho	-55757	5	5.8 s 0.4	(11/2 <sup>-</sup> )	09	1982	$\beta^+$ =100	
<sup>147</sup> Ho <sup>m</sup>	-53070	5	2687.1 0.4	315 ns 30	(27/2 <sup>-</sup> )	09	IT=100	
<sup>147</sup> Er	-46610	40	3.2 s 1.2	(1/2 <sup>+</sup> )	09 10Ma27 T	1992	$\beta^+$ =100; $\beta^+$ p=?	
<sup>147</sup> Er <sup>m</sup>	-46510#	60#	100# 50#	1.6 s 0.2	(11/2 <sup>-</sup> )	09 10Ma27 T	$\beta^+$ =100; $\beta^+$ p=?	
<sup>147</sup> Tm	-35974	7	580 ms 30	11/2 <sup>-</sup>	09	1982	$\beta^+$ =85 5; p=15 5	
<sup>147</sup> Tm <sup>m</sup>	-35913	7	62 5 p	360 $\mu$ s 40	3/2 <sup>+</sup>	09	p=100	
* <sup>147</sup> Xe	T : other 03Be05=100(+100-50)						**	
* <sup>147</sup> Cs	T : average 20Wu04=255(5) 17Wu04=234(14) 93Ru01=235(3) 86ReZU=229(1)						**	
* <sup>147</sup> Cs	T : 79Ri09=214(30) 78Ko29=235(10)						**	
* <sup>147</sup> Cs	D : % $\beta^-_n$ average 93Ru01=30.7(2.0) 86ReZU=26.4(2.9) 79Ri09=25.4(3.2)						**	
* <sup>147</sup> Cs <sup>m</sup>	E : from 16Ya.A=701.4(0.4)						**	
* <sup>147</sup> Ba	T : average 17Wu04=921(47) 93Ru01=894(10) 86Wa17=893(1), supersedes 86ReZU						**	
* <sup>147</sup> Ba	D : % $\beta^-_n$ unweighted average 93Ru01=0.110(0.016) 86Wa17=0.019(1);						**	
* <sup>147</sup> Ba	D : Birge ratio=5.68; other 81En05=5.21(52), outlier						**	
* <sup>147</sup> La	T : average 93Ru01=4.100(0.021) 86Wa17=4.015(0.008) 81En05=4.10(0.25);						**	
* <sup>147</sup> La	T : Birge ratio=2.68						**	
* <sup>147</sup> La	D : % $\beta^-_n$ average 93Ru01=0.043(0.004) 86Wa17=0.035(0.006)						**	
* <sup>147</sup> Pr	J : from 15Wa28						**	
* <sup>147</sup> Tb <sup>m</sup>	T : average 93Al03=1.92(0.07) 73Bo13=1.83(0.06)						**	
<sup>148</sup> Xe	-38650#	300#	85 ms 15	0 <sup>+</sup>	14 20Wu04 TD	2010	$\beta^-$ =100; $\beta^-_n$ ?; $\beta^-_{2n}$ ?	
<sup>148</sup> Cs	-46911	13	151.8 ms 1.0	(2 <sup>-</sup> )	14 20Wu04 T	1978	$\beta^-$ =100; $\beta^-_n$ =28.7 21; $\beta^-_{2n}$ ?	
<sup>148</sup> Cs <sup>m</sup>	-46866	13	45.2 0.1	4 <sup>-</sup> #	15YaZW TD	2015	IT=100	
<sup>148</sup> Ba	-57544.9	1.5	620 ms 5	0 <sup>+</sup>	14 17Wu04 T	1979	$\beta^-$ =100; $\beta^-_n$ =0.4 3	
<sup>148</sup> La	-62709	19	1.414 s 0.025	(2 <sup>-</sup> )	14 17Wu04 T	1969	$\beta^-$ =100; $\beta^-_n$ =0.18 7	
<sup>148</sup> Ce	-70398	11	56.8 s 0.3	0 <sup>+</sup>	14	1964	$\beta^-$ =100	

Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)		
<sup>148</sup> Pr	-72535	15	2.29 m	0.02	1 <sup>-</sup>	14	1964	$\beta^-$ =100	
<sup>148</sup> Pr <sup>m</sup>	-72458	15	2.01 m	0.07	(4)	14	1964	$\beta^-$ =64 10;IT=36 10	
<sup>148</sup> Nd	-77408.1	2.1	STABLE	>3.0Ey	0 <sup>+</sup>	14 82Be20	T 1937	IS=5.756 21;2 $\beta^-$ ?; $\alpha$ ?	*
<sup>148</sup> Pm	-76866	6	5.368 d	0.007	1 <sup>-*</sup>	14	1947	$\beta^-$ =100	
<sup>148</sup> Pm <sup>m</sup>	-76728	6	41.29 d	0.11	5 <sup>-</sup> ,6 <sup>-</sup>	14	1951	$\beta^-$ =95.8 6;IT=4.2 6	
<sup>148</sup> Sm	-79336.1	1.2	6.3 Py	1.3	0 <sup>+</sup>	14 16Ca43	T 1933	IS=11.25 9; $\alpha$ =100	*
<sup>148</sup> Eu	-76297	10	54.5 d	0.5	5 <sup>-*</sup>	14	1951	$\beta^+$ =100; $\alpha$ =9.4e-7 28	
<sup>148</sup> Eu <sup>m</sup>	-75577	10	162 ns	8	9 <sup>+</sup>	14	1980	IT=100	
<sup>148</sup> Gd	-76269.4	1.5	71.3 y	1.0	0 <sup>+</sup>	14 03Fu10	T 1953	$\alpha$ =100;2 $\beta^+$ ?	*
<sup>148</sup> Tb	-70537	12	60 m	1	2 <sup>-</sup>	14	1960	$\beta^+$ =100	
<sup>148</sup> Tb <sup>m</sup>	-70447	12	2.20 m	0.05	(9) <sup>+</sup>	14	1973	$\beta^+$ =100	
<sup>148</sup> Tb <sup>n</sup>	-61918	12	1.310 $\mu$ s	0.007	(27) <sup>+</sup>	14	1980	IT=100	
<sup>148</sup> Dy	-67859	9	3.3 m	0.2	0 <sup>+</sup>	14	1974	$\beta^+$ =100	
<sup>148</sup> Dy <sup>m</sup>	-64940	9	471 ns	20	10 <sup>+</sup>	14	1978	IT=100	
<sup>148</sup> Ho	-57990	80	2.2 s	1.1	(1) <sup>+</sup>	14	1979	$\beta^+$ =100	
<sup>148</sup> Ho <sup>n</sup>	-57740#	130#	9.49 s	0.12	(5 <sup>-</sup> )	14 93Al03	T 1979	$\beta^+$ =100; $\beta^+$ p=0.08 1	*
<sup>148</sup> Ho <sup>n</sup>	-57050#	130#	2.36 ms	0.06	(10) <sup>+</sup>	14	1984	IT=100[gs=0,m=100]	*
<sup>148</sup> Er	-51479	10	4.6 s	0.2	0 <sup>+</sup>	14	1982	$\beta^+$ =100; $\beta^+$ p $\approx$ 0.15	
<sup>148</sup> Er <sup>m</sup>	-48566	10	13 $\mu$ s	3	(10 <sup>+</sup> )	14	1982	IT=100	
<sup>148</sup> Tm	-38765	10	700 ms	200	(10 <sup>+</sup> )	14	1982	$\beta^+$ =100; $\beta^+$ p ?	
<sup>148</sup> Yb	-30230#	400#	250# ms		0 <sup>+</sup>			$\beta^+$ ?; $\beta^+$ p ?	
* <sup>148</sup> Cs	T : average 20Wu04=144(9) 18Li06=158(6) 17Li06=152(1) 17Wu04=144(5)							**	
* <sup>148</sup> Cs	T : 93Ru01=140(12) 86Hi08=158(7); others 86Wa17=130(10) 78Ko29=130(40)							**	
* <sup>148</sup> Cs	J : direct $\beta^-$ decay feeding to 1- and 3- levels in <sup>148</sup> Ba in 18Li06							**	
* <sup>148</sup> Cs	D : % $\beta^-$ -n average 86RuZU=25.1(2.5) 93Ru01=24(17) 18Li06=38(4)							**	
* <sup>148</sup> Cs <sup>m</sup>	E : from 16Ya.A=45.2(0.1)							**	
* <sup>148</sup> Ba	T : average 17Wu04=621(11) 86Wa17=620(5) 84Ch02=607(25) 82Ga24=630(50)							**	
* <sup>148</sup> La	T : average 17Wu04=1.27(+0.10-0.09) 93Ru01=1.428(0.012) 86Wa17=1.40(0.02)							**	
* <sup>148</sup> La	T : 83Mu19=1.34(0.02) 82Ga24=1.55(0.03) 69Wi.A=1.29(0.08);							**	
* <sup>148</sup> La	T : Birge ratio=2.83							**	
* <sup>148</sup> La	D : % $\beta^-$ -n unweighted average 93Ru01=0.24(0.02) 86Wa17=0.11(0.01);							**	
* <sup>148</sup> La	D : Birge ratio=5.81							**	
* <sup>148</sup> Nd	T : lower limit is for 2 $\beta^-$ decay							**	
* <sup>148</sup> Sm	T : symmetrized from 16Ca43=6.4(+1.2-1.3)							**	
* <sup>148</sup> Gd	T : average 03Fu10=70.9(1.0) 81Pr06=74.6(3.0)							**	
* <sup>148</sup> Ho <sup>m</sup>	T : average 93Al03=9.30(0.20) 89Ta11=9.59(0.15)							**	
* <sup>148</sup> Ho <sup>n</sup>	E : 694.4 keV above <sup>148</sup> Ho <sup>m</sup> from Ensdf2014							**	
<sup>149</sup> Xe	-33000#	300#	50# ms	>550ns	3/2 <sup>-</sup> #	18Sh11	I 2018	$\beta^-$ ?; $\beta^-$ n ?; $\beta^-$ 2n ?	
<sup>149</sup> Cs	-43300#	400#	112.3 ms	2.5	3/2 <sup>+</sup> #	17 17Li06	TD 1979	$\beta^-$ =100; $\beta^-$ -n=25 4; $\beta^-$ 2n ?	*
<sup>149</sup> Ba	-52830.6	2.5	349 ms	4	3/2 <sup>-</sup> #	04 20Wu04	T 1993	$\beta^-$ =100; $\beta^-$ -n=3.9 12	*
<sup>149</sup> La	-60220	200	1.071 s	0.022	(3/2 <sup>-</sup> )	07 17Wu04	T 1979	$\beta^-$ =100; $\beta^-$ -n=1.43 28	*
<sup>149</sup> Ce	-66670	10	4.94 s	0.04	3/2 <sup>-</sup> #	04 96Ya.A	T 1974	$\beta^-$ =100	
<sup>149</sup> Pr	-71039	10	2.26 m	0.07	(5/2 <sup>+</sup> )	04	1964	$\beta^-$ =100	
<sup>149</sup> Nd	-74375.5	2.1	1.728 h	0.001	5/2 <sup>-*</sup>	04	1938	$\beta^-$ =100	
<sup>149</sup> Pm	-76064.4	2.2	53.08 h	0.05	7/2 <sup>+</sup> *	04	1947	$\beta^-$ =100	
<sup>149</sup> Pm <sup>m</sup>	-75824.2	2.2	35 $\mu$ s	3	11/2 <sup>-</sup>	04	1966	IT=100	
<sup>149</sup> Sm	-77135.9	1.2	STABLE	>2Py	7/2 <sup>-*</sup>	04	1933	IS=13.82 10; $\alpha$ ?	
<sup>149</sup> Eu	-76441	4	93.1 d	0.4	5/2 <sup>+</sup> *	04	1959	$\epsilon$ =100	
<sup>149</sup> Eu <sup>m</sup>	-75945	4	2.45 $\mu$ s	0.05	11/2 <sup>-</sup>	04	1961	IT=100	
<sup>149</sup> Gd	-75127	3	9.28 d	0.10	7/2 <sup>-*</sup>	04	1951	$\beta^+$ =100; $\alpha$ =4.3e-4 10	
<sup>149</sup> Tb	-71489	4	4.118 h	0.025	1/2 <sup>+</sup>	04	1950	$\beta^+$ =83.3 17; $\alpha$ =16.7 17	
<sup>149</sup> Tb <sup>m</sup>	-71453	4	4.16 m	0.04	11/2 <sup>-</sup>	04	1962	$\beta^+$ $\approx$ 100; $\alpha$ =0.022 3	
<sup>149</sup> Dy	-67694	9	4.20 m	0.14	7/2 <sup>-</sup>	04	1958	$\beta^+$ =100	
<sup>149</sup> Dy <sup>m</sup>	-65033	9	490 ms	15	27/2 <sup>-</sup>	04 80Da18	J 1976	IT=99.3 3; $\beta^+$ =0.7 3	*
<sup>149</sup> Ho	-61646	12	21.1 s	0.2	(11/2 <sup>-</sup> )	04	1979	$\beta^+$ =100	
<sup>149</sup> Ho <sup>m</sup>	-61597	12	56 s	3	(1/2 <sup>+</sup> )	04	1988	$\beta^+$ =100	
<sup>149</sup> Er	-53742	28	4 s	2	(1/2 <sup>+</sup> )	04	1984	$\beta^+$ =100; $\beta^+$ p=7 2	
<sup>149</sup> Er <sup>m</sup>	-53000	28	8.9 s	0.2	(11/2 <sup>-</sup> )	04	1984	$\beta^+$ =96.5 7;IT=3.5 7; $\beta^+$ p=0.18 7	
<sup>149</sup> Er <sup>n</sup>	-51131	28	610 ns	80	(19/2 <sup>+</sup> )	04	1987	IT=100	
<sup>149</sup> Er <sup>p</sup>	-50440	30	4.8 $\mu$ s	0.1	(27/2 <sup>-</sup> )	04 87Br14	EJD 1987	IT=100	*

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)			Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)		
<sup>149</sup> Tm	-43940#	200#			900 ms 200	11/2 <sup>-</sup>	04		1987	$\beta^+=100;\beta^+p=0.26$ 15	*	
<sup>149</sup> Tm <sup>m</sup>	-43840#	210#	100#	50#	500# ms	1/2 <sup>+</sup> #				$\beta^+ ?;\beta^+p ?$	*	
<sup>149</sup> Yb	-33330#	300#			700 ms 200	(1/2 <sup>+</sup> )	04	05Xu04	J 2001	$\beta^+=100;\beta^+p\approx 100$	*	
* <sup>149</sup> Cs	T : average 20Wu04=113(6) 17Wu04=113(8) 00KoZH=112(3),114(16); other											
* <sup>149</sup> Cs	T : 18Li06=80(3) 17Li06=80(4), outliers											
* <sup>149</sup> Ba	T : average 20Wu04=368(19) 17Wu04=352(6) 93Ru01=324(18) 86Wa17=346(6)											
* <sup>149</sup> Ba	D : % $\beta^-$ n from 93Ru01; other 86Wa17=0.43(12), discrepant											
* <sup>149</sup> La	T : average 17Wu04=1.11(0.04) 93Ru01=1.066(0.034) 86Wa17=1.04(0.04)											
* <sup>149</sup> La	D : % $\beta^-$ n average 93Ru01=1.74(0.13) 86Wa17=1.17(12), Birge ratio=3.22											
* <sup>149</sup> Dy <sup>m</sup>	T : other 03Li42=11(1) s for q=66+ (bare ion)											
* <sup>149</sup> Er <sup>p</sup>	E : 661+x keV above <sup>149</sup> Er <sup>n</sup> and x<60 keV in 87Br14											
* <sup>149</sup> Tm	D : symmetrized from $\beta^+p=0.2(+0.2-0.1)\%$											
* <sup>149</sup> Tm	J : favorite $\alpha$ decay from <sup>153</sup> Lu (J=11/2-)											
* <sup>149</sup> Tm <sup>m</sup>	I : probably fed by $\alpha$ -decaying isomer in <sup>153</sup> Lu											
* <sup>149</sup> Yb	J : (1/2+,3/2+) in Ensdf2004 and 1/2 in 05Xu04; 06Xu07=(1/2-), however,											
* <sup>149</sup> Yb	J : no 1/2- gs or isomer for N=79 isotones											
<sup>150</sup> Xe	-28990#	300#			40# ms >550ns	0 <sup>+</sup>		18Sh11	I 2018	$\beta^- ?;\beta^-n ?;\beta^-2n ?$		
<sup>150</sup> Cs	-38170#	400#			81.0 ms 2.6	(2 <sup>-</sup> )	17	18Li06	TD 1979	$\beta^-=100;\beta^-n\approx 44;\beta^-2n ?$	*	
<sup>150</sup> Ba	-49890	6			258 ms 5	0 <sup>+</sup>	17	02Pf04	TD 1994	$\beta^-=100;\beta^-n=1.0$ 5	*	
<sup>150</sup> La	-56311.1	2.5			504 ms 15	(3 <sup>+</sup> )	13	17Wu04	T 1993	$\beta^-=100;\beta^-n=2.7$ 3	*	
<sup>150</sup> Ce	-64847	12			6.05 s 0.07	0 <sup>+</sup>	13	15Ko23	T 1970	$\beta^-=100$		
<sup>150</sup> Pr	-68301	9			6.19 s 0.16	1 <sup>-</sup>	13	15Ko23	J 1970	$\beta^-=100$		
<sup>150</sup> Nd	-73680.0	1.1			9.3 Ey 0.7	0 <sup>+</sup>	13	20Ba.A	T 1937	IS=5.638 28;2 $\beta^-$ =100		
<sup>150</sup> Pm	-73597	20			2.698 h 0.015	(1 <sup>-</sup> )	13		1952	$\beta^-=100$		
<sup>150</sup> Sm	-77051.3	1.1			STABLE	0 <sup>+</sup>	13		1934	IS=7.37 9		
<sup>150</sup> Eu	-74792	6			36.9 y 0.9	5 <sup>-</sup>	13		1950	$\beta^+=100$		
<sup>150</sup> Eu <sup>m</sup>	-74750	6	41.7	1.0	12.8 h 0.1	0 <sup>-</sup> *	13		1953	$\beta^-=89$ 2; $\beta^+=11$ 2;IT ?		
<sup>150</sup> Gd	-75764	6			1.79 My 0.08	0 <sup>+</sup>	13		1953	$\alpha=100;2\beta^+ ?$		
<sup>150</sup> Tb	-71106	7			3.48 h 0.16	(2 <sup>-</sup> )	13		1959	$\beta^+\approx 100;\alpha ?$		
<sup>150</sup> Tb <sup>m</sup>	-70645	26	461	27	MD	5.8 m 0.2	9 <sup>+</sup>	13	1993	$\beta^+\approx 100;IT ?$		
<sup>150</sup> Dy	-69310	4			7.17 m 0.05	0 <sup>+</sup>	13		1959	$\beta^+=66.4$ 18; $\alpha=33.6$ 18	*	
<sup>150</sup> Ho	-61946	14			76.8 s 1.8	(2 <sup>-</sup> )	13	93Al03	T 1963	$\beta^+=100$	*	
<sup>150</sup> Ho <sup>m</sup>	-61950	50	0	50	BD*	23.3 s 0.3	(9 <sup>+</sup> )	13	1980	$\beta^+=100$		
<sup>150</sup> Ho <sup>n</sup>	-54050	50	7900	50		787 ns 36	(28 <sup>-</sup> )	13	06Fu06	JTE 2006	IT=100	*
<sup>150</sup> Er	-57831	17			18.5 s 0.7	0 <sup>+</sup>	13		1982	$\beta^+=100$		
<sup>150</sup> Er <sup>m</sup>	-55035	17	2796.5	0.5		2.55 $\mu$ s 0.10	10 <sup>+</sup>	13	1984	IT=100		
<sup>150</sup> Tm	-46490#	200#			3# s	(1 <sup>+</sup> )		88Ni02	Jl 1982	$\beta^+=100$		
<sup>150</sup> Tm <sup>m</sup>	-46350#	240#	140#	140#	2.20 s 0.06	(6 <sup>-</sup> )	13		1981	$\beta^+=100;\beta^+p=1.1$ 3	*	
<sup>150</sup> Tm <sup>n</sup>	-45680#	240#	811#	140#	5.2 ms 0.3	10 <sup>+</sup> #	13		1984	IT=100[gs=0,m=100]	*	
<sup>150</sup> Yb	-38830#	300#			700# ms >200ns	0 <sup>+</sup>	13		2000	$\beta^+ ?$		
<sup>150</sup> Lu	-24770#	300#			45 ms 3	(5 <sup>-</sup> )	13	03Gi10	J 1993	$p\approx 100;\beta^+ ?$		
<sup>150</sup> Lu <sup>m</sup>	-24750#	300#	22	5	p	40 $\mu$ s 7	(8 <sup>+</sup> )	13	03Gi10	J 1998	$p=100$	*
* <sup>150</sup> Cs	T : average 20Wu04=90(15) 18Li06=80(3) 17Wu04=84.4(8.2) 00KoZH=82(7)											
* <sup>150</sup> Cs	D : % $\beta^-$ n other 00KoZH=20(10)											
* <sup>150</sup> Cs	J : direct $\beta^-$ decay feeding to 1- and 3- levels in <sup>150</sup> Ba in 18Li06											
* <sup>150</sup> Ba	T : average 20Wu04=245(16) 17Wu04=259(5); other 02Pf04 300, compilation											
* <sup>150</sup> La	T : average 17Wu04=510(+10-22) 95Ok02=510(30)											
* <sup>150</sup> Dy	D : % $\alpha$ average 74To07=31(3),36(3) 73Bi06=32(5) 77Ha48=36(5)											
* <sup>150</sup> Ho	T : average 93Al03=78(2) 82No08=72(4)											
* <sup>150</sup> Ho <sup>n</sup>	E : 7912.1(2.3) keV above <sup>150</sup> Ho <sup>m</sup> from Ensdf2013											
* <sup>150</sup> Tm <sup>m</sup>	D : % $\beta^+p$ symmetrized from 88Ni02=1.2(+2-4)											
* <sup>150</sup> Tm <sup>n</sup>	E : 671.3(1.0) keV above <sup>150</sup> Tm <sup>m</sup> from Ensdf2013											
* <sup>150</sup> Lu <sup>m</sup>	T : symmetrized from 03Gi10=39(+8-6)											
<sup>151</sup> Cs	-34280#	500#			59 ms 19	3/2 <sup>+</sup> #	17	20Wu04	TD 1979	$\beta^-=100;\beta^-n ?;\beta^-2n ?$	*	
<sup>151</sup> Ba	-44940#	400#			167 ms 5	3/2 <sup>-</sup> #	17	17Wu04	T 1994	$\beta^-=100;\beta^-n ?$	*	
<sup>151</sup> La	-53310	440			465 ms 24	1/2 <sup>+</sup> #	17	17Wu04	TD 1994	$\beta^-=100;\beta^-n ?$	*	
<sup>151</sup> Ce	-61225	18			1.76 s 0.06	(3/2 <sup>-</sup> )	09	10Si03	J 1997	$\beta^-=100$	*	
<sup>151</sup> Pr	-66780	12			18.90 s 0.07	(3/2 <sup>-</sup> )	09		1990	$\beta^-=100$		
<sup>151</sup> Pr <sup>m</sup>	-66745	12	35.10	0.10	50 $\mu$ s 8	(7/2 <sup>+</sup> )	09	12Ma03	T 2006	IT=100		



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>151</sup> Nd	-70943.2	1.1	12.44 m 0.07	3/2 <sup>+</sup>	09	1938	$\beta^- = 100$	
<sup>151</sup> Pm	-73386	5	28.40 h 0.04	5/2 <sup>+</sup> *	09	1952	$\beta^- = 100$	
<sup>151</sup> Sm	-74576.5	1.1	94.6 y 0.6	5/2 <sup>+</sup> *	09 15Be23 T	1947	$\beta^- = 100$	
<sup>151</sup> Sm <sup>m</sup>	-74315.4	1.1	1.4 $\mu$ s 0.1	(11/2) <sup>-</sup>	09	1973	IT=100	
<sup>151</sup> Eu	-74653.1	1.2	4.6 Ey 1.2	5/2 <sup>+</sup> *	09 14Ca13 T	1933	IS=47.81 6; $\alpha = 100$	
<sup>151</sup> Eu <sup>m</sup>	-74456.9	1.2	196.245 0.010	58.9 $\mu$ s 0.5	11/2 <sup>-</sup>	09	1958	IT=100
<sup>151</sup> Gd	-74188.9	3.0	123.9 d 1.0	7/2 <sup>-</sup> *	09	1950	$\epsilon = 100; \alpha \approx 1.1e-6$	
<sup>151</sup> Tb	-71624	4	17.609 h 0.001	1/2 <sup>+</sup> *	09	1953	$\beta^+ 99.9905 15; \alpha = 0.0095 15$	
<sup>151</sup> Tb <sup>m</sup>	-71524	4	25 s 3	11/2 <sup>-</sup>	09	1978	IT=93.4 20; $\beta^+ = 6.6 20$	
<sup>151</sup> Dy	-68752	3	17.9 m 0.3	7/2 <sup>-</sup> *	09	1959	$\beta^+ = 94.4 6; \alpha = 5.6 4$	
<sup>151</sup> Ho	-63623	8	35.2 s 0.1	11/2 <sup>-</sup> *	09 87NeZZ J	1963	$\beta^+ = 88 3; \alpha = 22 3$	
<sup>151</sup> Ho <sup>m</sup>	-63582	8	47.2 s 1.3	1/2 <sup>+</sup> *	09 87NeZZ J	1963	$\alpha = 77 18; \beta^+ = 23 18$	
<sup>151</sup> Er	-58266	16	23.5 s 2.0	(7/2) <sup>-</sup>	09	1970	$\beta^+ = 100$	
<sup>151</sup> Er <sup>m</sup>	-55680	16	2586.0 0.5	580 ms 20	(27/2) <sup>-</sup>	09	1980	IT=95.3 3; $\beta^+ = 4.7 3$
<sup>151</sup> Er <sup>n</sup>	-47979	16	10286.6 1.0	420 ns 50	(65/2 <sup>-</sup> , 61/2 <sup>+</sup> )	09 09Fu05 J	1990	IT=100
<sup>151</sup> Tm	-50772	19	4.17 s 0.11	(11/2) <sup>-</sup>	09	1982	$\beta^+ = 100$	
<sup>151</sup> Tm <sup>m</sup>	-50679	20	93 6 AD	(1/2 <sup>+</sup> )	09	1987	$\beta^+ = 100$	
<sup>151</sup> Tm <sup>n</sup>	-48116	19	2655.67 0.22	451 ns 34	(27/2) <sup>-</sup>	09	1982	IT=100
<sup>151</sup> Yb	-41540	300	1.6 s 0.5	(1/2 <sup>+</sup> )	09 86To12 T	1985	$\beta^+ = 100; \beta^+ p = ?$	
<sup>151</sup> Yb <sup>m</sup>	-40800#	320#	740# 100#	1.6 s 0.5	(11/2) <sup>-</sup>	09 86To12 T	1986	$\beta^+ \approx 100; \beta^+ p = ?; IT ?$
<sup>151</sup> Yb <sup>n</sup>	-38910#	330#	2630# 141#	2.6 $\mu$ s 0.7	19/2 <sup>-</sup> #	09	1993	IT=100
<sup>151</sup> Yb <sup>p</sup>	-38250#	330#	3287# 141#	20 $\mu$ s 1	27/2 <sup>-</sup> #	09	1987	IT=100
<sup>151</sup> Lu	-30300#	300#	78.4 ms 0.9	11/2 <sup>-</sup>	09 15Ta12 TJ	1982	$p = ?; \beta^+ = ?$	
<sup>151</sup> Lu <sup>m</sup>	-30240#	300#	57 4 p	16.0 $\mu$ s 0.5	3/2 <sup>+</sup>	09 17Wa18 T	1998	$p = 100$
* <sup>151</sup> Cs	T : average 20Wu04=48(28) 17Wu04=69(26)						**	
* <sup>151</sup> Ba	T : average 20Wu04=166(11) 17Wu04=167(5)						**	
* <sup>151</sup> La	T : symmetrized from 457(+30-18); other 20Wu04=510(330)						**	
* <sup>151</sup> Ce	T : average 17Wu04=1.71(0.09) 06Ko25=1.76 (0.06)						**	
* <sup>151</sup> Ce	I : isomer with T1/2=1.02(0.06)s suggested in Ensdf2009, but no sufficient						**	
* <sup>151</sup> Ce	I : experimental evidence exists, so it is not trusted by Nubase						**	
* <sup>151</sup> Sm	T : other (recent) 09He22=96.6(2.4)						**	
* <sup>151</sup> Eu	J : 90Al34=5/2						**	
* <sup>151</sup> Gd	D : % $\alpha$ symmetrized from $\alpha/KX$ rays=0.8(+0.8-0.4)e-8 in 65Si06						**	
* <sup>151</sup> Tb <sup>m</sup>	J : E3 to 5/2+ following by E2 to 1/2+						**	
* <sup>151</sup> Ho <sup>m</sup>	D : % $\alpha$ symmetrized from $\alpha = 80(+15-20)$						**	
* <sup>151</sup> Er <sup>m</sup>	T : other 03Li42=19(3) s for q=68+ (bare ion)						**	
* <sup>151</sup> Yb	T : derived from 1.6(0.1)s for a mixture of gs and isomer that have almost						**	
* <sup>151</sup> Yb	T : the same half-life						**	
* <sup>151</sup> Yb <sup>m</sup>	E : 740# keV estimated in 90Ak01						**	
* <sup>151</sup> Yb <sup>n</sup>	E : 1790+x keV above <sup>151</sup> Yb <sup>m</sup> in 93Ni05; x=100#(100#)						**	
* <sup>151</sup> Yb <sup>p</sup>	E : 657 keV above <sup>151</sup> Yb <sup>n</sup> in 93Ni05						**	
* <sup>151</sup> Lu	T : average 15Ta12=78(1) 99Bi14=80(2)						**	
* <sup>151</sup> Lu <sup>m</sup>	T : average 17Wa18=15.4(0.8) 15Ta12=17(1) 99Bi14=16(1)						**	
<sup>152</sup> Cs	-29130#	500#	17# ms		18 87Ra12 I	1987	$\beta^- ?; \beta^- n ?$	
<sup>152</sup> Ba	-41610#	400#	139 ms 8	0 <sup>+</sup>	17 17Wu04 TD	2010	$\beta^- = 100; \beta^- n ?$	
<sup>152</sup> La	-49290#	300#	287 ms 16	2 <sup>-</sup> #	17 17Wu04 TD	1994	$\beta^- = 100; \beta^- n ?$	
<sup>152</sup> Ce	-58980#	200#	1.42 s 0.02	0 <sup>+</sup>	13 17Wu04 T	1990	$\beta^- = 100$	
<sup>152</sup> Pr	-63758	19	3.57 s 0.11	4 <sup>+</sup>	13 99To04 J	1983	$\beta^- = 100$	
<sup>152</sup> Pr <sup>m</sup>	-63643	19	115.1 0.3	4.16 $\mu$ s 0.10	(1 <sup>+</sup> )	13 18Al14 TJE	1990	IT=100
<sup>152</sup> Nd	-70150	24	11.4 m 0.2	0 <sup>+</sup>	13	1969	$\beta^- = 100$	
<sup>152</sup> Pm	-71254	26	4.12 m 0.08	1 <sup>+</sup>	13	1958	$\beta^- = 100$	
<sup>152</sup> Pm <sup>m</sup>	-71110	80	140 90 BD*	7.52 m 0.08	4(-)	13	1971	$\beta^- = 100$
<sup>152</sup> Pm <sup>n</sup>		non-exist	EU	13.8 m 0.2	(8)	13	1971	$\beta^- = 100; IT ?$
<sup>152</sup> Sm	-74763.0	1.0	STABLE	0 <sup>+</sup>	13	1933	IS=26.74 9	
<sup>152</sup> Eu	-72888.5	1.2	13.517 y 0.006	3 <sup>-</sup> *	13 FGK209 T	1938	$\beta^+ = 72.08 13; \beta^- = 27.92 13$	
<sup>152</sup> Eu <sup>m</sup>	-72842.9	1.2	45.5998 0.0004	9.3116 h 0.0013	0 <sup>-</sup> *	13	1958	$\beta^- = 73 3; \beta^+ = 27 3$
<sup>152</sup> Eu <sup>n</sup>	-72823.2	1.2	65.2969 0.0004	940 ns 80	1 <sup>-</sup>	13	1978	IT=100
<sup>152</sup> Eu <sup>p</sup>	-72810.3	1.2	78.2331 0.0004	165 ns 10	1 <sup>+</sup>	13	1978	IT=100
<sup>152</sup> Eu <sup>q</sup>	-72798.7	1.2	89.8496 0.0004	384 ns 10	4 <sup>+</sup>	13	1970	IT=100
<sup>152</sup> Eu <sup>r</sup>	-72740.6	1.2	147.86 0.10	95.8 m 0.4	8 <sup>-</sup>	13 15Hu02 T	1963	IT=100
<sup>152</sup> Gd	-74707.3	1.0	108 Ty 8	0 <sup>+</sup>	13	1938	IS=0.20 3; $\alpha = 100; 2\beta^+ ?$	



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
<sup>153</sup> Hf <sup>m</sup>	-26550# 320#	750# 100#	500# ms	11/2 <sup>-</sup> #			$\beta^+ \text{ ?}; IT \text{ ?}$
* <sup>153</sup> Ba	T : average 20Wu04=109(59) 17Wu04=116(52)						
* <sup>153</sup> La	T : other 20Wu04=210(120)						
* <sup>153</sup> Pr	T : other 17Wu04=4.68(0.70) 87Gr12=4.3(0.2)						
* <sup>153</sup> Nd <sup>m</sup>	T : average 10Si03=1.17(0.07) 96Ya12=1.06(0.05)						
* <sup>153</sup> Eu	J : 85Ah03,90A134=5/2						
* <sup>153</sup> Gd	T : unweighted average 14Un01=239.29(0.10) 72Em01=241.6(0.2),240.9(0.6);						
* <sup>153</sup> Gd	T : Birge ratio=7.4; other (discrepant) 89Po21=226.7(4.2)						
* <sup>153</sup> Er	J : also 89Ot.A						
* <sup>153</sup> Yb <sup>m</sup>	E : from 93Mc03=2579 (23/2-) + x keV; x=50#(50#) keV by Nubase, probably						
* <sup>153</sup> Yb <sup>m</sup>	E : overlaps with 51-keV E1						
* <sup>153</sup> Lu	D : %p from 97Ir01=0						
<sup>154</sup> Ba	-32920# 500#		53 ms 48	0 <sup>+</sup>	17 17Wu04 TD	2017	$\beta^- = 100$
<sup>154</sup> La	-41530# 300#		161 ms 15	2 <sup>-</sup> #	17 17Wu04 TD	2017	$\beta^- = 100; \beta^- n \text{ ?}; \beta^- 2n \text{ ?}$
<sup>154</sup> Ce	-52220# 200#		722 ms 14	0 <sup>+</sup>	17 17Wu04 TD	1994	$\beta^- = 100; \beta^- n \text{ ?}$
<sup>154</sup> Pr	-57860 100		2.30 s 0.09	(3 <sup>+</sup> )	09 17Wu04 T	1988	$\beta^- = 100; \beta^- n \text{ ?}$
<sup>154</sup> Nd	-65579.6 1.0		25.9 s 0.2	0 <sup>+</sup>	09 1970	1970	$\beta^- = 100$
<sup>154</sup> Nd <sup>m</sup>	-64281.7 1.1	1297.9 0.4	3.2 $\mu$ s 0.3	(4 <sup>-</sup> )	09 09Si21 ETJ	1970	IT=100
<sup>154</sup> Pm	-68267 25		2.68 m 0.07	(4 <sup>+</sup> )	09 12So10 J	1958	$\beta^- = 100$
<sup>154</sup> Pm <sup>m</sup>	-68490 40	-230 50	IT* 1.73 m 0.10	(1 <sup>-</sup> )	09 12So10 J	1958	$\beta^- = 100$
<sup>154</sup> Sm	-72455.6 1.3		STABLE >2.3Ey	0 <sup>+</sup>	09	1933	IS=22.74 14; $2\beta^- \text{ ?}$
<sup>154</sup> Eu	-71738.4 1.2		8.592 y 0.003	3 <sup>-*</sup>	09 FGK209 T	1947	$\beta^- = 99.982 12; \epsilon = 0.018 12$
<sup>154</sup> Eu <sup>m</sup>	-71670.2 1.2	68.1702 0.0004	2.2 $\mu$ s 0.1	2 <sup>+</sup>	09	1964	IT=100
<sup>154</sup> Eu <sup>m</sup>	-71593.1 1.2	145.3 0.3	46.3 m 0.4	(8 <sup>-</sup> )	09	1975	IT=100
<sup>154</sup> Gd	-73706.4 1.0		STABLE	0 <sup>+</sup>	09	1938	IS=2.18 2
<sup>154</sup> Tb	-70160 50		9.994 h 0.039	3 <sup>-*</sup>	09 09Gy01 T	1972	$\beta^+ = 100; \beta^- \text{ ?}$
<sup>154</sup> Tb <sup>m</sup>	-70030# 70#	130# 50#	* & 21.5 h 0.4	0 <sup>-*</sup>	09	1950	$\beta^+ \approx 100; IT \text{ ?}; \beta^- \text{ ?}$
<sup>154</sup> Tb <sup>n</sup>	-69960# 160#	200# 150#	* 22.7 h 0.5	7 <sup>-</sup>	09	1972	$\beta^+ \approx 100; IT \text{ ?}$
<sup>154</sup> Tb <sup>p</sup>	-69760# 160#	405# 150#	513 ns 4.2		09	1982	IT=100
<sup>154</sup> Dy	-70394 7		3.0 My 1.5	0 <sup>+</sup>	09	1961	$\alpha = 100; 2\beta^+ \text{ ?}$
<sup>154</sup> Ho	-64639 8		11.76 m 0.19	2 <sup>-*</sup>	09	1966	$\beta^+ = 99.981 5; \alpha = 0.019 5$
<sup>154</sup> Ho <sup>m</sup>	-64397 27	243 28	AD 3.10 m 0.14	8 <sup>+</sup> *	09	1968	$\beta^+ = 100; \alpha < 0.001; IT \approx 0$
<sup>154</sup> Er	-62605 5		3.73 m 0.09	0 <sup>+</sup>	09	1963	$\beta^+ \approx 100; \alpha = 0.47 13$
<sup>154</sup> Tm	-54427 14		* 8.1 s 0.3	(2) <sup>-</sup>	09	1964	$\alpha = 54 5; \beta^+ = 46 5$
<sup>154</sup> Tm <sup>m</sup>	-54350 50	70 50	BD* 3.30 s 0.07	(9) <sup>+</sup>	09	1964	$\alpha = 58 5; \beta^+ = 42 5; IT \text{ ?}$
<sup>154</sup> Yb	-49932 17		409 ms 2	0 <sup>+</sup>	09	1964	$\alpha = 92.6 12; \beta^+ = 7.4 12$
<sup>154</sup> Lu	-39670# 200#		1# s	(2) <sup>-</sup>	09	1981	$\beta^+ \text{ ?}; \alpha \text{ ?}$
<sup>154</sup> Lu <sup>m</sup>	-39600# 200#	62 12	AD 1.12 s 0.08	(9 <sup>+</sup> )	09 88Vi02 D	1981	$\beta^+ \approx 100; \beta^+ p = \text{?}; \beta^+ \alpha = \text{?}; \alpha \text{ ?}$
<sup>154</sup> Lu <sup>n</sup>	-36950# 220#	2724# 100#	35 $\mu$ s 3	(17 <sup>+</sup> )	09	1990	IT=100
<sup>154</sup> Hf	-32730# 300#		2 s 1	0 <sup>+</sup>	09	1981	$\beta^+ \approx 100; \alpha \approx 0$
<sup>154</sup> Hf <sup>m</sup>	-30010# 300#	2721# 50#	9 $\mu$ s 4	(10 <sup>+</sup> )	09	1989	IT=100
* <sup>154</sup> La	T : other 20Wu04=221(89)						
* <sup>154</sup> Pr	T : average 17Wu04=2.29(0.20) 88Ka16=2.3(0.1)						
* <sup>154</sup> Nd <sup>m</sup>	E : from a least-squares fit to gamma-ray energies in 09Si21						
* <sup>154</sup> Nd <sup>m</sup>	I : other Ensdf2009 quotes this isomer twice: 233.2+x keV (1.3 us) and						
* <sup>154</sup> Nd <sup>m</sup>	I : 1349 keV (5-, >1 us); not trusted						
* <sup>154</sup> Sm	T : 2v- $\beta\beta$ to 2+ from 96De60						
* <sup>154</sup> Tb	J : 70Ad09=3; conf p3/2[411]n3/2[521], K=3- and GM rule						
* <sup>154</sup> Tb <sup>p</sup>	E : 82Be46=53.9,60.4 abd 90.1-keV gammas show 500 ns half-life; assumed by						
* <sup>154</sup> Tb <sup>p</sup>	E : Nubase above <sup>154</sup> Tb <sup>n</sup> since the level is populated in the						
* <sup>154</sup> Tb <sup>p</sup>	E : ( <sup>11</sup> B,5n) reaction that favors high spin						
* <sup>154</sup> Lu <sup>m</sup>	D : % $\beta^+ p$ and % $\beta^+ \alpha$ modes observed in 88Vi02; $\beta^+ p$ confirmed in 90Sh.A						
* <sup>154</sup> Lu <sup>n</sup>	E : 2431.3 + 130.4 + z keV above <sup>154</sup> Lu <sup>m</sup> ; z=100#(100#) keV						
* <sup>154</sup> Hf <sup>m</sup>	E : 93Mc03=2671+x keV; x=50#(50#) keV by Nubase						
<sup>155</sup> La	-37930# 400#		101 ms 28	1/2 <sup>+</sup> #	19 17Wu04 T	2016	$\beta^- = 100; \beta^- n \text{ ?}; \beta^- 2n \text{ ?}$
<sup>155</sup> Ce	-47780# 300#		313 ms 7	5/2 <sup>-</sup> #	19	1994	$\beta^- = 100; \beta^- n \text{ ?}$
<sup>155</sup> Pr	-55415 17		1.47 s 0.3	3/2 <sup>-</sup> #	19	1992	$\beta^- = 100; \beta^- n \text{ ?}$
<sup>155</sup> Nd	-62284 9		8.9 s 0.2	(3/2 <sup>-</sup> )	19	1986	$\beta^- = 100$
<sup>155</sup> Pm	-66940 5		41.5 s 0.2	(5/2 <sup>-</sup> )	19	1982	$\beta^- = 100$



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
* <sup>156</sup> Nd	T : other: 17Wu04=5.2(1.4)						**	
* <sup>156</sup> Pm	T : unweighed average 16Ko.A=27.78(0.07) 87Gr12=26.70(0.10);						**	
* <sup>156</sup> Pm	T : Birge ratio=8.85						**	
* <sup>156</sup> Pm <sup>m</sup>	E : other 20Or03=152.2(2.7) (PI-ICR)						**	
* <sup>156</sup> Sm <sup>m</sup>	T : other (recent) 09Si21=186(44)						**	
* <sup>156</sup> Eu	J : 90A134=0						**	
* <sup>156</sup> Tb <sup>n</sup>	E : from 49.630+x keV; x=50#(50#) keV estimated by Nubase						**	
* <sup>156</sup> Dy	T : the lower limit is for $\alpha$ decay						**	
* <sup>156</sup> Ho <sup>m</sup>	E : uncertainty estimated by Nubase						**	
* <sup>156</sup> Tm <sup>m</sup>	E : 203.6 keV above unknown level						**	
* <sup>156</sup> Lu <sup>m</sup>	E : 18Le10=2601.0(1.4) keV above <sup>156</sup> Lu <sup>m</sup>						**	
* <sup>156</sup> Hf <sup>m</sup>	T : average 96Pa01=520(10) 81Ho.A=444(17)						**	
<sup>157</sup> La	-29070#	300#	30# ms >550ns	1/2 <sup>+</sup> #	18Sh11 I	2018	$\beta^- ?; \beta^- n ?$	
<sup>157</sup> Ce	-39930#	400#	175 ms 41	7/2 <sup>+</sup> #	17 17Wu04 TD	2017	$\beta^- =100; \beta^- n ?$	
<sup>157</sup> Pr	-48435	3	307 ms 21	3/2 <sup>-</sup> #	17 17Wu04 TD	2017	$\beta^- =100; \beta^- n ?$	
<sup>157</sup> Nd	-56494.1	2.1	1.15 s 0.03	5/2 <sup>-</sup> #	17 17Wu04 TD	1992	$\beta^- =100$	
<sup>157</sup> Pm	-62297	7	10.56 s 0.10	(5/2 <sup>-</sup> )	16	1987	$\beta^- =100$	
<sup>157</sup> Sm	-66678	4	8.03 m 0.07	3/2 <sup>-</sup> #	16	1973	$\beta^- =100$	
<sup>157</sup> Eu	-69459	4	15.18 h 0.03	5/2 <sup>+</sup> *	16	1951	$\beta^- =100$	
<sup>157</sup> Gd	-70823.9	1.0	STABLE	3/2 <sup>-</sup> *	16	1933	IS=15.65 4	
<sup>157</sup> Gd <sup>m</sup>	-70760.0	1.0	63.916 0.005	460 ns 40	5/2 <sup>+</sup>	16	IT=100	
<sup>157</sup> Gd <sup>m</sup>	-70397.4	1.0	426.539 0.023	18.5 $\mu$ s 2.3	11/2 <sup>-</sup>	16	IT=100	
<sup>157</sup> Tb	-70763.8	1.0	71 y 7	3/2 <sup>+</sup>	16	1960	$\epsilon =100$	
<sup>157</sup> Dy	-69425	5	8.14 h 0.04	3/2 <sup>-</sup> *	16	1953	$\beta^+ =100$	
<sup>157</sup> Dy <sup>m</sup>	-69263	5	161.99 0.03	1.3 $\mu$ s 0.2	9/2 <sup>+</sup>	16	IT=100	
<sup>157</sup> Dy <sup>n</sup>	-69226	5	199.38 0.07	21.6 ms 1.6	11/2 <sup>-</sup>	16	IT=100	
<sup>157</sup> Ho	-66833	23	12.6 m 0.2	7/2 <sup>-</sup> *	16	1966	$\beta^+ =100$	
<sup>157</sup> Er	-63414	27	18.65 m 0.10	3/2 <sup>-</sup> *	16	1966	$\beta^+ =100$	
<sup>157</sup> Er <sup>m</sup>	-63259	27	155.4 0.3	76 ms 6	9/2 <sup>+</sup>	16	IT=100	
<sup>157</sup> Tm	-58709	28	3.63 m 0.09	1/2 <sup>+</sup> *	16 96By.A D	1974	$\beta^+ =100; \alpha =7.5e-4 25$	
<sup>157</sup> Tm <sup>m</sup>	-58610#	60#	100# 50# *	1.6 s	7/2 <sup>-</sup> #	08VaZV TJ	2008	$\beta^+ ?; IT ?$
<sup>157</sup> Yb	-53420	11	38.6 s 1.0	7/2 <sup>-</sup> *	16	1970	$\beta^+ \approx 100; \alpha = ?$	
<sup>157</sup> Lu	-46440	12	7.7 s 2.0	(1/2 <sup>+</sup> )	16	1977	$\beta^+ ?; \alpha = ?$	
<sup>157</sup> Lu <sup>m</sup>	-46419	12	20.9 2.0 AD	4.79 s 0.12	(11/2 <sup>-</sup> )	16	1972	$\beta^+ =92.3 19; \alpha =7.7 19$
<sup>157</sup> Hf	-38860#	200#	115 ms 1	7/2 <sup>-</sup>	16	1965	$\alpha =94 4; \beta^+ =14 4$	
<sup>157</sup> Ta	-29600	150	10.1 ms 0.4	1/2 <sup>+</sup>	16	1979	$\alpha =96.6 12; p =3.4 12; \beta^+ ?$	
<sup>157</sup> Ta <sup>m</sup>	-29570	150	22 5 AD	4.3 ms 0.1	11/2 <sup>-</sup>	16	1996	$\alpha \approx 100; \beta^+ ?; p =0$
<sup>157</sup> Ta <sup>n</sup>	-28000	150	1593 9 AD	1.7 ms 0.1	25/2 <sup>-</sup> #	16	1996	$\alpha =100$
<sup>157</sup> W	-19690#	400#	275 ms 40	(7/2 <sup>-</sup> )	16 10Bi03 D	2010	$\beta^+ =100; \alpha =0$	
<sup>157</sup> W <sup>p</sup>	-19370#	400#	320 30 AD	(9/2 <sup>-</sup> )	16	2010	IT ?	
* <sup>157</sup> Pr	T : symmetrized from 17Wu04=295(+29-11)						**	
* <sup>157</sup> Lu	T : unweighted average 91To09=5.7(0.5) 91Le15,92Po14=9.6(0.8);						**	
* <sup>157</sup> Lu	T : Birge ratio=4.13						**	
* <sup>157</sup> Lu <sup>m</sup>	D : % $\alpha$ average 91To09=18(5) 79Ho10=6(2); Birge ratio=2.23						**	
* <sup>157</sup> Hf	J : favored $\alpha$ decay to J=7/2- gs in <sup>153</sup> Yb						**	
<sup>158</sup> Ce	-36540#	400#	99 ms 93	0 <sup>+</sup>	17	2016	$\beta^- =100; \beta^- n ?$	
<sup>158</sup> Pr	-44150#	300#	181 ms 14	5 <sup>-</sup> #	17	2016	$\beta^- =100; \beta^- n ?$	
<sup>158</sup> Nd	-53835.1	1.3	810 ms 30	0 <sup>+</sup>	17 17Wu04 TD	1992	$\beta^- =100$	
<sup>158</sup> Nd <sup>m</sup>	-52187.0	1.9	339 ns 20	(6 <sup>-</sup> )	17	2016	IT=100	
<sup>158</sup> Pm	-59106.1	0.9	4.8 s 0.5	(0 <sup>+</sup> , 1 <sup>+</sup> )#	17	1987	$\beta^- =100$	
<sup>158</sup> Pm <sup>m</sup>	-58960#	50#	150# 50#	> 16 $\mu$ s	5 <sup>+</sup> #	17 15YoZX EDT2015	IT=?; $\beta^- ?$	
<sup>158</sup> Sm	-65252	5	5.30 m 0.03	0 <sup>+</sup>	17	1970	$\beta^- =100$	
<sup>158</sup> Eu	-67270.5	2.0	45.9 m 0.2	1 <sup>-</sup> *	17	1951	$\beta^- =100$	
<sup>158</sup> Gd	-70690.0	1.0	STABLE	0 <sup>+</sup>	17	1933	IS=24.84 8	
<sup>158</sup> Tb	-69470.9	1.3	180 y 11	3 <sup>-</sup> *	04	1957	$\beta^+ =83.4 7; \beta^- =16.6 7$	
<sup>158</sup> Tb <sup>m</sup>	-69360.6	1.8	110.3 1.2	10.70 s 0.17	0 <sup>-</sup>	17	1957	IT $\approx$ 100; $\beta^- ?; \beta^+ ?$
<sup>158</sup> Tb <sup>n</sup>	-69082.5	1.3	388.39 0.11	400 $\mu$ s 40	7 <sup>-</sup>	17	1961	IT=100
<sup>158</sup> Dy	-70407.2	2.3	STABLE	0 <sup>+</sup>	17	1938	IS=0.095 3; $\alpha ?; \beta^+ ?$	
<sup>158</sup> Ho	-66187	27	11.3 m 0.4	5 <sup>+</sup> *	17	1961	$\beta^+ \approx 100; \alpha ?$	

Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)

Nuclide	Mass excess (keV)	Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{158}\text{Ho}^m$	-66120	27	67.20	0.01	28 m 2	$2^-*$	17	1960	IT≈91 6;β <sup>+</sup> ≈9 6	*
$^{158}\text{Ho}^n$	-66095	27	91.595	0.012	140 ns 25	$(2^-)$	17	2005	IT=100	*
$^{158}\text{Ho}^p$	-66010#	80#	180#	70#	21.3 m 2.3	$(9^+)$	17	1970	β <sup>+</sup> ≈100;IT ?	
$^{158}\text{Er}$	-65304	25			2.29 h 0.06	$0^+$	17	1961	ε=100	
$^{158}\text{Tm}$	-58703	25		*	3.98 m 0.06	$2^-*$	17	1970	β <sup>+</sup> =100	
$^{158}\text{Tm}^m$	-58600#	60#	100#	50#	~ 20 s	$5^-#$	17 81Dr07	IT 1981	IT ?;ε ?	*
$^{158}\text{Yb}$	-56010	8			1.49 m 0.13	$0^+$	17	1967	β <sup>+</sup> ≈100;α≈0.0021 12	
$^{158}\text{Lu}$	-47212	15			10.6 s 0.3	$(2)^-$	17	1979	β <sup>+</sup> =99.09 20;α=0.91 20	
$^{158}\text{Hf}$	-42102	17			2.85 s 0.07	$0^+$	17	1965	β <sup>+</sup> =55.7 19;α=44.3 19	
$^{158}\text{Ta}$	-31120#	200#			49 ms 4	$(2)^-$	17 97Da07	TD 1979	α≈100;β <sup>+</sup> ?	*
$^{158}\text{Ta}^m$	-30980#	200#	141	11	AD	$(9)^+$	17 97Da07	ETJ 1979	α=95 5;β <sup>+</sup> ?;IT ?	*
$^{158}\text{Ta}^n$	-28310#	200#	2808	16		$(19)^-$	17	2014	IT=98.6 2;α=1.4 2	*
$^{158}\text{W}$	-23690#	300#			1.43 ms 0.18	$0^+$	17 19Hi06	T 1981	α=100	*
$^{158}\text{W}^m$	-21800#	300#	1889	8	AD	$(8^+)$	17	1995	α=100;IT ?	
* $^{158}\text{Nd}$	T : symmetrized from 17Wu04=820(+15-36)									**
* $^{158}\text{Pm}^m$	E : 15YoZX=121+x (121-keV gamma ray below the isomer); x=30#(50#) by Nubase									**
* $^{158}\text{Eu}$	J : 90A134=1									**
* $^{158}\text{Tb}$	J : 68Ea04=3									**
* $^{158}\text{Ho}^m$	D : %IT from Ensdf2017>81									**
* $^{158}\text{Ho}^n$	J : E1 from 1+; not fed directly in $^{158}\text{Er}$ (J=0+) β <sup>+</sup> decay									**
* $^{158}\text{Tm}^m$	I : 20 s activity, following observation of gammas in $^{158}\text{Er}$ ε decay									**
* $^{158}\text{Tm}^m$	I : in 81Dr07, is adopted. Note, that 20 ns appears in the level scheme									**
* $^{158}\text{Tm}^m$	I : in Fig. 2 (81Dr07), which seems to be a misprint. This is a spin-trap									**
* $^{158}\text{Tm}^m$	I : isomer and the suggested 20 ns half-life in Ensdf17 is unrealistic.									**
* $^{158}\text{Tm}^m$	I : The configuration is the same as that for the ground state,									**
* $^{158}\text{Tm}^m$	I : p7/2[404] n3/2[521], but K=5-. 75Ag01 also cannot rule out the									**
* $^{158}\text{Tm}^m$	I : existence of two ε-decaying states									**
* $^{158}\text{Ta}$	T : average 97Da07=72(12) 96Pa01=46(4); Birge ratio B=2.06									**
* $^{158}\text{Ta}^m$	T : average 97Da07=37.7(1.5) 96Pa01=35(1) 79Ho10=36.8(1.6)									**
* $^{158}\text{Ta}^n$	E : from Ensdf2017=2664.5(0.4) keV above $^{158}\text{Ta}^m$									**
* $^{158}\text{W}$	T : average 19Hi06=1.9(+1.2-0.6) 00Ma95=1.5(0.2) 96Pa01=0.9(+0.4-0.3)									**
$^{159}\text{Ce}$	-31340#	500#				$1/2^-#$			β <sup>-</sup> ?;β <sup>-</sup> n ?	
$^{159}\text{Pr}$	-40770#	400#			134 ms 43	$3/2^-#$	17 17Wu04	TD 2017	β <sup>-</sup> =100;β <sup>-</sup> n ?	
$^{159}\text{Nd}$	-49724	30			500 ms 30	$7/2^+#$	17 17Wu04	TD 2012	β <sup>-</sup> =100;β <sup>-</sup> n ?	*
$^{159}\text{Pm}$	-56554	10			1.49 s 0.13	$(5/2^-)$	12 17Wu04	T 1998	β <sup>-</sup> =100	*
$^{159}\text{Pm}^m$	-55089	10	1465.0	0.5	4.42 μs 0.17	$17/2^+#$	15YoZX	ETD2015	IT=100	*
$^{159}\text{Sm}$	-62208	6			11.37 s 0.15	$5/2^-$	12	1986	β <sup>-</sup> =100	
$^{159}\text{Sm}^m$	-60932	6	1276.5	0.8	116 ns 8	$(15/2^+)$	12 17Pa25	EJ 2009	IT=100	*
$^{159}\text{Eu}$	-66043	4			18.1 m 0.1	$5/2^+*$	12	1961	β <sup>-</sup> =100	*
$^{159}\text{Gd}$	-68561.9	1.0			18.479 h 0.004	$3/2^-*$	12	1949	β <sup>-</sup> =100	
$^{159}\text{Tb}$	-69532.6	1.1			STABLE	$3/2^+*$	12 12Vi10	J 1933	IS=100	
$^{159}\text{Dy}$	-69167.2	1.4			144.4 d 0.2	$3/2^-*$	12	1951	ε=100	
$^{159}\text{Dy}^m$	-68814.4	1.4	352.77	0.14	122 μs 3	$11/2^-$	12	1965	IT=100	
$^{159}\text{Ho}$	-67330	3			33.05 m 0.11	$7/2^-*$	12	1958	β <sup>+</sup> =100	
$^{159}\text{Ho}^m$	-67124	3	205.91	0.05	8.30 s 0.08	$1/2^+$	12	1966	IT=100	
$^{159}\text{Er}$	-64561	4			36 m 1	$3/2^-*$	12	1962	β <sup>+</sup> =100	
$^{159}\text{Er}^m$	-64378	4	182.602	0.024	337 ns 14	$9/2^+$	12	1971	IT=100	
$^{159}\text{Er}^n$	-64132	4	429.05	0.03	590 ns 60	$11/2^-$	12	1971	IT=100	
$^{159}\text{Tm}$	-60570	28			9.13 m 0.16	$5/2^+*$	12	1971	β <sup>+</sup> =100	
$^{159}\text{Yb}$	-55834	18			1.67 m 0.09	$5/2^-*$	12	1975	β <sup>+</sup> =100	
$^{159}\text{Lu}$	-49710	40		*	12.1 s 1.0	$1/2^+$	12 FGK12a	J 1980	β <sup>+</sup> ≈100;α ?	*
$^{159}\text{Lu}^m$	-49610#	90#	100#	80#	10# s	$11/2^-#$			β <sup>+</sup> ?;IT ?;α ?	
$^{159}\text{Hf}$	-42853	17			5.20 s 0.10	$7/2^-$	12 96Pa01	T 1973	β <sup>+</sup> =65 7;α=35 7	
$^{159}\text{Ta}$	-34439	20			1.04 s 0.09	$1/2^+$	12 97Da07	T 1979	β <sup>+</sup> =66 5;α=34 5	*
$^{159}\text{Ta}^m$	-34375	19	64	5	AD	$11/2^-$	12	1994	α=55 1;β <sup>+</sup> =45 1	
$^{159}\text{W}$	-25430#	300#			560 ms 60	$7/2^-#$	12 96Pa01	TD 1981	α≈100;β <sup>+</sup> ?	*
$^{159}\text{Re}$	-14810#	310#			40# μs	$1/2^+*$	12	2006	p ?;α ?	
$^{159}\text{Re}^m$	-14600#	300#	210#	50#	20 μs 4	$11/2^-$	12 07Pa27	TD 2006	p=92.5 35;α=7.5 35	*
* $^{159}\text{Nd}$	T : symmetrized from 17Wu04=485(+39-20)									**
* $^{159}\text{Pm}$	T : average 17Wu04=1.48(0.18) 05Ic02=1.5(0.2), supersedes 01AsZY									**
* $^{159}\text{Pm}^m$	J : 99.6-keV gamma to (15/2-), conf=p5/2[532] n(5/2[523],7/2[633]), K=17/2+									**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	J <sup>π</sup>	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>159</sup> Sm <sup>m</sup>	E : average 17Pa25=1275.9(1.4) 09Ur04=1276.8(1.0)						**
* <sup>159</sup> Sm <sup>m</sup>	T : from 09Ur04; other 17Pa25=50(17)						**
* <sup>159</sup> Eu	J : 90A134=5/2						**
* <sup>159</sup> Lu	J : favored α decay from <sup>163</sup> Ta (J=1/2+)						**
* <sup>159</sup> Ta	T : average 97Da07=0.83(0.18) 96Pa01=1.10(0.10)						**
* <sup>159</sup> W	T : others 19Hi06=10.3(+2.1-1.5) 81Ho10=7.3(2.7)						**
* <sup>159</sup> Re <sup>m</sup>	T : average 07Pa27=16(9) 06Jo10=21(4)						**
<sup>160</sup> Pr	-36200#	400#	170 ms 140	1 <sup>+</sup> #	17 17Wu04	TD 2017	β <sup>-</sup> =100;β <sup>-</sup> <sub>n</sub> ?
<sup>160</sup> Nd	-46720	50	439 ms 37	0 <sup>+</sup>	17 17Wu04	TD 1985	β <sup>-</sup> =100;β <sup>-</sup> <sub>n</sub> ?
<sup>160</sup> Nd <sup>m</sup>	-45610	50	1107.9 0.9	1.63 μs 0.21	(4 <sup>-</sup> )	17 16Id02	ETJ 2016 IT=100
<sup>160</sup> Pm	-52894.6	2.0	725 ms 57	6 <sup>-</sup> #	17 17Wu04	TD 2012	β <sup>-</sup> =100;β <sup>-</sup> <sub>n</sub> ?
<sup>160</sup> Pm <sup>m</sup>	-52704	11	> 700 ms	1 <sup>-</sup> #	20Or03	EJT 2020	β <sup>-</sup> ?; IT ?;β <sup>-</sup> <sub>n</sub> ?
<sup>160</sup> Sm	-60233.2	2.0	9.6 s 0.3	0 <sup>+</sup>	05	1986	β <sup>-</sup> =100
<sup>160</sup> Sm <sup>m</sup>	-58871.9	2.0	1361.3 0.4	120 ns 46	(5 <sup>-</sup> )	09Si21	ETJ 2009 IT=100
<sup>160</sup> Sm <sup>m</sup>	-57475.9	2.0	2757.3 0.4	1.8 μs 0.4	(11 <sup>+</sup> )	16Pa01	ETJ 2016 IT=100
<sup>160</sup> Eu	-63493.4	0.9	42.6 s 0.5	(5 <sup>-</sup> )	05 18Ha19	TJ 1973	β <sup>-</sup> =100
<sup>160</sup> Eu <sup>m</sup>	-63400.4	0.8	93.0 1.2 MD	30.8 s 0.5	(1 <sup>-</sup> )	05 18Ha19	ETJ 2016 β <sup>-</sup> =100
<sup>160</sup> Gd	-67942.1	1.1	STABLE	>31Ey	0 <sup>+</sup>	05 01Da22	T 1933 IS=21.86 3;2β <sup>-</sup> ?
<sup>160</sup> Tb	-67836.5	1.1	72.3 d 0.2	3 <sup>-*</sup>	05	1943	β <sup>-</sup> =100
<sup>160</sup> Dy	-69672.4	0.7	STABLE	0 <sup>+</sup>	05	1938	IS=2.329 18
<sup>160</sup> Ho	-66382	15	25.6 m 0.3	5 <sup>+</sup> *	05	1950	β <sup>+</sup> =100
<sup>160</sup> Ho <sup>m</sup>	-66322	15	59.98 0.03	5.02 h 0.05	2 <sup>-*</sup>	05	1955 IT=73 3;β <sup>+</sup> =27 3
<sup>160</sup> Ho <sup>n</sup>	-66185	22	197 16	~ 3 s	(9 <sup>+</sup> )	05 88Bh05	TD 1988 IT=100
<sup>160</sup> Er	-66064	24	28.58 h 0.09	0 <sup>+</sup>	05	1954	ε=100
<sup>160</sup> Tm	-60300	30	9.4 m 0.3	1 <sup>-*</sup>	05	1970	β <sup>+</sup> =100
<sup>160</sup> Tm <sup>m</sup>	-60230	30	67 14	74.5 s 1.5	(5 <sup>+</sup> )	05	1983 IT=85 5;β <sup>+</sup> =15 5
<sup>160</sup> Tm <sup>n</sup>	-60090#	60#	215# 52#	~ 200 ns	(8)	05	1986 IT=100
<sup>160</sup> Yb	-58163	5	4.8 m 0.2	0 <sup>+</sup>	05	1967	β <sup>+</sup> =100
<sup>160</sup> Lu	-50270	60	36.1 s 0.3	2 <sup>-</sup> #	05	1979	β <sup>+</sup> =100;α ?
<sup>160</sup> Lu <sup>m</sup>	-50270#	120#	0# 100#	40 s 1	05	1980	β <sup>+</sup> ≈100;α ?
<sup>160</sup> Hf	-45939	10	13.6 s 0.2	0 <sup>+</sup>	05	1973	β <sup>+</sup> =99.3 2;α=0.7 2
<sup>160</sup> Ta	-35820	50	* & 1.70 s 0.20	(2 <sup>-</sup> )	05 96Pa01	TD 1979	β <sup>+</sup> ?;α=?
<sup>160</sup> Ta <sup>m</sup>	-35710	240	110 250	* & 1.55 s 0.04	(9,10) <sup>+</sup>	05 96Pa01	TD 1979 β <sup>+</sup> ?;α=?
<sup>160</sup> W	-29330	150	90 ms 5	0 <sup>+</sup>	05 96Pa01	TD 1979	α=87 8;β <sup>+</sup> ?
<sup>160</sup> Re	-16880#	300#	611 μs 7	(4 <sup>-</sup> )	05 11Da12	TJD 1992	p=89 1;α=11 1
<sup>160</sup> Re <sup>m</sup>	-16700#	300#	177 15	2.8 μs 0.1	(9 <sup>+</sup> )	11Da01	JT 2011 IT=100
* <sup>160</sup> Pm	T : the value of 17Wu04 probably includes both the gs and isomer						**
* <sup>160</sup> Gd	T : value quoted at 68% CL						**
* <sup>160</sup> Ho <sup>n</sup>	E : from 169.61 keV + x with x<55 keV from Ensdf2005						**
* <sup>160</sup> Tm <sup>n</sup>	E : from 42.10+x keV above gs; x<50 keV from Ensdf2005						**
* <sup>160</sup> Tm <sup>m</sup>	E : 98.2 keV+x keV above <sup>160</sup> Tm <sup>m</sup> ; x=50#(50#) keV by Nubase						**
* <sup>160</sup> Ta	J : favored α decay to <sup>156</sup> Lu [J=(2-)]						**
* <sup>160</sup> Ta <sup>m</sup>	J : favored α decay to <sup>156</sup> Lu <sup>m</sup> [J=10+]						**
* <sup>160</sup> W	T : average 96Pa01=91(5) 81Ho10=81(15)						**
* <sup>160</sup> Re	J : other 92Pa05=(2-)						**
<sup>161</sup> Pr	-32490#	500#	90# ms >550ns	3/2 <sup>-</sup> #	18Fu08	I 2018	β <sup>-</sup> ?;β <sup>-</sup> <sub>n</sub> ?
<sup>161</sup> Nd	-42230#	400#	215 ms 76	1/2 <sup>-</sup> #	17 17Wu04	TD 2012	β <sup>-</sup> =100;β <sup>-</sup> <sub>n</sub> ?
<sup>161</sup> Pm	-50087	9	1.05 s 0.15	(5/2 <sup>-</sup> )	17 17Wu04	TD 2012	β <sup>-</sup> =100;β <sup>-</sup> <sub>n</sub> ?
<sup>161</sup> Pm <sup>m</sup>	-49121	9	0.89 μs 0.09	(13/2 <sup>+</sup> )	17 15YoZX	TJ 2015	IT=100
<sup>161</sup> Sm	-56672	7	4.8 s 0.4	7/2 <sup>+</sup> #	11	1998	β <sup>-</sup> =100
<sup>161</sup> Sm <sup>m</sup>	-55284	7	1388.1 0.6	2.6 μs 0.4	(17/2 <sup>-</sup> )	17Pa25	ETJ 2017 IT=100
<sup>161</sup> Eu	-61792	10	26.2 s 2.3	5/2 <sup>+</sup> #	11 17Wu04	T 1986	β <sup>-</sup> =100
<sup>161</sup> Gd	-65506.1	1.5	3.646 m 0.003	5/2 <sup>-</sup>	11 94It.A	T 1949	β <sup>-</sup> =100
<sup>161</sup> Tb	-67461.8	1.2	6.948 d 0.005	3/2 <sup>+</sup> *	11 FGK204	T 1949	β <sup>-</sup> =100
<sup>161</sup> Dy	-68055.5	0.7	STABLE	5/2 <sup>+</sup> *	11	1934	IS=18.889 42
<sup>161</sup> Dy <sup>m</sup>	-67569.9	0.7	485.56 0.16	760 ns 170	11/2 <sup>-</sup>	11 12Sw01	T 2012 IT=100
<sup>161</sup> Ho	-67196.3	2.2	2.48 h 0.05	7/2 <sup>-*</sup>	11	1954	ε=100
<sup>161</sup> Ho <sup>m</sup>	-66985.2	2.2	211.15 0.03	6.76 s 0.07	1/2 <sup>+</sup>	11	1965 IT=100
<sup>161</sup> Er	-65201	9	3.21 h 0.03	3/2 <sup>-*</sup>	11	1954	β <sup>+</sup> =100





Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{163}\text{Eu}$	-56573.8	0.9	7.7 s 0.4	5/2 <sup>+</sup> #	10 08Os02	T 2007	$\beta^- = 100$
$^{163}\text{Eu}^m$	-55609.3	1.0	911 ns 24	(13/2 <sup>-</sup> )	17Pa25	TJE 2017	IT=100
$^{163}\text{Gd}$	-61388.6	0.8	68 s 3	7/2 <sup>+</sup>	10 20Za04	J 1982	$\beta^- = 100$
$^{163}\text{Gd}^m$	-61250.4	0.8	23.5 s 1.0	1/2 <sup>-</sup>	14Ha38	TD 2014	IT=?; $\beta^-$ ?
$^{163}\text{Tb}$	-64596	4	19.5 m 0.3	3/2 <sup>+</sup>	10	1966	$\beta^- = 100$
$^{163}\text{Dy}$	-66380.9	0.7	STABLE	5/2 <sup>-</sup> *	10	1934	IS=24.896 42
$^{163}\text{Ho}$	-66378.0	0.7	4.570 ky 0.025	7/2 <sup>-</sup> *	10	1957	$\varepsilon = 100$
$^{163}\text{Ho}^m$	-66080.1	0.7	1.09 s 0.03	1/2 <sup>+</sup>	10	1957	IT=100
$^{163}\text{Ho}^n$	-64268.6	0.8	800 ns 150	(23/2 <sup>+</sup> )	12Sw01	ETJ 2012	IT=100
$^{163}\text{Er}$	-65167	5	75.0 m 0.4	5/2 <sup>-</sup> *	10	1953	$\beta^+ = 100$
$^{163}\text{Er}^m$	-64722	5	445.5 ns 100	(11/2 <sup>-</sup> )	10	1974	IT=100
$^{163}\text{Tm}$	-62728	6	1.810 h 0.005	1/2 <sup>+</sup> *	10	1959	$\beta^+ = 100$
$^{163}\text{Tm}^m$	-62641	6	380 ns 30	(7/2 <sup>-</sup> )	10	1975	IT=100
$^{163}\text{Yb}$	-59294	15	11.05 m 0.35	3/2 <sup>-</sup> *	10	1967	$\beta^+ = 100$
$^{163}\text{Lu}$	-54791	28	3.97 m 0.13	1/2 <sup>+</sup> *	10	1979	$\beta^+ = 100$
$^{163}\text{Hf}$	-49269	26	40.0 s 0.6	(5/2 <sup>-</sup> )	15	1982	$\beta^+ = 100$ ; $\alpha$ ?
$^{163}\text{Ta}$	-42530	40	10.6 s 1.8	1/2 <sup>+</sup>	10 FGK12a	J 1985	$\beta^+ \approx 100$ ; $\alpha$ ?
$^{163}\text{Ta}^m$	-42400#	40#	10# s	9/2 <sup>-</sup>	FGK12a	J	$\beta^+ ?$ ; $\alpha ?$ ; IT ?
$^{163}\text{W}$	-34910	60	2.63 s 0.09	7/2 <sup>-</sup>	10	1973	$\beta^+ ?$ ; $\alpha = 14.2$
$^{163}\text{W}^m$	-34430	60	154 ns 3	13/2 <sup>+</sup>	10	2010	IT=100
$^{163}\text{Re}$	-26002	19	390 ms 70	1/2 <sup>+</sup>	10	1979	$\beta^+ ?$ ; $\alpha = 32.3$
$^{163}\text{Re}^m$	-25882	19	214 ms 5	11/2 <sup>-</sup>	10	1979	$\alpha = 66.4$ ; $\beta^+ ?$
$^{163}\text{Os}$	-16340#	300#	5.7 ms 0.5	7/2 <sup>-</sup>	10 13Dr06	J 1981	$\alpha \approx 100$ ; $\beta^+ ?$
$^{163}\text{Ir}$	-5310#	400#		1/2 <sup>+</sup> #			p ?
* $^{163}\text{Pm}$	T : other 17Wu04=430(350)						**
* $^{163}\text{Sm}$	T : symmetrized from 17Wu04=1.23(+0.51-0.47)						**
* $^{163}\text{Eu}^m$	T : average 17Pa25=990(40) 17Yo01=869(29)						**
* $^{163}\text{Gd}^m$	J : 20Za04=1/2-						**
* $^{163}\text{Ho}$	T : other: 92Ju01=47(+5-4) d for q=66+ (bare ion)						**
* $^{163}\text{Ta}$	J : favored $\alpha$ decay from $^{167}\text{Re}^m$ (J=1/2+)						**
* $^{163}\text{Ta}^m$	J : favored $\alpha$ decay from $^{167}\text{Re}$ (J=9/2-)						**
* $^{163}\text{Os}$	T : average 19Hi06=6.2(+1.3-0.9) 96Bi07=5.5(0.6)						**
$^{164}\text{Pm}$	-38360#	400#	300# ms > 550ns	5 <sup>-</sup> #	18Fu08	I 2018	$\beta^- ?$ ; $\beta^-_n ?$
$^{164}\text{Sm}$	-47925	4	1.43 s 0.24	0 <sup>+</sup>	17	2012	$\beta^- = 100$ ; $\beta^-_n ?$
$^{164}\text{Sm}^m$	-46440	4	600 ns 140	(6 <sup>-</sup> )	17	2014	IT=100
$^{164}\text{Eu}$	-53232.1	2.1	4.16 s 0.19	3 <sup>-</sup> #	17 17Wu04	T 2007	$\beta^- = 100$
$^{164}\text{Gd}$	-59693.7	1.0	45 s 3	0 <sup>+</sup>	17	1988	$\beta^- = 100$
$^{164}\text{Gd}^m$	-58597.9	1.1	589 ns 18	(4 <sup>-</sup> )	17 18Ga18	T 2017	IT=100
$^{164}\text{Tb}$	-62105.0	1.9	3.0 m 0.1	(5 <sup>+</sup> )	17	1968	$\beta^- = 100$
$^{164}\text{Tb}^m$	-61960	12	2# m	2 <sup>+</sup> #	20Or03	EJ 2020	$\beta^- ?$ ; IT ?
$^{164}\text{Dy}$	-65967.6	0.7	STABLE	0 <sup>+</sup>	17	1934	IS=28.260 54
$^{164}\text{Ho}$	-64980.5	1.4	28.8 m 0.5	1 <sup>+</sup> *	17	1938	$\varepsilon = 61.1$ ; $\beta^+ = 39.1$
$^{164}\text{Ho}^m$	-64840.7	1.4	36.6 m 0.3	6 <sup>-</sup> *	17	1966	IT=100
$^{164}\text{Er}$	-65942.6	0.7	STABLE	0 <sup>+</sup>	17	1938	IS=1.601 3; $\alpha = 2$ ; $\beta^+ ?$
$^{164}\text{Tm}$	-61909	25	2.0 m 0.1	1 <sup>+</sup> *	17	1960	$\beta^+ = 100$ ; $\varepsilon = 61.1$ ; $e^+ = 39.1$
$^{164}\text{Tm}^m$	-61889	28	5.1 m 0.1	6 <sup>-</sup> *	17	1971	IT $\approx 80$ ; $\beta^+ \approx 20$
$^{164}\text{Yb}$	-61012	15	75.8 m 1.7	0 <sup>+</sup>	17	1960	$\varepsilon = 100$
$^{164}\text{Lu}$	-54642	28	3.14 m 0.03	1 <sup>-</sup> *	17	1977	$\beta^+ = 100$
$^{164}\text{Hf}$	-51818	16	111 s 8	0 <sup>+</sup>	17	1981	$\beta^+ = 100$
$^{164}\text{Ta}$	-43283	28	14.2 s 0.3	(3 <sup>+</sup> )	17	1982	$\beta^+ = 100$
$^{164}\text{W}$	-38236	10	6.3 s 0.2	0 <sup>+</sup>	17	1973	$\beta^+ = 96.2$ 12; $\alpha = 3.8$ 12
$^{164}\text{Re}$	-27470	50	719 ms 89	(2) <sup>-</sup>	17	1979	$\alpha = ?$ ; $\beta^+ ?$
$^{164}\text{Re}^m$	-27520	240	890 ms 130	(9, 10) <sup>+</sup>	17 09Ha42	TD 2009	$\beta^+ ?$ ; $\alpha = 3.1$
$^{164}\text{Os}$	-20420	150	21 ms 1	0 <sup>+</sup>	17	1981	$\alpha = 96.4$ ; $\beta^+ ?$
$^{164}\text{Ir}$	-7480#	320#	1# ms	2 <sup>-</sup> #	17		p ?; $\alpha ?$ ; $\beta^+ ?$
$^{164}\text{Ir}^m$	-7220#	300#	260# 100#	(9 <sup>+</sup> )	17 14Dr02	TD 2001	p=?; $\alpha = 4.2$ ; $\beta^+ ?$
* $^{164}\text{Eu}$	T : average 17Wu04=3.80(0.56) 08Os02=4.2(0.2)						**
* $^{164}\text{Gd}^m$	T : average 18Ga18=605(30) 17Yo01=580(23); other 17Pa25=530(100)						**
* $^{164}\text{Tm}^m$	E : from 87Dr07 < 40 keV						**
* $^{164}\text{Re}$	T : average 09Ha42=848(+140-105) 96Pa01=380(160) 81Ho10=880(240)						**
* $^{164}\text{Re}$	J : favored $\alpha$ decay to $^{160}\text{Ta}$ [J=(2)-]						**









**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>170</sup> Au <sup>m</sup>	T : symmetrized from 04Ke06=617(+50-40); other 02Ma61=570(+310-150)						
* <sup>170</sup> Au <sup>m</sup>	D : %p other 02Ma61=75(15)%						
* <sup>170</sup> Au <sup>m</sup>	J : favored $\alpha$ decay to <sup>166</sup> Re <sup>m</sup> [J=(9)+]						
* <sup>170</sup> Hg	T : symmetrized from 19Hi06=80(+400-40)						
<sup>171</sup> Gd	-36210# 500#		300# ms >550ns	9/2 <sup>+</sup> #	18Fu08 I	2018	$\beta^-$ ?; $\beta^-$ n ?
<sup>171</sup> Tb	-43770# 400#		1.23 s 0.10	3/2 <sup>+</sup> #	18 17Wu04 TD	2012	$\beta^-$ =100; $\beta^-$ n ?
<sup>171</sup> Dy	-50010# 200#		4.07 s 0.40	7/2 <sup>-</sup> #	18 17Wu04 TD	2012	$\beta^-$ =100
<sup>171</sup> Ho	-54520 600		53 s 2	7/2 <sup>-</sup> #	18	1989	$\beta^-$ =100
<sup>171</sup> Er	-57717.8 1.4		7.516 h 0.002	5/2 <sup>-</sup> *	18	1938	$\beta^-$ =100
<sup>171</sup> Er <sup>m</sup>	-57519.2 1.4	198.61 0.09	210 ns 10	1/2 <sup>-</sup>	18	1969	IT=100
<sup>171</sup> Tm	-59210.3 1.0		1.92 y 0.01	1/2 <sup>+</sup> *	18	1948	$\beta^-$ =100
<sup>171</sup> Tm <sup>m</sup>	-58785.3 1.0	424.9557 0.0015	2.60 $\mu$ s 0.02	7/2 <sup>-</sup>	18	1948	IT=100
<sup>171</sup> Tm <sup>n</sup>	-57535.9 1.0	1674.43 0.13	1.7 $\mu$ s 0.2	19/2 <sup>+</sup>	18	2009	IT=100
<sup>171</sup> Yb	-59306.818 0.013		STABLE	1/2 <sup>-</sup> *	18	1934	IS=14.086 140
<sup>171</sup> Yb <sup>m</sup>	-59211.536 0.013	95.282 0.002	5.25 ms 0.24	7/2 <sup>+</sup>	18	1968	IT=100
<sup>171</sup> Yb <sup>n</sup>	-59184.402 0.013	122.416 0.002	265 ns 20	5/2 <sup>-</sup>	18	1968	IT=100
<sup>171</sup> Lu	-57828.5 1.9		8.247 d 0.023	7/2 <sup>+</sup> *	18	1951	$\beta^-$ =100
<sup>171</sup> Lu <sup>m</sup>	-57757.4 1.9	71.13 0.08	79 s 2	1/2 <sup>-</sup> *	18	1965	IT=100
<sup>171</sup> Hf	-55431 29		12.1 h 0.4	7/2 <sup>+</sup> *	18	1951	$\beta^-$ =100
<sup>171</sup> Hf <sup>m</sup>	-55409 29	21.93 0.09	29.5 s 0.9	1/2 <sup>-</sup> *	18	1997	IT $\approx$ 100; $\beta^+$ ?
<sup>171</sup> Ta	-51720 28		23.3 m 0.3	(5/2 <sup>+</sup> )	18	1969	$\beta^+$ =100
<sup>171</sup> W	-47086 28		2.38 m 0.04	(5/2 <sup>-</sup> )	18	1983	$\beta^+$ =100
<sup>171</sup> Re	-41250 28		15.2 s 0.4	(9/2 <sup>-</sup> )	18	1987	$\beta^+$ =100
<sup>171</sup> Os	-34297 18		8.3 s 0.2	(5/2 <sup>-</sup> )	18	1972	$\beta^+$ ?; $\alpha$ =1.80 21
<sup>171</sup> Ir	-26410 40		3.1 s 0.3	1/2 <sup>+</sup>	18 11Ko.B TJ	1967	$\beta^+$ ?; $\alpha$ =15 2
<sup>171</sup> Ir <sup>m</sup>	-26250# 40#	164# 11#	1.47 s 0.06	(11/2 <sup>-</sup> )	18 11Ko.B T	1967	$\alpha$ =54 5; $\beta^+$ ?; p ?
<sup>171</sup> Pt	-17470 80		45.5 ms 2.5	7/2 <sup>-</sup>	18 10Sc02 JD	1981	$\alpha$ =86 3; $\beta^+$ ?
<sup>171</sup> Pt <sup>m</sup>	-17060 80	412.6 1.0	901 ns 9	13/2 <sup>+</sup>	18 FGK128 J	2010	IT=100
<sup>171</sup> Au	-7562 21		22.3 $\mu$ s 2.4	1/2 <sup>+</sup>	18 04Ke06 TJ	1997	p $\approx$ 100; $\alpha$ ?
<sup>171</sup> Au <sup>m</sup>	-7308 18	255 10 p	1.036 ms 0.016	11/2 <sup>-</sup>	18 04Ke06 TDJ	1996	$\alpha$ =60 6; p=40 6
<sup>171</sup> Hg	3340# 310#		70 $\mu$ s 30	3/2 <sup>-</sup> #	04 04Ke06 TD	2004	$\alpha$ $\approx$ 100; $\beta^+$ ?
* <sup>171</sup> Tb	T : symmetrized from 17Wu04=1.24(+0.09-0.10)						
* <sup>171</sup> Hf	J : 00Ye02=7/2						
* <sup>171</sup> Hf	J : 00Ye02=1/2						
* <sup>171</sup> Ta	T : Ensdf18 assign this lifetime to an excited state ( $J^\pi=5/2^-$ ) that is						
* <sup>171</sup> Ta	T : 31.2 keV above the proposed ground state ( $J^\pi=5/2^+$ )						
* <sup>171</sup> Ir	T : other 02Ro17=3.2(+1.3-0.7)						
* <sup>171</sup> Ir	D : % $\alpha$ from 13An10=15(2)						
* <sup>171</sup> Ir <sup>m</sup>	D : % $\alpha$ average 10An01=53(5)% 96Pa01=58(11)%						
* <sup>171</sup> Ir <sup>m</sup>	T : average 11Ko.B=1.50(0.07) 10An01=1.40(0.10)						
* <sup>171</sup> Pt	D : % $\alpha$ average 10Sc02=83(3) 04GoZZ=96(5)						
* <sup>171</sup> Pt <sup>m</sup>	J : M2 to 9/2- followed by M1 to 7/2- gs						
* <sup>171</sup> Au	T : average 04Ke06=22(+3-2) 99Po09=17(+9-5)						
* <sup>171</sup> Au	T : other 03Ba20=37(+7-5) conflicting, not used						
* <sup>171</sup> Au <sup>m</sup>	T : average 04Ke06=1.09(0.03) 03Ba20=1.014(0.019)						
* <sup>171</sup> Au <sup>m</sup>	D : %p average 04Ke06=34(4) 97Da07=46(4)						
* <sup>171</sup> Hg	T : symmetrized from 04Ke06=59(+36-16)						
<sup>172</sup> Gd	-32970# 300#		160# ms >550ns	0 <sup>+</sup> #	18Fu08 I	2018	$\beta^-$ ?; $\beta^-$ n ?
<sup>172</sup> Tb	-39690# 500#		760 ms 190	6 <sup>+</sup> #	17 17Wu04 TD	2012	$\beta^-$ =100; $\beta^-$ n ?
<sup>172</sup> Dy	-47760# 300#		3.4 s 0.2	0 <sup>+</sup>	13 16Wa19 TD	2012	$\beta^-$ =100
<sup>172</sup> Dy <sup>m</sup>	-46480# 300#	1278 1	710 ms 50	(8 <sup>-</sup> )	16Wa19 ETJ	2016	$\beta^-$ =19 3; IT=81 3
<sup>172</sup> Ho	-51480# 200#		25 s 3	0 <sup>+</sup> #	15	1991	$\beta^-$ =100
<sup>172</sup> Er	-56483 4		49.3 h 0.5	0 <sup>+</sup>	15	1956	$\beta^-$ =100
<sup>172</sup> Er <sup>m</sup>	-54982 4	1500.9 0.3	579 ns 62	(6 <sup>+</sup> )	15 10Dr02 ETJ	2006	IT=100
<sup>172</sup> Tm	-57374 5		63.6 h 0.3	2 <sup>-</sup>	15	1956	$\beta^-$ =100
<sup>172</sup> Tm <sup>m</sup>	-56898 5	476.2 0.2	132 $\mu$ s 7	(6 <sup>+</sup> )	15	2008	IT=100
<sup>172</sup> Yb	-59255.456 0.014		STABLE	0 <sup>+</sup>	95	1934	IS=21.686 130
<sup>172</sup> Yb <sup>m</sup>	-57705.03 0.06	1550.43 0.06	3.6 $\mu$ s 0.1	6 <sup>-</sup>	95	1969	IT=100
<sup>172</sup> Lu	-56736.1 2.3		6.70 d 0.03	4 <sup>-</sup> *	95	1951	$\beta^+$ =100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{172}\text{Lu}^m$	-56694.2	2.3	41.86 0.04	3.7 m 0.5			
$^{172}\text{Lu}^u$	-56670.3	2.3	65.79 0.04	332 ns 20			
$^{172}\text{Lu}^p$	-56626.7	2.3	109.41 0.10	440 $\mu\text{s}$ 12			
$^{172}\text{Lu}^q$	-56522.5	2.3	213.57 0.17	150 ns			
$^{172}\text{Hf}$	-56402	24		1.87 y 0.03			
$^{172}\text{Hf}^m$	-54396	24	2005.84 0.11	163 ns 3			
$^{172}\text{Ta}$	-51330	28		36.8 m 0.3			
$^{172}\text{W}$	-49097	28		6.6 m 0.9			
$^{172}\text{Re}$	-41570	40		55 s 5			
$^{172}\text{Re}^m$	-41460#	60#	110# 50#	15 s 3			
$^{172}\text{Os}$	-37244	13		19.2 s 0.9			
$^{172}\text{Ir}$	-27380	30		4.4 s 0.3			
$^{172}\text{Ir}^m$	-27240	30	139 10 AD	2.19 s 0.07			
$^{172}\text{Pt}$	-21107	10		97.6 ms 1.3			
$^{172}\text{Au}$	-9320	60		28 ms 4			
$^{172}\text{Au}^m$	-9160	240	160 250 AD*	11.0 ms 1.0			
$^{172}\text{Hg}$	-1060	150		231 $\mu\text{s}$ 9			
* $^{172}\text{Ho}$	T : other 17Wu04=27(11)						
* $^{172}\text{Re}$	J : direct $\beta^+$ feeding to 2+; conf=p9/2[514]n5/2[523] (N=97), K=2+						
* $^{172}\text{Re}^m$	J : direct $\beta^+$ feeding to 6+ and 8+; conf=p9/2[514]n5/2[523] (N=97), K=7+						
* $^{172}\text{Os}$	D : % $\alpha$ average 04GoZZ=1.4(0.3) 95Hi02=1.1(0.2)						
* $^{172}\text{Pt}$	D : % $\alpha$ average 10An02=97(3) 04GoZZ=94(6) 99Po09=94(12)						
* $^{172}\text{Au}$	T : symmetrized from 09Ha42=22(+6-4)						
* $^{172}\text{Au}$	J : favored $\alpha$ decay to $^{168}\text{Ir}$ [J=(2)-]						
* $^{172}\text{Au}^m$	T : average 09Ha42=9(+2-1) 09Ha42=8(+5-2) (independent measurements);						
* $^{172}\text{Au}^m$	T : others 96Pa01=6.3(1.5) 93Se09=4(1)						
* $^{172}\text{Au}^m$	J : favored $\alpha$ decay to $^{168}\text{Ir}^m$ [J=(9,10)+]						
$^{173}\text{Tb}$	-36510#	500#		400# ms >550ns			
$^{173}\text{Dy}$	-43740#	400#		1.43 s 0.20			
$^{173}\text{Ho}$	-49350#	300#		7.1 s 0.4			
$^{173}\text{Ho}^m$	-48950#	300#	405 1	3.7 $\mu\text{s}$ 1.2			
$^{173}\text{Er}$	-53650#	200#		1.434 m 0.017			
$^{173}\text{Tm}$	-56256	4		8.24 h 0.08			
$^{173}\text{Tm}^m$	-55938	4	317.73 0.20	10.7 $\mu\text{s}$ 1.7			
$^{173}\text{Tm}^n$	-54350	4	1905.7 0.4	250 ns 69			
$^{173}\text{Tm}^p$	-52208	4	4047.9 0.5	121 ns 28			
$^{173}\text{Yb}$	-57551.234	0.011		STABLE			
$^{173}\text{Yb}^m$	-57152.3	0.5	398.9 0.5	2.9 $\mu\text{s}$ 0.1			
$^{173}\text{Lu}$	-56881.0	1.6		1.37 y 0.01			
$^{173}\text{Lu}^m$	-56757.3	1.6	123.672 0.013	74.2 $\mu\text{s}$ 1.0			
$^{173}\text{Hf}$	-55412	28		23.6 h 0.1			
$^{173}\text{Hf}^m$	-55305	28	107.16 0.05	180 ns 8			
$^{173}\text{Hf}^n$	-55215	28	197.47 0.10	160 ns 40			
$^{173}\text{Ta}$	-52397	28		3.14 h 0.13			
$^{173}\text{Ta}^m$	-52224	28	173.10 0.21	205.2 ns 5.6			
$^{173}\text{Ta}^n$	-50680	28	1717.2 0.4	132 ns 3			
$^{173}\text{W}$	-48727	28		7.6 m 0.2			
$^{173}\text{Re}$	-43554	28		2.0 m 0.3			
$^{173}\text{Os}$	-37438	15		22.4 s 0.9			
$^{173}\text{Ir}$	-30268	11		9.0 s 0.8			
$^{173}\text{Ir}^m$	-30042	11	226 9 AD	2.20 s 0.05			
$^{173}\text{Pt}$	-21940	60		382 ms 2			
$^{173}\text{Au}$	-12832	23		25.5 ms 0.8			
$^{173}\text{Au}^m$	-12618	12	214 21 AD	12.2 ms 0.1			
$^{173}\text{Hg}$	-2660#	200#		800 $\mu\text{s}$ 80			
* $^{173}\text{Ho}$	T : average 20Li28=7.5(0.7) 17Wu04=6.9(0.5)						
* $^{173}\text{Tm}^m$	T : average 12Hu10=11.1(2.8) 72Pu02=10.4(2.1)						
* $^{173}\text{Ta}^m$	T : average 17Wo02=202(6) 91Ku12=225(15)						
* $^{173}\text{Ta}^n$	T : other 17Wo02=148(9)						
* $^{173}\text{Ir}$	J : $\alpha$ decay from $^{177}\text{Au}$ (J=1/2+)						
* $^{173}\text{Ir}$	D : % $\alpha$ from Ensdf2015<7						







**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	J <sup>π</sup>	Ens Reference	Year of discovery	Decay modes and intensities (%)
<sup>176</sup> Hf	-54576.4	1.5		STABLE	0 <sup>+</sup>	06	1934	IS=5.26 70
<sup>176</sup> Hf <sup>m</sup>	-53243.3	1.5	1333.07 0.07	9.6 μs 0.3	6 <sup>+</sup>	06	1964	IT=100
<sup>176</sup> Hf <sup>n</sup>	-53017.1	1.5	1559.31 0.09	9.9 μs 0.2	8 <sup>-</sup>	06	1967	IT=100
<sup>176</sup> Hf <sup>p</sup>	-51710.6	1.7	2865.8 0.7	401 μs 6	14 <sup>-</sup>	06	1975	IT=100
<sup>176</sup> Hf <sup>q</sup>	-49712.8	1.7	4863.6 0.9	43 μs 4	22 <sup>-</sup>	06 10Mu13 JT	1976	IT=100
<sup>176</sup> Ta	-51370	30		8.09 h 0.05	(1) <sup>-</sup>	06	1948	β <sup>+</sup> =100
<sup>176</sup> Ta <sup>m</sup>	-51270	30	103.0 1.0	1.08 ms 0.07	7 <sup>+</sup>	06 78Du06 ET	1971	IT=100
<sup>176</sup> Ta <sup>n</sup>	-49900	30	1474.0 1.4	3.8 μs 0.4	14 <sup>-</sup>	06	1978	IT=100
<sup>176</sup> Ta <sup>p</sup>	-48500	30	2874.0 1.4	970 μs 70	20 <sup>-</sup>	06	1994	IT=100
<sup>176</sup> W	-50642	28		2.5 h 0.1	0 <sup>+</sup>	06	1950	ε=100
<sup>176</sup> Re	-45063	28		5.3 m 0.3	(3 <sup>+</sup> )	06	1967	β <sup>+</sup> =100
<sup>176</sup> Os	-42131	11		3.6 m 0.5	0 <sup>+</sup>	06	1970	β <sup>+</sup> =100
<sup>176</sup> Ir	-33882	8		8.7 s 0.5	(3 <sup>+</sup> )	06 FGK208 J	1967	β <sup>+</sup> =96.9 6;α=3.1 6
<sup>176</sup> Ir <sup>m</sup>	-33830#	50#	50# 50#	10# s	(7 <sup>+</sup> )	06 FGK208 IJ		β <sup>+</sup> = ?;IT ?
<sup>176</sup> Pt	-28934	13		6.33 s 0.15	0 <sup>+</sup>	06	1966	β <sup>+</sup> ?;α=40 2
<sup>176</sup> Au	-18520	30		1.05 s 0.01	(3 <sup>-</sup> , 4 <sup>-</sup> )	06 14An10 JD	1975	α=75 8;β <sup>+</sup> ?
<sup>176</sup> Au <sup>m</sup>	-18380	30	139 13	1.36 s 0.02	(8 <sup>+</sup> , 9 <sup>+</sup> )	06 14An10 DJ	2002	α=?;β <sup>+</sup> ?
<sup>176</sup> Hg	-11785	11		20.3 ms 1.4	0 <sup>+</sup>	06	1983	α=90 9;β <sup>+</sup> ?
<sup>176</sup> Tl	580	80		6.2 ms 2.3	(3 <sup>-</sup> , 4 <sup>-</sup> )	09 04Ke06 TD	2004	p≈100;α ?;β <sup>+</sup> ?
* <sup>176</sup> Yb <sup>m</sup>	J : 12F105=8							**
* <sup>176</sup> Lu	T : average 37.20(0.23), evaluated by FGK using the world counting data,							**
* <sup>176</sup> Lu	T : 06Al03=37.13(0.26), using the world data on age comparison of							**
* <sup>176</sup> Lu	T : terrestrial minerals, 06Al03=35.40(0.80), using the world data on							**
* <sup>176</sup> Lu	T : age comparison of meteorites							**
* <sup>176</sup> Lu	D : %β <sup>+</sup> from 05Am04<0.9							**
* <sup>176</sup> Ta <sup>m</sup>	T : average 78Du06=1.05(0.10) 71Go21=1.1(0.1)							**
* <sup>176</sup> Ta <sup>n</sup>	E : 1371(1) keV above <sup>176</sup> Ta <sup>m</sup>							**
* <sup>176</sup> Ta <sup>p</sup>	E : 2771(1) keV above <sup>176</sup> Ta <sup>m</sup>							**
* <sup>176</sup> Ir	J : 205.2-keV gamma, most likely E2, from (1+) state populated by favored							**
* <sup>176</sup> Ir	J : α decay from <sup>180</sup> Au gs [J=(1+)]							**
* <sup>176</sup> Ir <sup>m</sup>	J : direct β <sup>+</sup> feeding to J=6 and 7 states in <sup>176</sup> Os implies existence							**
* <sup>176</sup> Ir <sup>m</sup>	J : of a higher-spin β <sup>+</sup> decaying isomer							**
* <sup>176</sup> Au <sup>m</sup>	T : from 04GoZZ; other 02Ro17=840(+170-140)							**
* <sup>176</sup> Hg	D : %α symmetrized from 99Po09=94(+6-12)%							**
* <sup>176</sup> Tl	T : symmetrized from 04Ke06=5.2(+3.0-1.4)							**
* <sup>176</sup> Tl	J : I <sub>p</sub> =0 to <sup>175</sup> Hg (J=7/2-) in 04Ke06							**
<sup>177</sup> Ho	-36280#	500#		1# s >550ns	7/2 <sup>-</sup> #	19 18Fu08 I	2018	β <sup>-</sup> =100;n ?
<sup>177</sup> Er	-42860#	500#		8# s >300ns	1/2 <sup>-</sup> #	19 12Ku26 I	2012	β <sup>-</sup> ?
<sup>177</sup> Tm	-47570#	200#		95 s 7	1/2 <sup>+</sup> #	19	1989	β <sup>-</sup> =100
<sup>177</sup> Tm <sup>m</sup>	-47470#	220#	100# 100#	77 s 11	7/2 <sup>-</sup> #	19	1989	β <sup>-</sup> =100
<sup>177</sup> Yb	-50986.40	0.22		1.911 h 0.003	9/2 <sup>+</sup> *	19	1945	β <sup>-</sup> =100
<sup>177</sup> Yb <sup>m</sup>	-50654.9	0.4	331.5 0.3	6.41 s 0.02	1/2 <sup>-</sup> *	19	1962	IT=100
<sup>177</sup> Lu	-52383.9	1.2		6.6443 d 0.0009	7/2 <sup>+</sup> *	19	1945	β <sup>-</sup> =100
<sup>177</sup> Lu <sup>m</sup>	-52233.5	1.2	150.3984 0.0010	130.1 ns 2.4	9/2 <sup>-</sup>	19	1949	IT=100
<sup>177</sup> Lu <sup>n</sup>	-51814.2	1.2	569.6721 0.0015	155 μs 7	1/2 <sup>+</sup>	19	1965	IT=100
<sup>177</sup> Lu <sup>p</sup>	-51413.7	1.2	970.1757 0.0024	160.4 d 0.3	23/2 <sup>-</sup> *	19	1962	β <sup>-</sup> =77.30 8;IT=22.70 8
<sup>177</sup> Lu <sup>q</sup>	-49612.2	1.3	2771.7 0.5	625 ns 62	33/2 <sup>+</sup>	19	2004	IT=100
<sup>177</sup> Lu <sup>r</sup>	-48853.5	1.3	3530.4 0.6	6 μs 2	39/2 <sup>-</sup>	19	2003	IT=100
<sup>177</sup> Hf	-52880.7	1.4		STABLE >1.3Ey	7/2 <sup>-</sup> *	19 20Da04 T	1934	IS=18.60 16
<sup>177</sup> Hf <sup>m</sup>	-51565.2	1.4	1315.4502 0.0008	1.09 s 0.05	23/2 <sup>+</sup>	19	1966	IT=100
<sup>177</sup> Hf <sup>n</sup>	-51538.3	1.7	1342.4 1.0	55.9 μs 1.2	(19/2 <sup>-</sup> )	19	1976	IT=100
<sup>177</sup> Hf <sup>p</sup>	-50140.7	1.4	2740.02 0.15	51.4 m 0.5	37/2 <sup>-</sup>	19	1971	IT=100
<sup>177</sup> Ta	-51715	3		56.36 h 0.13	7/2 <sup>+</sup> *	19	1948	β <sup>+</sup> =100
<sup>177</sup> Ta <sup>m</sup>	-51642	3	73.16 0.07	410 ns 7	9/2 <sup>-</sup>	19	1973	IT=100
<sup>177</sup> Ta <sup>n</sup>	-51529	3	186.16 0.06	3.62 μs 0.10	5/2 <sup>-</sup>	19	1971	IT=100
<sup>177</sup> Ta <sup>p</sup>	-50360	3	1354.8 0.3	5.30 μs 0.11	21/2 <sup>-</sup>	19	1971	IT=100
<sup>177</sup> Ta <sup>q</sup>	-47059	3	4656.3 0.8	133 μs 4	49/2 <sup>-</sup>	19	1994	IT=100
<sup>177</sup> W	-49702	28		132.4 m 2.0	1/2 <sup>-</sup>	19	1950	β <sup>+</sup> =100
<sup>177</sup> Re	-46269	28		14 m 1	5/2 <sup>-</sup>	19	1957	β <sup>+</sup> =100
<sup>177</sup> Re <sup>m</sup>	-46170#	60#	100# 50#	> 100 ns	9/2 <sup>-</sup>	19		IT=100
<sup>177</sup> Re <sup>n</sup>	-46184	28	84.70 0.10	50 μs 10	5/2 <sup>+</sup>	19	1972	IT=100



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
<sup>179</sup> Er	-36080#	500#	3# s >550ns	3/2 <sup>-</sup> #	18Fu08 I	2018	$\beta^- ?; \beta^- n ?$
<sup>179</sup> Tm	-41900#	400#	18# s >300ns	1/2 <sup>+</sup> #	13 12Ku26 I	2012	$\beta^- ?; \beta^- n ?$
<sup>179</sup> Yb	-46640#	200#	8.0 m 0.4	(1/2 <sup>-</sup> )	09	1982	$\beta^- =100$
<sup>179</sup> Lu	-49059	5	4.59 h 0.06	7/2 <sup>+</sup> *	09	1961	$\beta^- =100$
<sup>179</sup> Lu <sup>m</sup>	-48467	5	592.4 0.4	1/2 <sup>+</sup>	09	1982	IT=100
<sup>179</sup> Hf	-50463.0	1.4	STABLE	9/2 <sup>+</sup> *	09	1934	IS=13.62 11
<sup>179</sup> Hf <sup>m</sup>	-50088.0	1.4	18.67 s 0.04	1/2 <sup>-</sup>	09	1962	IT=100
<sup>179</sup> Hf <sup>m</sup>	-49356.6	1.4	1106.412 0.033	25/2 <sup>-</sup>	09 19Kr06 ET	1970	IT=100 *
<sup>179</sup> Hf <sup>p</sup>	-46687.8	2.5	3775.2 2.1	15 $\mu$ s 5	(43/2 <sup>+</sup> )	09	2000 IT=100
<sup>179</sup> Ta	-50357.5	1.5	1.82 y 0.03	7/2 <sup>+</sup>	09	1950	$\epsilon =100$
<sup>179</sup> Ta <sup>m</sup>	-50326.8	1.5	1.42 $\mu$ s 0.08	9/2 <sup>-</sup>	09	1964	IT=100
<sup>179</sup> Ta <sup>n</sup>	-49837.3	1.5	520.23 0.18	1/2 <sup>+</sup>	09 FGK128 J	1974	IT=100
<sup>179</sup> Ta <sup>p</sup>	-49104.9	1.5	1252.60 0.23	322 ns 16	21/2 <sup>-</sup>	09 97Ko13 J	1982 IT=100
<sup>179</sup> Ta <sup>q</sup>	-49040.3	1.6	1317.2 0.4	9.0 ms 0.2	25/2 <sup>+</sup>	09 97Ko13 J	1982 IT=100
<sup>179</sup> Ta <sup>r</sup>	-49029.5	1.6	1328.0 0.4	1.6 $\mu$ s 0.4	23/2 <sup>-</sup>	09 97Ko13 J	1982 IT=100
<sup>179</sup> Ta <sup>x</sup>	-47718.2	1.6	2639.3 0.5	54.1 ms 1.7	37/2 <sup>+</sup>	09 97Ko13 J	1982 IT=100
<sup>179</sup> W	-49295	15	37.05 m 0.16	7/2 <sup>-</sup>	09	1950	$\beta^+ =100$
<sup>179</sup> W <sup>m</sup>	-49073	15	221.91 0.03	6.40 m 0.07	1/2 <sup>-</sup>	09	1950 IT≈100; $\beta^+ =0.29$ 4
<sup>179</sup> W <sup>n</sup>	-47663	15	1631.90 0.08	390 ns 30	21/2 <sup>+</sup>	09 94Wa05 J	1978 IT=100
<sup>179</sup> W <sup>p</sup>	-45947	15	3348.41 0.14	750 ns 80	35/2 <sup>-</sup>	09 94Wa05 J	1978 IT=100
<sup>179</sup> Re	-46584	25	19.5 m 0.1	5/2 <sup>+</sup>	09	1960	$\beta^+ =100$
<sup>179</sup> Re <sup>m</sup>	-46519	25	65.35 0.09	95 $\mu$ s 25	(5/2 <sup>-</sup> )	09	1972 IT=100
<sup>179</sup> Re <sup>n</sup>	-44760#	60#	1822# 50#	408 ns 12	(23/2 <sup>+</sup> )	09	1972 IT=100 *
<sup>179</sup> Re <sup>p</sup>	-41176	25	5408.0 0.5	466 $\mu$ s 15	(47/2,49/2 <sup>+</sup> )	09 15Ko14 J	1989 IT=100
<sup>179</sup> Os	-43020	16	6.5 m 0.3	1/2 <sup>-</sup>	09	1968	$\beta^+ =100$
<sup>179</sup> Os <sup>m</sup>	-42875	16	145.41 0.12	500 ns	(7/2 <sup>-</sup> )	09	1983 IT=100
<sup>179</sup> Os <sup>n</sup>	-42777	16	243.0 0.8	783 ns 14	(9/2 <sup>+</sup> )	09	1983 IT=100
<sup>179</sup> Ir	-38082	10	79 s 1	(5/2 <sup>-</sup> )	09	1992	$\beta^+ =100$
<sup>179</sup> Pt	-32268	8	21.2 s 0.4	1/2 <sup>-</sup>	09	1966	$\beta^+ \approx 100; \alpha =0.24$ 3
<sup>179</sup> Au	-24989	12	7.1 s 0.3	1/2 <sup>+</sup> *	09 18Cu04 J	1968	$\beta^+ =78.0$ 9; $\alpha =22.0$ 9
<sup>179</sup> Au <sup>m</sup>	-24900	12	89.5 0.3	327 ns 5	(3/2 <sup>-</sup> )	11Ve01 ETD2011	IT=100 *
<sup>179</sup> Au <sup>p</sup>	-24856	19	133.5 15.0	(9/2 <sup>-</sup> )	09 11Ve01 EJD 1980		IT ? *
<sup>179</sup> Hg	-16933	28	1.05 s 0.03	7/2 <sup>-</sup> *	09 12Ve04 D	1970	$\alpha =75$ 4; $\beta^+ =25$ 4; $\beta^+ p \approx 0.15$ *
<sup>179</sup> Hg <sup>m</sup>	-16762	28	171.4 0.4	6.4 $\mu$ s 0.9	13/2 <sup>+</sup>	09 02Je09 J	2002 IT=100
<sup>179</sup> Tl	-8270	40	437 ms 9	1/2 <sup>+</sup> *	09	1983	$\alpha =60$ 2; $\beta^+ ?$ *
<sup>179</sup> Tl <sup>m</sup>	-7450#	40#	825# 10#	1.41 ms 0.02	(11/2 <sup>-</sup> )	09 18Ba46 TJ	1983 $\alpha \approx 100; IT ?; \beta^+ ?$ *
<sup>179</sup> Tl <sup>n</sup>	-7370	40	904.5 0.9	119 ns 14	(9/2 <sup>-</sup> )	09 18Ba46 TJD 2018	IT=100 *
<sup>179</sup> Pb	2050	80	2.7 ms 0.2	(9/2 <sup>-</sup> )	10 18Ba46 TJD 2010		$\alpha =100$ *
* <sup>179</sup> Hf <sup>m</sup>	T : average 19Kr06=24.91(0.27) 70KaZV=25.3(0.3) 73Ch18=24.8(0.3).						
* <sup>179</sup> Hf <sup>m</sup>	T : other 70Hu04=29(1)						
* <sup>179</sup> Re <sup>n</sup>	E : from 1772.20(0.22)+x keV; x=50#(50#) estimated by Nubase						
* <sup>179</sup> Au <sup>m</sup>	E : from 19Mo.B						
* <sup>179</sup> Au <sup>m</sup>	T : average 19Mo.B=304(9) 11Ve01=328(2); Birge ratio=2.6						
* <sup>179</sup> Au <sup>p</sup>	E : from 44(15) above 89.5 keV level						
* <sup>179</sup> Hg	J : 19Se04=7/2						
* <sup>179</sup> Tl	T : average 17Ba46=426(10) 11Ko.B=489(21); other 02Ro17=415(55)						
* <sup>179</sup> Tl	T : 13An10=265(10) 98To14=230(40) 83Sc24=160(+90-40)						
* <sup>179</sup> Tl	J : 17Ba04=1/2; favored $\alpha$ decay to <sup>175</sup> Au (J=1/2+)						
* <sup>179</sup> Tl	D : % $\alpha$ from 13An10						
* <sup>179</sup> Tl <sup>m</sup>	J : from favored $\alpha$ decay to <sup>175</sup> Au <sup>m</sup> [J=(11/2-)]						
* <sup>179</sup> Tl <sup>m</sup>	T : average 18Ba46=1.40(0.03) 11Ko.B=1.36(0.04) 10An01=1.46(0.04)						
* <sup>179</sup> Tl <sup>n</sup>	T : symmetrized from 18Ba46=114(+18-10)						
* <sup>179</sup> Pb	T : other 10An01=3.5(+1.4-0.8)						
<sup>180</sup> Er	-33180#	500#	2# s >550ns	0 <sup>+</sup>	18Fu08 I	2018	$\beta^- ?; \beta^- n ?$
<sup>180</sup> Tm	-38170#	400#	3# s >300ns		15	2012	$\beta^- ?; \beta^- n ?$
<sup>180</sup> Yb	-44720#	300#	2.4 m 0.5	0 <sup>+</sup>	15	1987	$\beta^- =100$
<sup>180</sup> Lu	-46680	70	5.7 m 0.1	5 <sup>+</sup>	15	1971	$\beta^- =100$
<sup>180</sup> Lu <sup>m</sup>	-46670	70	~ 1 s	3 <sup>-</sup>	15 95Me03 JT	1995	IT ?; $\beta^- ?$
<sup>180</sup> Lu <sup>n</sup>	-46060	70	624.0 0.5	> 1 ms	(9 <sup>-</sup> )	15 01Wh02 EJT 2001	IT=100



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>181</sup> Au	-27871	20		13.7 s 1.4	(5/2 <sup>-</sup> )	06	1968	$\beta^+$ =97.3 5; $\alpha$ =2.7 5	*
<sup>181</sup> Au <sup>p</sup>	-27660#	200#	210# 200#		(11/2 <sup>-</sup> )	06			
<sup>181</sup> Hg	-20661	15		3.6 s 0.1	1/2 <sup>-*</sup>	06	1969	$\beta^+$ =73 2; $\alpha$ =27 2; $\beta^+$ p=0.014 3; $\beta^+$ $\alpha$ =9e-6 3	*
<sup>181</sup> Hg <sup>m</sup>	-20450	50	210 50 IT	480 $\mu$ s 20	13/2 <sup>+</sup>	06 09An17 T	2009	IT=100	
<sup>181</sup> Tl	-12799	9		2.9 s 0.1	1/2 <sup>+</sup> *	09 18Cu04 TD	1996	$\beta^+$ ?; $\alpha$ =8.6 6	*
<sup>181</sup> Tl <sup>m</sup>	-11963	9	835.9 0.4	1.40 ms 0.03	(9/2 <sup>-</sup> )	09 09An14 JD	1984	IT=99.60 4; $\alpha$ =0.40 6; $\beta^+$ ?	*
<sup>181</sup> Pb	-3110	90		39.0 ms 0.8	(9/2 <sup>-</sup> )	06 09An20 TJ	1989	$\alpha$ $\approx$ 100; $\beta^+$ ?	*
<sup>181</sup> Pb <sup>m</sup>			non-exist RN		13/2 <sup>+</sup> #	96To01 I			
* <sup>181</sup> Ta <sup>p</sup>	E : 1403.2(0.6)+x keV with x<50 keV in 98Sa60								
* <sup>181</sup> Ta <sup>p</sup>	J : most likely E1 for the depopulating gamma to the 1403.2-keV (J=17/2-)								
* <sup>181</sup> Ta <sup>p</sup>	J : level, as proposed in 98Dr09, and hence J=19/2+; 98Sa60 suggests								
* <sup>181</sup> Ta <sup>p</sup>	J : J=15/2- for the 1403.2-keV level, but then J=19/2+ seems unlikely,								
* <sup>181</sup> Ta <sup>p</sup>	J : since the depopulating gamma would be M2 and much longer half-life								
* <sup>181</sup> Ta <sup>p</sup>	J : would be expected								
* <sup>181</sup> Ta <sup>q</sup>	T : average 98Wh02=25(2) 98Dr09=23(+6-2)								
* <sup>181</sup> W	T : average 14Un01=121.03(0.07) 73My02=120.95(0.02); other								
* <sup>181</sup> W	T : 72Em01=121.53(0.09), outlier								
* <sup>181</sup> Pt	J : 99Le52=1/2								
* <sup>181</sup> Au	J : favored $\alpha$ decay to <sup>177</sup> Ir (J=5/2-)								
* <sup>181</sup> Hg	J : 19Se04=1/2								
* <sup>181</sup> Hg	D : % $\beta^+$ p from I(p)/I( $\alpha$ )=5.0(0.8)e-4 in 71Ho07;								
* <sup>181</sup> Hg	D : % $\beta^+$ $\alpha$ from I( $\alpha$ )/I( $\beta^+$ )=1.2e-7 in 75Ho02								
* <sup>181</sup> Tl	J : 17Ba04=1/2								
* <sup>181</sup> Tl	T : others 98To14=3.2(0.3) 92Bo.D=3.4(0.6)								
* <sup>181</sup> Pb	T : average 09An20=36(2) 05Ca.A=39.6(0.9)								
<sup>182</sup> Tm	-31490#	500#						$\beta^-$ ?; $\beta^-$ n ?	
<sup>182</sup> Yb	-38900#	400#		30# s >300ms	0 <sup>+</sup>	15 12Ku26 I	2012	$\beta^-$ ?	
<sup>182</sup> Lu	-41770#	200#		2.0 m 0.2	1 <sup>-</sup> #	15	1982	$\beta^-$ =100	
<sup>182</sup> Hf	-46050	6		8.90 My0.09	0 <sup>+</sup>	15	1961	$\beta^-$ =100	
<sup>182</sup> Hf <sup>m</sup>	-44877	6	1172.87 0.18	61.5 m 1.5	8 <sup>-</sup>	15 15Ko14 J	1971	$\beta^-$ =54 2;IT=46 2	*
<sup>182</sup> Hf <sup>m</sup>	-43479	6	2571.3 1.2	40 $\mu$ s 10	(13 <sup>+</sup> )	15	1999	IT=100	
<sup>182</sup> Ta	-46430.7	1.6		114.74 d 0.12	3 <sup>-</sup>	15	1938	$\beta^-$ =100	
<sup>182</sup> Ta <sup>m</sup>	-46414.4	1.6	16.273 0.004	283 ms 3	5 <sup>+</sup>	15	1968	IT=100	
<sup>182</sup> Ta <sup>n</sup>	-45911.1	1.6	519.577 0.016	15.84 m 0.10	10 <sup>-</sup>	15	1947	IT=100	
<sup>182</sup> W	-48246.1	0.7		STABLE >7.7Zy	0 <sup>+</sup>	15 04Co26 T	1930	IS=26.50 16; $\alpha$ ?	*
<sup>182</sup> W <sup>m</sup>	-46015.5	0.7	2230.65 0.14	1.3 $\mu$ s 0.1	10 <sup>+</sup>	15 15Ko14 J	1969	IT=100	
<sup>182</sup> Re	-45450	100		64.2 h 0.5	7 <sup>+</sup> *	15	1950	$\beta^+$ =100	
<sup>182</sup> Re <sup>m</sup>	-45386	20	60 100 BD*	14.14 h 0.45	2 <sup>+</sup> *	15	1950	$\beta^+$ =100	
<sup>182</sup> Re <sup>n</sup>	-45150	140	296 100	585 ns 30	(2) <sup>-</sup>	15	1969	IT=100	*
<sup>182</sup> Re <sup>p</sup>	-44930	140	521 100	780 ns 90	(4) <sup>-</sup>	15	1984	IT=100	*
<sup>182</sup> Os	-44609	22		21.84 h 0.20	0 <sup>+</sup>	15	1950	$\epsilon$ =100	
<sup>182</sup> Os <sup>m</sup>	-42778	22	1831.4 0.3	780 $\mu$ s 70	8 <sup>-</sup>	15 15Ko14 J	1966	IT=100	
<sup>182</sup> Os <sup>n</sup>	-37560	22	7049.5 0.4	150 ns 10	25 <sup>+</sup>	15 15Ko14 J	1988	IT=100	
<sup>182</sup> Ir	-39052	21		15.0 m 1.0	3 <sup>+</sup>	15	1961	$\beta^+$ =100	
<sup>182</sup> Ir <sup>m</sup>	-38981	21	71.02 0.17	170 ns 40	(5) <sup>+</sup>	15	1990	IT=100	
<sup>182</sup> Ir <sup>n</sup>	-38876	21	176.4 0.3	130 ns 50	(6) <sup>-</sup>	15	1990	IT=100	
<sup>182</sup> Pt	-36168	13		2.67 m 0.12	0 <sup>+</sup>	15	1963	$\beta^+$ =0.962 2; $\alpha$ =0.038 2	
<sup>182</sup> Au	-28304	19		15.5 s 0.4	(2 <sup>+</sup> ) <sup>*</sup>	15 20Ha24 J	1970	$\beta^+$ $\approx$ 100; $\alpha$ =0.13 5	
<sup>182</sup> Au <sup>m</sup>	-28180	30	120 40	10# s	5 <sup>-</sup> #			$\beta^+$ =?; IT ?	*
<sup>182</sup> Hg	-23577	10		10.83 s 0.06	0 <sup>+</sup>	15 71Ho07 D	1968	$\beta^+$ =86.2 9; $\alpha$ =13.8 9; $\beta^+$ p<1e-5	*
<sup>182</sup> Tl	-13327	12	*&	1.9 s 0.1	(4) <sup>-</sup> *	10 16Va01 TD	1991	$\beta^+$ $\approx$ 100; $\alpha$ >0.49; $\beta^+$ SF<3.4e-6	*
<sup>182</sup> Tl <sup>m</sup>	-13280#	50#	50# 50# *&	3.1 s 1.0	(7 <sup>+</sup> ) <sup>*</sup>	91Bo02 TD	1991	$\beta^+$ $\approx$ 100; $\alpha$ =2.5 14;IT ?	*
<sup>182</sup> Tl <sup>p</sup>	-12830#	100#	500# 100#		(10) <sup>-</sup>				
<sup>182</sup> Pb	-6825	12		55 ms 5	0 <sup>+</sup>	15	1986	$\alpha$ $\approx$ 100; $\beta^+$ ?	
* <sup>182</sup> Hf <sup>m</sup>	J : E1 to 8+								
* <sup>182</sup> W	T : 04Co26>7.7Zy; others 03Da05>170Ey 03Ce01>25Ey 97Ge15>8.3Ey								
* <sup>182</sup> Re <sup>n</sup>	E : 235.732(0.022) keV above <sup>182</sup> Re <sup>m</sup>								
* <sup>182</sup> Re <sup>p</sup>	E : 461.3(0.1) keV above <sup>182</sup> Re <sup>m</sup>								

Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
* $^{182}\text{Au}^m$	I: direct $\beta^+$ feeding to 4+ and 5+ levels in $^{182}\text{Pt}$ (99Da18)								
* $^{182}\text{Au}^m$	I: indicates the existence of an isomer								
* $^{182}\text{Hg}$	D: % $\alpha$ average 97Ba21=13.3(0.5) 80Sc09=15.2(0.8); $\beta^+$ p 71Ho07<1e-5								
* $^{182}\text{Tl}$	J: 17Ba04=(4)								
* $^{182}\text{Tl}^m$	J: 17Ba04=(7)								
* $^{182}\text{Tl}^m$	D: % $\alpha$ from 91Bo01<5%								
$^{183}\text{Yb}$	-35000#	400#			30# s >300ns	3/2 <sup>-</sup> #	16 12Ku26 I	2012	$\beta^-$ ?
$^{183}\text{Lu}$	-39720	80			58 s 4	7/2 <sup>+</sup> #	16	1983	$\beta^-$ =100
$^{183}\text{Hf}$	-43280	30			1.018 h 0.002	(3/2 <sup>-</sup> )	16	1956	$\beta^-$ =100
$^{183}\text{Hf}^m$	-41820	70	1464	64	40 s 30	27/2 <sup>-</sup> #	16 10Re07 ETJ	2010	IT $\approx$ 100; $\beta^-$ ?
$^{183}\text{Ta}$	-45293.5	1.6			5.1 d 0.1	7/2 <sup>+</sup> *	16	1950	$\beta^-$ =100
$^{183}\text{Ta}^m$	-45220.3	1.6	73.164	0.014	106 ns 10	9/2 <sup>-</sup>	16	1967	IT=100
$^{183}\text{Ta}^n$	-43959	14	1335	14	900 ns 300	(19/2 <sup>+</sup> )	16 09Sh17 ETJ	2009	IT=100
$^{183}\text{W}$	-46365.7	0.7			STABLE >670Ey	1/2 <sup>-</sup> *	16	1930	IS=14.31 4; $\alpha$ ?
$^{183}\text{W}^m$	-46056.2	0.7	309.492	0.004	5.30 s 0.08	11/2 <sup>+</sup>	16	1961	IT=100
$^{183}\text{Re}$	-45810	8			70.0 d 1.4	5/2 <sup>+</sup> *	16	1950	$\epsilon$ =100
$^{183}\text{Re}^m$	-43903	8	1907.21	0.15	1.04 ms 0.04	25/2 <sup>+</sup>	16 15Ko14 J	1966	IT=100
$^{183}\text{Os}$	-43660	50			13.0 h 0.5	9/2 <sup>+</sup> *	16	1950	$\beta^+$ =100
$^{183}\text{Os}^m$	-43490	50	170.73	0.07	9.9 h 0.3	1/2 <sup>-</sup> *	16	1958	$\beta^+$ =85 2;IT=15 2
$^{183}\text{Ir}$	-40202	25			58 m 5	5/2 <sup>-</sup>	16 61Di04 T	1961	$\beta^+$ $\approx$ 100; $\alpha$ ?
$^{183}\text{Pt}$	-35773	14			6.5 m 1.0	1/2 <sup>-</sup> *	16	1963	$\beta^+$ $\approx$ 100; $\alpha$ =0.0096 5
$^{183}\text{Pt}^m$	-35738	14	34.74	0.07	43 s 5	7/2 <sup>-</sup> *	16	1979	$\beta^+$ =96.9 8;IT=3.1 8; $\alpha$ ?
$^{183}\text{Pt}^n$	-35577	14	195.90	0.10	> 150 ns	9/2 <sup>+</sup>	16	1990	IT=100
$^{183}\text{Au}$	-30191	9			42.8 s 1.0	5/2 <sup>-</sup> *	16	1968	$\beta^+$ =99.45 25; $\alpha$ =0.55 25
$^{183}\text{Au}^m$	-30118	9	73.10	0.01	> 1 $\mu$ s	(1/2 <sup>+</sup> )	16 17Ve04 E	1984	IT=100
$^{183}\text{Au}^p$	-29960	9	230.6	0.6	< 1 $\mu$ s	(11/2 <sup>-</sup> )	16	1984	IT=100
$^{183}\text{Hg}$	-23805	7			9.4 s 0.7	1/2 <sup>-</sup> *	16	1969	$\beta^+$ =88.3 20; $\alpha$ =11.7 20; $\beta^+$ p=2.6e-4 6
$^{183}\text{Hg}^m$	-23601	13	204	14	> 5# $\mu$ s	13/2 <sup>+</sup> #	81Mi12 I	1981	IT ?; $\beta^+$ ?
$^{183}\text{Tl}$	-16587	9			6.9 s 0.7	1/2 <sup>+</sup> *	16	1980	$\beta^+$ =?; $\alpha$ ?
$^{183}\text{Tl}^m$	-15959	9	628.7	0.5	53.3 ms 0.3	(9/2 <sup>-</sup> )	16 11Ve.A E	1980	IT ?; $\alpha$ =1.5 3; $\beta^+$ ?
$^{183}\text{Tl}^n$	-15612	9	975.3	0.6	1.48 $\mu$ s 0.10	(13/2 <sup>+</sup> )	16	2001	IT=100
$^{183}\text{Pb}$	-7580	29			535 ms 30	3/2 <sup>-</sup> *	16	1980	$\alpha$ $\approx$ 100; $\beta^+$ ?
$^{183}\text{Pb}^m$	-7486	28	94	8	415 ms 20	13/2 <sup>+</sup> *	16	1980	$\alpha$ $\approx$ 100; $\beta^+$ ?;IT ?
* $^{183}\text{Hf}^m$	T: symmetrized from 10Re07=10(+48-5), value for q=71+ (H+ like ion); the								
* $^{183}\text{Hf}^m$	T: actual half-life could be shorter								
* $^{183}\text{Ta}^n$	E: from 1310.16 + x keV with x<50 keV in 09Sh17								
* $^{183}\text{Os}^m$	J: 75Ru06,78Ru04=1/2								
* $^{183}\text{Ir}$	T: average 61Di04=55(7) 61La05=60(6)								
* $^{183}\text{Pt}$	J: 92Hi07,99Le52=1/2								
* $^{183}\text{Pt}^m$	J: 99Le52=7/2								
* $^{183}\text{Au}$	J: 89Wa11,94Pa37=5/2								
* $^{183}\text{Hg}$	J: 72Bo09,76Bo09,19Se04=1/2								
* $^{183}\text{Hg}$	D: % $\beta^+$ p from 71Ho07=l(p)/l(a)=2.2(0.3)e-5								
* $^{183}\text{Hg}^m$	I: lack of 6073 $\alpha$ -gamma coincidences in $^{187}\text{Pb}^m$ decay								
* $^{183}\text{Tl}$	J: 17Ba04j,13Ba41=1/2+								
* $^{183}\text{Tl}^m$	E: uncertainty estimated by Nubase								
* $^{183}\text{Pb}$	J: 09Se13=3/2								
* $^{183}\text{Pb}^m$	J: 09Se13=13/2								
$^{184}\text{Yb}$	-32600#	500#			7# s >300ns	0 <sup>+</sup>	13 12Ku26 I	2012	$\beta^-$ ?
$^{184}\text{Lu}$	-36300#	200#			20 s 3	(3 <sup>+</sup> )	10 95Kr04 TJ	1989	$\beta^-$ =100
$^{184}\text{Hf}$	-41500	40			4.12 h 0.05	0 <sup>+</sup>	10	1973	$\beta^-$ =100
$^{184}\text{Hf}^m$	-40230	40	1272.2	0.4	48 s 10	8 <sup>-</sup>	10 12Re.A D	1995	IT $\approx$ 100; $\beta^-$ ?
$^{184}\text{Hf}^n$	-39020	40	2477	10	16 m 7	15 <sup>+</sup> #	10 10Re07 ET	2010	$\beta^-$ ?; IT ?
$^{184}\text{Ta}$	-42839	26			8.7 h 0.1	(5 <sup>-</sup> )	10	1955	$\beta^-$ =100
$^{184}\text{W}$	-45705.5	0.7			STABLE >8.9Zy	0 <sup>+</sup>	10 04Co26 T	1930	IS=30.64 2; $\alpha$ ?
$^{184}\text{W}^m$	-44420.5	0.7	1284.997	0.008	8.33 $\mu$ s 0.18	5 <sup>-</sup>	10	1969	IT=100
$^{184}\text{W}^n$	-41577.8	0.9	4127.7	0.5	188 ns 38	(14 <sup>+</sup> )	10 15Ko14 JE	2004	IT=100
$^{184}\text{Re}$	-44220	4			35.4 d 0.7	3 <sup>-</sup>	10	1940	$\beta^+$ =100





**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	J <sup>π</sup>	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>185</sup> Ir <sup>m</sup>	E : 2157.3(0.5) + x keV; x<80 keV						**
* <sup>185</sup> Pt	J : 92Hi07,99Le52=9/2						**
* <sup>185</sup> Pt	D : %α from E(a)=4444(10) keV in 91Bi04						**
* <sup>185</sup> Pt <sup>m</sup>	J : 92Hi07,99Le52=1/2						**
* <sup>185</sup> Hg	J : 19Se04=1/2						**
* <sup>185</sup> Hg <sup>m</sup>	J : 19Se04=13/2						**
* <sup>185</sup> Pb	J : 09Se13=3/2						**
* <sup>185</sup> Pb <sup>m</sup>	T : average 02An15=4.3(0.2) 80Sc09=3.73(0.24) (excluding the 6.1 s activity)						**
* <sup>185</sup> Pb <sup>m</sup>	J : 09Se13=13/2						**
* <sup>185</sup> Bi	T : estimated from J=9/2- isomers in odd Bi and Tl isotopes						**
* <sup>185</sup> Bi <sup>n</sup>	E : 100 keV above <sup>185</sup> Bi <sup>m</sup>						**
<sup>186</sup> Lu	-30320#	400#	6# s >300ns		13 12Ku26 I	2012	β <sup>-</sup> ?;β <sup>-</sup> n ?
<sup>186</sup> Hf	-36420	50	2.6 m 1.2	0 <sup>+</sup>	03	1998	β <sup>-</sup> =100
<sup>186</sup> Hf <sup>m</sup>	-33450	70	2968 43	> 20 s	17 <sup>+</sup> #	10Re07 ET	2010 β <sup>-</sup> ?; IT ?
<sup>186</sup> Ta	-38610	60	10.5 m 0.3	3#	03	1955	β <sup>-</sup> =100
<sup>186</sup> Ta <sup>m</sup>	-38270	60	1.54 m 0.05	9 <sup>+</sup> #	04Xu08 T	2010	β <sup>-</sup> ?; IT ?
<sup>186</sup> W	-42508.6	1.2	STABLE	>4.1EY	0 <sup>+</sup>	03 03Da09 T	1930 IS=28.43 19;2β <sup>-</sup> ?;α ?
<sup>186</sup> W <sup>m</sup>	-40991.4	1.3	1517.2 0.6	18 μs 1	7 <sup>-</sup>	03 12La.A J	1998 IT=100
<sup>186</sup> W <sup>n</sup>	-38965.8	2.4	3542.8 2.1	2.0 s 0.2	16 <sup>+</sup>	03 12La.A TJ	1998 IT=100
<sup>186</sup> Re	-41927.3	0.8	3.7185 d 0.0005	1 <sup>-</sup> *	03 FGK204 T	1939	β <sup>-</sup> =92.53 10;ε=7.47 10
<sup>186</sup> Re <sup>m</sup>	-41779.1	0.9	148.2 0.5	~ 200 ky	(8 <sup>+</sup> )	03 15Ma60 E	1972 IT≈100;β <sup>-</sup> ?
<sup>186</sup> Os	-43000.0	0.8	2.0 Py 1.1	0 <sup>+</sup>	03	1931	IS=1.59 64;α=100
<sup>186</sup> Ir	-39172	17	16.64 h 0.03	5 <sup>+</sup> *	03	1958	β <sup>+</sup> =100
<sup>186</sup> Ir <sup>m</sup>	-39171	17	0.8 0.4	1.92 h 0.05	2 <sup>-</sup>	03 91Be25 ET	1962 β <sup>+</sup> ≈75;IT≈25
<sup>186</sup> Pt	-37864	22	10.8 h 0.05	0 <sup>+</sup>	03	1961	β <sup>+</sup> =100;α≈1.4e-4
<sup>186</sup> Au	-31715	21	2.07 m 0.5	3 <sup>-</sup> *	03	1960	β <sup>+</sup> =100;α≈0.0008 2
<sup>186</sup> Au <sup>m</sup>	-31670#	40#	50# 30#	> 1 μs	6 <sup>-</sup> #		IT ?;β <sup>+</sup> ?
<sup>186</sup> Au <sup>n</sup>	-31487	21	227.77 0.07	110 ns 10	2 <sup>+</sup>	03	1983 IT=100
<sup>186</sup> Hg	-28539	12	1.38 m 0.06	0 <sup>+</sup>	03	1960	β <sup>+</sup> ≈100;α=0.016 5
<sup>186</sup> Hg <sup>m</sup>	-26322	12	2217.3 0.4	82 μs 5	(8 <sup>-</sup> )	03	1984 IT=100
<sup>186</sup> Tl	-19883	21	3.5 s 0.5	(2 <sup>-</sup> )	03 20St11 TDJ	1975	β <sup>+</sup> ?;α=?
<sup>186</sup> Tl <sup>m</sup>	-19860	30	20 40	27.5 s 1.0	7 <sup>+</sup> *	03 13Ba41 J	1975 β <sup>+</sup> ≈100;α≈0.006
<sup>186</sup> Tl <sup>n</sup>	-19490	30	390 40 MD	3.40 s 0.09	10 <sup>-</sup> *	03 20St11 TD	1977 IT<94.1 3;β <sup>+</sup> >5.9 3
<sup>186</sup> Pb	-14681	11	4.82 s 0.03	0 <sup>+</sup>	03	1972	β <sup>+</sup> ?;α=40 8
<sup>186</sup> Bi	-3145	17	14.8 ms 0.7	(3 <sup>+</sup> )	03 13La02 D	1997	α≈100;β <sup>+</sup> =?;β <sup>+</sup> SF≈0.011
<sup>186</sup> Bi <sup>m</sup>	-2980#	100#	170# 100#	9.8 ms 0.4	(10 <sup>-</sup> )	03 13La02 D	1984 α≈100;β <sup>+</sup> =?;β <sup>+</sup> SF≈0.011
<sup>186</sup> Po	4102	18	34 μs 12	0 <sup>+</sup>	13 13An13 T	2005	α≈100;p ?
* <sup>186</sup> Hf <sup>m</sup>	T : for q=72+ (bare ion) in 10Re07; the actual half-life could be shorter						**
* <sup>186</sup> Ta	J : direct β <sup>-</sup> feeding to 3- in <sup>186</sup> Hf; conf p7/2[404]n1/2[510],K=3- or						**
* <sup>186</sup> Ta	J : p9/2[514]n3/2[512],K=3+						**
* <sup>186</sup> Ta <sup>m</sup>	T : other 12Re19=3.0(+1.5-0.8) q=72+(H+ like ion); supersedes						**
* <sup>186</sup> Ta <sup>m</sup>	T : 10Re07=3.4(+2.4-1.4)						**
* <sup>186</sup> Ta <sup>m</sup>	E : from 10Re07						**
* <sup>186</sup> W	T : the limit given is for 2β <sup>-</sup> decay; α decay 04Co26>8.2 Zy						**
* <sup>186</sup> W	T : 03Da05>170 Ey 03Ce01>27 Ey 97Ge15>6.5 Ey						**
* <sup>186</sup> Ir <sup>m</sup>	T : average 91Be25=1.90(0.05) 70Fi.A=2.0(0.1)						**
* <sup>186</sup> Au <sup>m</sup>	I : floated strongly-coupled band in 92Ja01; conf p3/2[532]n9/2[624],K=6-,						**
* <sup>186</sup> Au <sup>m</sup>	I : same as the ground state where K=3-						**
* <sup>186</sup> Tl	T : symmetrized from 20St11=3.4(+0.5-0.4)						**
* <sup>186</sup> Tl <sup>n</sup>	E : 374.2(0.1) keV above <sup>186</sup> Tl <sup>m</sup>						**
* <sup>186</sup> Tl <sup>n</sup>	J : 13Ba41=10						**
* <sup>186</sup> Bi	T : average 03An27=14.8(0.8) 97Ba21=15.0(1.7)						**
* <sup>186</sup> Bi	D : %β <sup>+</sup> SF 13La02=0.022 13 for both isomers						**
* <sup>186</sup> Bi <sup>m</sup>	T : from 03An27						**
* <sup>186</sup> Bi <sup>m</sup>	D : %β <sup>+</sup> SF 13La02=0.022 13 for both isomers						**
* <sup>186</sup> Po	T : symmetrized from 13An13=28(+16-6)						**
<sup>187</sup> Lu	-27770#	400#	7# s >300ns	7/2 <sup>+</sup> #	13 12Ku26 I	2012	β <sup>-</sup> ?
<sup>187</sup> Hf	-33000#	200#	14# s >300ns	9/2 <sup>-</sup> #	09 99Be63 I	1999	β <sup>-</sup> ?
<sup>187</sup> Hf <sup>m</sup>	-32500#	360#	500# 300#	270 ns 80	3/2 <sup>-</sup> #	09A130 TD	2009 IT=100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)				
$^{187}\text{Ta}$	-36900	60		2.3 m	6	(7/2 <sup>+</sup> )	09 10Re07 T	1999	$\beta^-$ =100	*		
$^{187}\text{Ta}^m$	-35110	60	1778	1	7.3 s	0.9	(25/2 <sup>-</sup> )	20Wa29 ETJ	2010	IT≈100; $\beta^-$ ?	*	
$^{187}\text{Ta}^n$	-33970	60	2935	14	> 5 m		41/2 <sup>+</sup> #	10Re07 ET	2010	$\beta^-$ ?; IT ?	*	
$^{187}\text{W}$	-39904.0	1.2			23.809 h	0.025	3/2 <sup>-*</sup>	09 19Kr02 T	1940	$\beta^-$ =100	*	
$^{187}\text{W}^m$	-39493.9	1.2	410.06	0.04	1.38 μs	0.07	11/2 <sup>+</sup>	09 08Bo26 J	2008	IT=100	*	
$^{187}\text{Re}$	-41216.5	0.7			41.6 Gy	0.02	5/2 <sup>+</sup> *	09 01Be81 T	1931	IS=62.60 5; $\beta^-$ =100; $\alpha$ =0	*	
$^{187}\text{Re}^m$	-41010.3	0.7	206.2473	0.0010	555.3 ns	1.7	9/2 <sup>-</sup>	09	1949	IT=100	*	
$^{187}\text{Re}^n$	-39534.5	0.9	1682.0	0.6	354 ns	62	21/2 <sup>+</sup>	09 16Re02 ETJ	2003	IT=100	*	
$^{187}\text{Os}$	-41219.0	0.7			STABLE	>3.2Py	1/2 <sup>-*</sup>	09	1931	IS=1.96 17; $\alpha$ ?	*	
$^{187}\text{Os}^m$	-41118.6	0.7	100.45	0.04	112 ns	6	7/2 <sup>-</sup>	09	1964	IT=100	*	
$^{187}\text{Os}^n$	-40961.9	0.7	257.10	0.07	231 μs	2	11/2 <sup>+</sup>	09	1964	IT=100	*	
$^{187}\text{Ir}$	-39549	28			10.5 h	0.3	3/2 <sup>+</sup> *	09	1958	$\beta^+$ =100	*	
$^{187}\text{Ir}^m$	-39363	28	186.16	0.04	30.3 ms	0.6	9/2 <sup>-</sup>	09	1963	IT=100	*	
$^{187}\text{Ir}^n$	-39115	28	433.75	0.06	152 ns	12	11/2 <sup>-</sup>	09	1969	IT=100	*	
$^{187}\text{Ir}^p$	-37061	28	2487.7	0.4	1.8 μs	0.5	29/2 <sup>-</sup>	10Mo09 ETJ	2010	IT=100	*	
$^{187}\text{Pt}$	-36685	24			2.35 h	0.03	3/2 <sup>-*</sup>	09	1961	$\beta^+$ =100	*	
$^{187}\text{Pt}^m$	-36511	24	174.38	0.22	311 μs	15	11/2 <sup>+</sup>	09	1976	IT=100	*	
$^{187}\text{Au}$	-33029	22			8.3 m	0.2	1/2 <sup>+</sup> *	09	1955	$\beta^+$ ≈100; $\alpha$ ?	*	
$^{187}\text{Au}^m$	-32909	22	120.33	0.14	2.3 s	0.1	9/2 <sup>-*</sup>	09 20Ba29 J	1983	IT=100	*	
$^{187}\text{Hg}$	-28119	13			1.9 m	0.3	3/2 <sup>-*</sup>	09 70Ha18 TD	1960	$\beta^+$ =100; $\alpha$ ?	*	
$^{187}\text{Hg}^m$	-28060	15	58	14	MD	2.4 m	13/2 <sup>+</sup> *	09 70Ha18 D	1970	$\beta^+$ =100; $\alpha$ ?	*	
$^{187}\text{Tl}$	-22445	8			~ 51 s		1/2 <sup>+</sup>	09	1976	$\beta^+$ ≈100; $\alpha$ ?	*	
$^{187}\text{Tl}^m$	-22111	8	334	3	AD	15.60 s	9/2 <sup>-*</sup>	09 13Ba41 J	1976	IT=?; $\beta^+$ =?; $\alpha$ =0.15 5	*	
$^{187}\text{Tl}^n$	-20570#	50#	1875#	50#	1.11 μs	0.7		09 00By02 T	2000	IT=100	*	
$^{187}\text{Tl}^p$	-19863	8	2582.5	0.3	693 ns	38	29/2 <sup>+</sup> #	09	2000	IT=100	*	
$^{187}\text{Pb}$	-14987	5			*	15.2 s	3/2 <sup>-*</sup>	09	1972	$\beta^+$ =90.5 20; $\alpha$ =9.5 20	*	
$^{187}\text{Pb}^m$	-14968	11	19	10	MD*	18.3 s	13/2 <sup>+</sup> *	09	1972	$\beta^+$ =88 2; $\alpha$ =12 2	*	
$^{187}\text{Bi}$	-6383	10			37 ms	2	(9/2 <sup>-</sup> )	09	1999	$\alpha$ =100	*	
$^{187}\text{Bi}^m$	-6275	12	108	8	AD	370 μs	20	1/2 <sup>+</sup>	09	1984	$\alpha$ =100	*
$^{187}\text{Bi}^n$	-6131	10	252	3		7 μs	5	(13/2 <sup>+</sup> )	09 02Hu14 ETJ	2002	IT=100	*
$^{187}\text{Po}$	2820	30			*	1.40 ms	0.25	1/2 <sup>-</sup> , 5/2 <sup>-</sup>	09	2005	$\alpha$ ≈100; $\beta^+$ ?	*
$^{187}\text{Po}^m$	2830	40	4	27	AD*	0.5 ms		13/2 <sup>+</sup> #	06An11 ETD	2006	$\alpha$ ≈100; $\beta^+$ ?	*
* $^{187}\text{Ta}$	J : from 20Wa29									**		
* $^{187}\text{Ta}^m$	TE : other 10Re07=22(9) s for q=73+ (bare ion); E=1789(13) keV in 10Re07									**		
* $^{187}\text{Ta}^n$	T : from 10Re07 for q=73+ (bare ion)									**		
* $^{187}\text{W}$	T : average 19Kr0=23.80(0.03) 64An02=23.72(0.06) 57Wr37=24.04(0.09)									**		
* $^{187}\text{W}^m$	T : 53Ei02=23.85(0.08)									**		
* $^{187}\text{Re}$	J : E1 to 9/2 <sup>-</sup> ; l(d,p)=(6)									**		
* $^{187}\text{Re}^m$	T : recommended in 01Be81, based on 96Sm.A data, in agreement									**		
* $^{187}\text{Re}^n$	T : with 01Ga01=41.2(1.1) (direct measurement); other: 96Bo37=32.9(2.0) y									**		
* $^{187}\text{Re}^p$	T : for q=75+ (bare ion)									**		
* $^{187}\text{Os}$	T : from 20Be23 for T1/2( $\alpha$ , 1/2 <sup>-</sup> →3/2 <sup>-</sup> )									**		
* $^{187}\text{Pt}$	J : 92Hi07=3/2									**		
* $^{187}\text{Pt}^m$	J : M2 to 7/2-									**		
* $^{187}\text{Hg}$	T : from 70Ha18; other 98Ru04=2.4 m, but no uncertainty given									**		
* $^{187}\text{Hg}^m$	T : from 70Ha18; other 98Ru04=2.2 m, but no uncertainty given									**		
* $^{187}\text{Tl}^n$	E : 1433.23(0.19)+191+201+x keV; x=50#(50#) keV estimated by Nubase									**		
* $^{187}\text{Pb}$	J : 09Se13=3/2									**		
* $^{187}\text{Pb}^m$	J : 09Se13=13/2									**		
* $^{187}\text{Bi}$	J : favored $\alpha$ decay to $^{183}\text{Tl}^m$ [J=(9/2-)]									**		
* $^{187}\text{Bi}^m$	J : favored $\alpha$ decay to $^{183}\text{Tl}$ (J=1/2+)									**		
* $^{187}\text{Bi}^n$	T : symmetrized from 02Hu14=3.2(+7.6-2.0)									**		
* $^{187}\text{Bi}^p$	E : 02Hu14=252 keV gamma at the focal plane of RITU separator;									**		
* $^{187}\text{Bi}^m$	E : uncertainty estimated by Nubase									**		
* $^{187}\text{Po}$	J : favored $\alpha$ decay to J=1/2-, 5/2- level in $^{183}\text{Pb}$									**		
$^{188}\text{Lu}$	-23820#	400#			1#	s >300ns		18 12Ku26 I	2012	$\beta^-$ ?; $\beta^-$ n ?	*	
$^{188}\text{Hf}$	-30830#	300#			7#	s >300ns	0 <sup>+</sup>	18 99Be63 I	1999	$\beta^-$ ?	*	
$^{188}\text{Ta}$	-33910#	200#			19.6 s	2.0	(1 <sup>-</sup> )	18	1999	$\beta^-$ =100	*	
$^{188}\text{Ta}^m$	-33810#	200#	99	33	19.6 s	2.0	(7 <sup>-</sup> )	18	2005	IT ?; $\beta^-$ ?	*	
$^{188}\text{Ta}^n$	-33520#	200#	391	33	3.6 μs	0.4	10 <sup>+</sup> #	18	2005	IT=100	*	
$^{188}\text{W}$	-38668	3			69.77 d	0.05	0 <sup>+</sup>	18 14Un01 T	1951	$\beta^-$ =100	*	

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{188}\text{W}^m$	-36741	3	1926.7	0.8	109.5 ns 3.5	$8^-$	18	2010	IT=100
$^{188}\text{Re}$	-39016.9	0.7			17.005 h 0.003	$1^-*$	18	1939	$\beta^-$ =100
$^{188}\text{Re}^m$	-38844.8	0.7	172.0848	0.0024	18.59 m 0.04	$6^-$	18	1953	IT=100
$^{188}\text{Os}$	-41137.3	0.7			STABLE >3.3Ey	$0^+$	18	1931	IS=13.24 27; $\alpha$ ? *
$^{188}\text{Ir}$	-38345	9			41.5 h 0.5	$1^-*$	18	1950	$\beta^+$ =100
$^{188}\text{Ir}^m$	-37381	25	964	23	4.15 ms 0.15	$11^- \#$	18	1971	IT $\approx$ 100; $\beta^+$ ? *
$^{188}\text{Pt}$	-37821	5			10.16 d 0.18	$0^+$	18	1954	$\epsilon$ =100; $\alpha$ =2.6e-5 3
$^{188}\text{Au}$	-32371.3	2.7			8.84 m 0.06	$1^-*$	18	1955	$\beta^+$ =100
$^{188}\text{Hg}$	-30198	7			3.25 m 0.15	$0^+$	18	1960	$\beta^+$ =100; $\alpha$ =3.7e-5 8
$^{188}\text{Hg}^m$	-27474	7	2724.1	0.4	142 ns 14	$12^+$	18 83Ha15 T	1983	IT=100 *
$^{188}\text{Tl}$	-22336	30		*	71 s 2	$2^- \#$	18	1970	$\beta^+$ =100
$^{188}\text{Tl}^m$	-22308	9	30	30 MD*	71.5 s 1.5	$7^+*$	18 13Ba41 J	1970	$\beta^+$ =100 *
$^{188}\text{Tl}^n$	-22040	40	299	30	41 ms 4	$9^-$	18	1981	IT $\approx$ 100; $\beta^+$ ? *
$^{188}\text{Pb}$	-17811	10			25.1 s 0.1	$0^+$	18	1972	$\beta^+$ =91.5 5; $\alpha$ =8.5 5
$^{188}\text{Pb}^m$	-15234	10	2577.2	0.4	800 ns 20	$8^-$	18	1999	IT=100
$^{188}\text{Pb}^n$	-15101	10	2709.8	0.5	94 ns 12	$12^+$	18	2004	IT=100
$^{188}\text{Pb}^p$	-13028	10	4783.4	0.7	440 ns 60	$(19^-)$	18	2000	IT=100
$^{188}\text{Bi}$	-7195	11		&	60 ms 3	$(3^+)$	18 13La02 TD	1980	$\alpha$ $\approx$ 100; $\beta^+$ ?; $\beta^+$ SF=0.0014 7 *
$^{188}\text{Bi}^m$	-7130	30	66	30 AD	> 5 $\mu$ s	$7^+ \#$	18	1984	IT ?; $\alpha$ ?; $\beta^+$ ?
$^{188}\text{Bi}^n$	-7040	30	153	30 AD &	265 ms 15	$(10^-)$	18 13La02 TD	1984	$\alpha$ $\approx$ 100; $\beta^+$ ?; $\beta^+$ SF=0.0046 9 *
$^{188}\text{Po}$	-544	20			270 $\mu$ s 30	$0^+$	18	1999	$\alpha$ $\approx$ 100; $\beta^+$ ?
* $^{188}\text{Os}$	T : from 20Be23 for T1/2( $\alpha,0^+ \rightarrow 2^+$ ) **								
* $^{188}\text{Ir}^m$	E : 923.53(0.22) + x; x<80 keV **								
* $^{188}\text{Hg}^m$	T : average 83Ha15=134(15) 04G104=187(35) **								
* $^{188}\text{Tl}^m$	J : 92Me07.13Ba41=7 **								
* $^{188}\text{Tl}^n$	E : 268.8(0.2) keV above $^{188}\text{Tl}^m$ from 91Va04 **								
* $^{188}\text{Bi}$	D : % $\beta^+$ SF from 20An12=0.0004(0.0002)-0.0018(0.0007); other **								
* $^{188}\text{Bi}$	D : 13La02=0.0032(0.0016) for both $\beta^+$ SF decaying isomers **								
* $^{188}\text{Bi}$	J : from 20An12 **								
* $^{188}\text{Bi}^n$	D : % $\beta^+$ SF from 20An12; other 13La02=0.0032(0.0016) for both $\beta^+$ SF decaying **								
<hr/>									
$^{189}\text{Hf}$	-27150#	300#			400# ms >300ns	$3/2^- \#$	17 09A130 I	2009	$\beta^-$ =100
$^{189}\text{Ta}$	-31960#	200#			20# s >300ns	$7/2^+ \#$	17 99Be63 I	1999	$\beta^-$ =100
$^{189}\text{Ta}^m$	-30310#	220#	1650#	100#	1.6 $\mu$ s 0.2	$21/2^- \#$	09A130 TD	2009	IT=100 *
$^{189}\text{W}$	-35810#	200#			11.6 m 0.2	$9/2^- \#$	17	1963	$\beta^-$ =100
$^{189}\text{Re}$	-37979	8			24.3 h 0.4	$5/2^+$	17	1963	$\beta^-$ =100
$^{189}\text{Re}^m$	-37854	9	125	3	2# $\mu$ s	$9/2^-$	FGK209 TIJ		IT ? *
$^{189}\text{Re}^n$	-36208	8	1770.9	0.6	223 $\mu$ s 14	$29/2^+$	17 16Re02 JTE	2016	IT=100
$^{189}\text{Os}$	-38986.8	0.7			STABLE >3.3Py	$3/2^-*$	17	1931	IS=16.15 23; $\alpha$ ? *
$^{189}\text{Os}^m$	-38956.0	0.7	30.82	0.02	5.81 h 0.10	$9/2^-$	17	1960	IT $\approx$ 100; $\beta^-$ ?
$^{189}\text{Ir}$	-38450	13			13.2 d 0.1	$3/2^+*$	17	1955	$\epsilon$ =100
$^{189}\text{Ir}^m$	-38078	13	372.17	0.04	13.3 ms 0.3	$11/2^-$	17	1960	IT=100
$^{189}\text{Ir}^n$	-36117	13	2332.8	0.3	3.7 ms 0.2	$25/2^+$	17 75Ke06 J	1975	IT=100
$^{189}\text{Pt}$	-36469	10			10.87 h 0.12	$3/2^-*$	17	1955	$\beta^+$ =100 *
$^{189}\text{Pt}^m$	-36296	10	172.79	0.06	464 ns 25	$9/2^-$	17	1970	IT=100
$^{189}\text{Pt}^n$	-36278	10	191.4	0.2	143 $\mu$ s 5	$(13/2^+)$	17	1976	IT=100
$^{189}\text{Au}$	-33582	20			28.7 m 0.4	$1/2^+*$	17	1955	$\beta^+$ =100; $\alpha$ <3e-5
$^{189}\text{Au}^m$	-33335	20	247.25	0.16	4.59 m 0.11	$11/2^-*$	17	1966	$\beta^+$ $\approx$ 100;IT ?
$^{189}\text{Au}^n$	-33257	20	325.12	0.16	190 ns 15	$9/2^-$	17	1975	IT=100
$^{189}\text{Au}^p$	-31027	20	2554.8	0.8	242 ns 10	$31/2^+$	17	1975	IT=100
$^{189}\text{Hg}$	-29630	30			7.6 m 0.2	$3/2^-*$	17	1955	$\beta^+$ =100; $\alpha$ ?
$^{189}\text{Hg}^m$	-29548	18	80	30 MD	8.6 m 0.2	$13/2^+*$	17	1966	$\beta^+$ =100; $\alpha$ ? *
$^{189}\text{Tl}$	-24616	8			2.3 m 0.2	$1/2^+$	17	1972	$\beta^+$ =100
$^{189}\text{Tl}^m$	-24331	8	285	6 AD	1.4 m 0.1	$9/2^-*$	17	1972	$\beta^+$ $\approx$ 100;IT ? *
$^{189}\text{Pb}$	-17844	14			39 s 8	$3/2^-*$	17 09Sa09 T	1972	$\beta^+$ =99.58 15; $\alpha$ =0.42 15 *
$^{189}\text{Pb}^m$	-17804	14	40	4 AD	50.5 s 2.1	$13/2^+*$	17 09Sa09 T	2009	$\beta^+$ $\approx$ 100; $\alpha$ $\approx$ 0.4;IT ? *
$^{189}\text{Pb}^n$	-15369	15	2475	4	26 $\mu$ s 5	$31/2^-$	17 09Dr03 J	2005	IT=100 *
$^{189}\text{Bi}$	-10065	21			688 ms 5	$9/2^-*$	17	1973	$\alpha$ $\approx$ 100; $\beta^+$ ? *
$^{189}\text{Bi}^m$	-9881	21	184	5 AD	5.0 ms 0.1	$1/2^+$	17	1984	$\alpha$ =83 5; IT=17 5 *
$^{189}\text{Bi}^n$	-9707	21	357.6	0.5	880 ns 50	$13/2^+$	17	2001	IT=100 *
$^{189}\text{Po}$	-1422	22			3.5 ms 0.5	$(5/2^-)$	17	1999	$\alpha$ $\approx$ 100; $\beta^+$ ?

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>189</sup> Ta <sup>m</sup>	T : other 11St21=0.58(0.22), possibly a different isomer						**
* <sup>189</sup> Ta <sup>m</sup>	E : 11St11=154,284,343,389,482 keV gammas in a mutual coincidence; other						**
* <sup>189</sup> Ta <sup>m</sup>	E : 09A130=57,83,134,154,199,246,284,389,481 keV gammas in singles						**
* <sup>189</sup> Re <sup>m</sup>	IJ : M2 to 5/2+; existence of a similar isomer in <sup>187</sup> Re						**
* <sup>189</sup> Re <sup>m</sup>	T : estimated from B(M2)=0.7 (W.u.) for <sup>187</sup> Re <sup>m</sup>						**
* <sup>189</sup> Os	T : from 20Be23 for T1/2( $\alpha,3/2^- \rightarrow 3/2^-$ )						**
* <sup>189</sup> Pt	J : 75Ru06,92Hi07=3/2						**
* <sup>189</sup> Hg <sup>m</sup>	J : 79Da06=13/2						**
* <sup>189</sup> Tl <sup>m</sup>	J : 85Bo46,13Ba41=9/2						**
* <sup>189</sup> Pb	J : 09Se13=3/2						**
* <sup>189</sup> Pb	D : % $\alpha$ from 74Ho26						**
* <sup>189</sup> Pb <sup>m</sup>	T : average 09Sa09=50(3) 72Ga27=51(3)						**
* <sup>189</sup> Pb <sup>m</sup>	J : 09Se13=13/2						**
* <sup>189</sup> Pb <sup>n</sup>	E : 2434.50(0.18) keV above <sup>189</sup> Pb <sup>m</sup>						**
* <sup>189</sup> Pb <sup>n</sup>	T : symmetrized from $\tau=32(+10-2)$ us in 05Ba51						**
* <sup>189</sup> Bi	J : 17Ba12,12Ba32,95Ba75=(9/2); favored $\alpha$ decay to <sup>185</sup> Tl <sup>m</sup>						**
* <sup>189</sup> Bi	J : [J=(9/2-)]						**
* <sup>189</sup> Bi <sup>m</sup>	J : favored $\alpha$ decay to <sup>185</sup> Tl (J=1/2+)						**
* <sup>189</sup> Bi <sup>n</sup>	J : M2 to 9/2-						**
<sup>190</sup> Hf	-24800#	400#		600# ms > 300ns	0 <sup>+</sup>	13 12Ku26 I 2012	$\beta^- ?$
<sup>190</sup> Ta	-28720#	200#		5.3 s 0.7	(3)	10 09A130 TJD 2009	$\beta^- =100$
<sup>190</sup> W	-34370	40		30.0 m 1.5	0 <sup>+</sup>	20 1976	$\beta^- =100$
<sup>190</sup> W <sup>m</sup>	-32630	40	1743.6 1.0	111 ns 17	8 <sup>+</sup>	10La16 ETJ 2010	IT=100
<sup>190</sup> W <sup>n</sup>	-32530	40	1840.6 1.4	166 $\mu$ s 6	10 <sup>-</sup>	20 10La16 ETJ 2000	IT=100 *
<sup>190</sup> Re	-35583	5		3.0 m 0.2	(2) <sup>-</sup>	20 1955	$\beta^- =100$
<sup>190</sup> Re <sup>m</sup>	-35379	11	204 10	3.1 h 0.2	(6) <sup>-</sup>	20 12Re19 E 1962	$\beta^- =54.4 \text{ 20; IT}=45.6 \text{ 20}$
<sup>190</sup> Os	-38707.8	0.6		STABLE > 12Ey	0 <sup>+</sup>	20 1931	IS=26.26 20; $\alpha ?$ *
<sup>190</sup> Os <sup>m</sup>	-37002.1	0.6	1705.7 0.1	9.86 m 0.03	10 <sup>-</sup>	20 12Kr05 T 1950	IT=100 *
<sup>190</sup> Ir	-36753.6	1.4		11.78 d 0.10	4 <sup>-</sup>	20 1947	$\beta^+ =100; \alpha < 0.002$
<sup>190</sup> Ir <sup>m</sup>	-36727.5	1.4	26.1 0.1	1.120 h 0.003	1 <sup>-</sup>	20 1964	IT=100 *
<sup>190</sup> Ir <sup>n</sup>	-36717.4	1.4	36.154 0.025	> 2 $\mu$ s	4 <sup>+</sup>	20 1996	IT=100
<sup>190</sup> Ir <sup>p</sup>	-36377.2	1.4	376.4 0.1	3.087 h 0.012	11 <sup>-</sup>	20 1950	$\beta^+ =91.4 \text{ 2; IT}=8.6 \text{ 2}$ *
<sup>190</sup> Pt	-37306.5	0.7		483 Gy 3	0 <sup>+</sup>	20 FGK209 T 1949	IS=0.012 2; $\alpha =100; 2\beta^+ ?$
<sup>190</sup> Au	-32834	3		42.8 m 1.0	1 <sup>-*</sup>	20 1959	$\beta^+ =100; \alpha < 1e-6$
<sup>190</sup> Au <sup>m</sup>	-32630#	150#	200# 150# *	125 ms 20	11 <sup>-</sup> #	20 1982	IT $\approx$ 100; $\beta^+ ?$
<sup>190</sup> Hg	-31371	16		20.0 m 0.5	0 <sup>+</sup>	20 1959	$\beta^+ =100; \alpha ?$
<sup>190</sup> Tl	-24366	7		2.6 m 0.3	2 <sup>-*</sup>	20 13Ba41 J 1970	$\beta^+ =100$ *
<sup>190</sup> Tl <sup>m</sup>	-24296	5	70 7 MD	3.6 m 0.3	7 <sup>+</sup> *	20 13Ba41 J 1970	$\beta^+ =100$
<sup>190</sup> Tl <sup>n</sup>		<i>non-exist</i>		750 $\mu$ s 40	(8) <sup>-</sup>	20 1981	IT=100 *
<sup>190</sup> Tl <sup>p</sup>	-24052	16	306 10 AD	60# ms	(9) <sup>-</sup>	20 91Va04 EJT 1991	IT=100 *
<sup>190</sup> Pb	-20417	13		71 s 1	0 <sup>+</sup>	20 1972	$\beta^+ =99.60 \text{ 4; } \alpha =0.40 \text{ 4}$
<sup>190</sup> Pb <sup>m</sup>	-17802	13	2614.8 0.8	150 ns 14	10 <sup>+</sup>	20 01Dr05 J 1998	IT=100 *
<sup>190</sup> Pb <sup>n</sup>	-17750#	50#	2665# 50#	24.3 $\mu$ s 2.1	(12) <sup>+</sup>	20 1998	IT=100 *
<sup>190</sup> Pb <sup>p</sup>	-17759	13	2658.2 0.8	7.7 $\mu$ s 0.3	11 <sup>-</sup>	20 01Dr05 JT 1985	IT=100 *
<sup>190</sup> Bi	-10596	21		6.3 s 0.1	(3) <sup>+</sup> *	20 20An12 D 1972	$\alpha =77 \text{ 21; } \beta^+ =23 \text{ 21; } \beta^+ \text{ SF}=6e-6 \text{ 5}$ *
<sup>190</sup> Bi <sup>m</sup>	-10470	30	120 40 AD	6.2 s 0.1	10 <sup>-*</sup>	20 20An12 D 1988	$\alpha =70 \text{ 9; } \beta^+ ?; \beta^+ \text{ SF}=4e-6 \text{ 3}$ *
<sup>190</sup> Bi <sup>n</sup>	-10475	26	121 15	175 ns 8	(5) <sup>-</sup>	09An11 ET 2009	IT=100 *
<sup>190</sup> Bi <sup>p</sup>	-10200	50	394 40	1.3 $\mu$ s 0.8	(8) <sup>-</sup>	20 09An11 EJT 2001	IT=100 *
<sup>190</sup> Po	-4563	13		2.45 ms 0.05	0 <sup>+</sup>	20 1996	$\alpha =100; \beta^+ ?$
* <sup>190</sup> W <sup>n</sup>	T : others 11St21=108(9) 09A130=106(18) us 05Ca02=60(+1500-30) us 00Po26<3.1ms						**
* <sup>190</sup> W <sup>n</sup>	E : other 00Po26=2381						**
* <sup>190</sup> Os	T : from 20Be23 for T1/2( $\alpha,0^+ \rightarrow 2^+$ )						**
* <sup>190</sup> Os <sup>m</sup>	J : M2 + E3 to the 8+ member of the K=0+ gs band						**
* <sup>190</sup> Ir <sup>n</sup>	J : M3 to 4-; $l(d,t)=1$						**
* <sup>190</sup> Ir <sup>p</sup>	J : M4 to 7+; conf=p11/2[505]n11/2[615]; log ft=4.94 to <sup>190</sup> Os <sup>m</sup>						**
* <sup>190</sup> Ir <sup>p</sup>	J : [J=10-], conf=n9/2[505]n11/2[615], consistent with						**
* <sup>190</sup> Ir <sup>p</sup>	J : n9/2[505] $\rightarrow$ p11/2-[505] transition						**
* <sup>190</sup> Tl	J : also 92Me07=2						**
* <sup>190</sup> Tl <sup>m</sup>	J : also 92Me07=7						**
* <sup>190</sup> Tl <sup>n</sup>	I : introduced in 81Kr20, but not confirmed in 91Va04 and 05Xi06						**

Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>190</sup> Tl <sup>p</sup>	E : 236(7) keV above <sup>190</sup> Tl <sup>m</sup>						**
* <sup>190</sup> Tl <sup>p</sup>	T : from BM2=5.2(0.6)e-6 (W.u.) for a similar isomer in <sup>188</sup> Tl; other						**
* <sup>190</sup> Tl <sup>p</sup>	T : 91Va04>1 us						**
* <sup>190</sup> Pb <sup>m</sup>	T : uncertainty from 12Dr.A						**
* <sup>190</sup> Pb <sup>m</sup>	E : x keV above <sup>190</sup> Pb <sup>m</sup> ; x=50#(50#)						**
* <sup>190</sup> Pb <sup>n</sup>	T : uncertainty from 12Dr.A						**
* <sup>190</sup> Pb <sup>p</sup>	T : average 01Dr05=7.2(0.6) 85St16=7.9(0.4)						**
* <sup>190</sup> Bi	D : % $\alpha$ symmetrized from 91Va04=90(+10-30)%; % $\beta^+$ SF from T1/2( $\beta^+$ SF)						**
* <sup>190</sup> Bi	D : 20An12=2.8(+13.4-2.1)e7 s and T1/2 20An12=5.7(8)s						**
* <sup>190</sup> Bi	J : 17Ba12=(3); favored $\alpha$ decay to J=3+ in <sup>186</sup> Tl						**
* <sup>190</sup> Bi	T : other 13Ny01=7.7(+1.0-0.8) not used						**
* <sup>190</sup> Bi <sup>m</sup>	T : also 13Ny01=5.9(+1.0-0.8) not used						**
* <sup>190</sup> Bi <sup>m</sup>	J : 17Ba12=(10); favored $\alpha$ decay to <sup>186</sup> Tl <sup>n</sup> (J=10-)						**
* <sup>190</sup> Bi <sup>m</sup>	D : % $\beta^+$ SF from T1/2( $\beta^+$ SF) 20An12=4.7(+22.6-3.5)e7 s and T1/2 20An12=5.9(6)s						**
* <sup>190</sup> Bi <sup>n</sup>	J : E1 and M1(+E2) gammas in cascade to (3+), absence of gamma to (3+)						**
* <sup>190</sup> Bi <sup>n</sup>	E : 45(15) + 76 keV above <sup>190</sup> Bi						**
* <sup>190</sup> Bi <sup>p</sup>	E : 274(1) keV above <sup>190</sup> Bi <sup>m</sup>						**
* <sup>190</sup> Bi <sup>p</sup>	T : symmetrized from 09An11=1.0(+1.0-0.5)						**
<sup>191</sup> Ta	-26520# 300#		460# ms >300ns	7/2 <sup>+</sup> #	11 09St16 I	2009	$\beta^-$ ?
<sup>191</sup> W	-31180 40		14# s >300ns	3/2 <sup>-</sup> #	07 99Be63 I	1999	$\beta^-$ ?
<sup>191</sup> W <sup>m</sup>	-30950# 40#	235# 10#	340 ns 14	9/2 <sup>-</sup> #	11St21 ETD2009		IT=100
<sup>191</sup> Re	-34350 10		9.8 m 0.5	(3/2 <sup>+</sup> )	07 97Hi06 J	1963	$\beta^-$ =100
<sup>191</sup> Re <sup>m</sup>	-34205 10	145 3	20# $\mu$ s	9/2 <sup>-</sup>	FGK209 TIJ		IT ?
<sup>191</sup> Re <sup>m</sup>	-32749 10	1601.5 0.4	50.6 $\mu$ s 3.5	25/2 <sup>-</sup>	16Re02 EJT	2011	IT=100
<sup>191</sup> Os	-36395.2 0.7		14.99 d 0.02	9/2 <sup>-</sup>	07 12Kr05 T	1940	$\beta^-$ =100
<sup>191</sup> Os <sup>m</sup>	-36320.8 0.7	74.382 0.003	13.10 h 0.05	3/2 <sup>-</sup>	07 12Kr05 T	1952	IT=100
<sup>191</sup> Ir	-36708.8 1.3		STABLE	3/2 <sup>+</sup> *	07 12Kr05 T	1935	IS=37.3 2
<sup>191</sup> Ir <sup>m</sup>	-36537.5 1.3	171.29 0.04	4.899 s 0.023	11/2 <sup>-</sup>	07 12Dr02 ETJ	1979	IT=100
<sup>191</sup> Ir <sup>m</sup>	-34607.8 1.6	2101.0 0.9	5.7 s 0.4	31/2 <sup>(+)</sup>	07 12Dr02 ETJ	1979	IT=100
<sup>191</sup> Pt	-35698 4		2.83 d 0.02	3/2 <sup>-</sup> *	07 12Dr02 ETJ	1979	$\epsilon$ =100
<sup>191</sup> Pt <sup>m</sup>	-35597 4	100.663 0.020	> 1 $\mu$ s	9/2 <sup>-</sup>	07 12Dr02 ETJ	1979	IT=100
<sup>191</sup> Pt <sup>n</sup>	-35549 4	149.035 0.022	95 $\mu$ s 5	13/2 <sup>+</sup>	07 12Dr02 ETJ	1979	IT=100
<sup>191</sup> Au	-33798 5		3.18 h 0.08	3/2 <sup>+</sup> *	07 12Dr02 ETJ	1979	$\beta^+$ =100
<sup>191</sup> Au <sup>m</sup>	-33532 5	266.2 0.7	920 ms 110	11/2 <sup>-</sup> *	07 20Ba17 J	1971	IT=100
<sup>191</sup> Au <sup>n</sup>	-31308 5	2489.6 0.9	402 ns 20	31/2 <sup>+</sup> *	07 12Dr02 ETJ	1979	IT=100
<sup>191</sup> Hg	-30592 22		49 m 10	3/2 <sup>-</sup> *	07 12Dr02 ETJ	1979	$\beta^+$ =100; $\alpha$ ?
<sup>191</sup> Hg <sup>m</sup>	-30460 30	128 22	50.8 m 1.5	13/2 <sup>+</sup> *	07 01Sc41 E	1954	$\beta^+$ =100;IT ?; $\alpha$ ?
<sup>191</sup> Tl	-26283 7		20# m	1/2 <sup>+</sup> *	07 12Dr02 ETJ	1979	$\beta^+$ ?
<sup>191</sup> Tl <sup>m</sup>	-25986 7	297 7	5.22 m 0.16	9/2 <sup>-</sup> *	07 12Dr02 ETJ	1979	$\beta^+$ =100
<sup>191</sup> Pb	-20291 7		1.33 m 0.08	3/2 <sup>-</sup>	07 10Co13 JD	1974	$\beta^+$ $\approx$ 100; $\alpha$ =0.51 5
<sup>191</sup> Pb <sup>m</sup>	-20234 8	58 10	2.18 m 0.08	13/2 <sup>+</sup> *	07 17Al34 E	1975	$\beta^+$ $\approx$ 100; $\alpha$ $\approx$ 0.02
<sup>191</sup> Pb <sup>n</sup>	-17632 12	2659 10	180 ns 80	33/2 <sup>+</sup>	07 99La06 JT	1999	IT=100
<sup>191</sup> Bi	-13239 7		12.4 s 0.3	9/2 <sup>-</sup> *	16 12Dr02 ETJ	1972	$\alpha$ =51 10; $\beta^+$ ?
<sup>191</sup> Bi <sup>m</sup>	-12997 9	242 4	125 ms 8	1/2 <sup>+</sup>	16 12Dr02 ETJ	1972	$\alpha$ =68 5;IT ?; $\beta^+$ ?
<sup>191</sup> Bi <sup>n</sup>	-12809 7	429.7 0.5	562 ns 10	13/2 <sup>+</sup>	16 15Ny02 J	2001	IT=100
<sup>191</sup> Bi <sup>p</sup>	-11364# 26#	1875# 25#	400 ns 40	25/2 <sup>-</sup> #	16 12Dr02 ETJ	2016	IT=100
<sup>191</sup> Po	-5069 7		22 ms 1	3/2 <sup>-</sup>	07 12Dr02 ETJ	1993	$\alpha$ $\approx$ 100; $\beta^+$ ?
<sup>191</sup> Po <sup>m</sup>	-5008 12	61 11	93 ms 3	13/2 <sup>+</sup>	07 12Dr02 ETJ	1999	$\alpha$ $\approx$ 100; $\beta^+$ ?
<sup>191</sup> At	3864 16		2.1 ms 0.8	1/2 <sup>+</sup>	07 03Ke08 T	2003	$\alpha$ $\approx$ 100; $\beta^+$ ?
<sup>191</sup> At <sup>m</sup>	3922 18	58 20	2.2 ms 0.4	(7/2 <sup>-</sup> )	07 03Ke08 T	2003	$\alpha$ $\approx$ 100; $\beta^+$ ?
* <sup>191</sup> W <sup>m</sup>	T : average 11St21=360(20) 09Al30=320(20) ns						**
* <sup>191</sup> Re	J : measured ( <i>t</i> , $\alpha$ ) cross sections in 77Hi06 favor J=3/2 over 1/2						**
* <sup>191</sup> Re <sup>m</sup>	IJ : M2 to 5/2+ at 97(3) keV; existence of a similar isomer in <sup>187</sup> Re						**
* <sup>191</sup> Re <sup>m</sup>	T : estimated from B(M2)=0.7 (W.u.) from <sup>187</sup> Re						**
* <sup>191</sup> Re <sup>n</sup>	T : other 11St21=77(33) $\mu$ s						**
* <sup>191</sup> Os <sup>m</sup>	T : other 12Kr05=13.6(0.2) from the decay growth, less accurate						**
* <sup>191</sup> Os <sup>m</sup>	J : M3 + E4 to 9/2-						**
* <sup>191</sup> Ir <sup>m</sup>	J : E3 to 5/2+						**
* <sup>191</sup> Ir <sup>n</sup>	T : average 12Dr02=5.8(0.6) 79Lu01=5.5(0.7)						**
* <sup>191</sup> Ir <sup>n</sup>	E : from a least-squares fit to gamma-ray energies using data of 12Dr02						**
* <sup>191</sup> Pt	J : 92Hi07=3/2						**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)		
* <sup>191</sup> Pt <sup>m</sup>	J : E2 to 5/2-						**		
* <sup>191</sup> Pt <sup>m</sup>	T : other 2 us, estimated from B(E2)=0.088(5) (W.u.) in <sup>189</sup> Pt						**		
* <sup>191</sup> Pt <sup>n</sup>	J : M2 to 9/2-						**		
* <sup>191</sup> Au <sup>n</sup>	J : E2 to 27/2+; measured magnetic moment						**		
* <sup>191</sup> Hg <sup>m</sup>	E : original uncertainty (8 keV) increased by 20 keV for gs+m lines in trap						**		
* <sup>191</sup> Tl	J : 92Me07=1/2						**		
* <sup>191</sup> Tl <sup>m</sup>	J : 13Ba41,12Ba32=9/2						**		
* <sup>191</sup> Pb	J : favored $\alpha$ decay to <sup>187</sup> Hg <sup>m</sup> (J=3/2-)						**		
* <sup>191</sup> Pb	D : % $\alpha$ other 74Ho26=0.013(0.005)						**		
* <sup>191</sup> Pb <sup>m</sup>	J : 91Du07=13/2						**		
* <sup>191</sup> Pb <sup>n</sup>	E : 2602.31(0.24) above <sup>191</sup> Pb <sup>m</sup>						**		
* <sup>191</sup> Pb <sup>n</sup>	T : symmetrized from 99La06=150(+100-50)						**		
* <sup>191</sup> Bi	J : 17Ba12=(9/2); favored $\alpha$ decay to <sup>187</sup> Tl <sup>m</sup> (J=9/2-)						**		
* <sup>191</sup> Bi <sup>m</sup>	J : favored $\alpha$ decay to <sup>187</sup> Tl (J=1/2+)						**		
* <sup>191</sup> Bi <sup>p</sup>	E : 1825.1 + x keV; x=50#(25#) keV						**		
* <sup>191</sup> Po	J : favored $\alpha$ decay to <sup>187</sup> Pb (J=3/2-)						**		
* <sup>191</sup> Po <sup>m</sup>	J : favored $\alpha$ decay to <sup>187</sup> Pb <sup>m</sup> (J=13/2+)						**		
* <sup>191</sup> At	T : symmetrized from 03Ke08=1.7(+1.1-0.5)						**		
* <sup>191</sup> At	J : favored $\alpha$ decay to <sup>187</sup> Bi <sup>m</sup> (J=1/2+)						**		
* <sup>191</sup> At <sup>m</sup>	T : symmetrized from 03Ke08=2.1(+0.4-0.3)						**		
<sup>192</sup> Ta	-23100#	400#	2.2 s	0.7	(2)	12 09Al30 T	2009 $\beta^- = 100; \beta^- n ?$		
<sup>192</sup> W	-29620#	200#	40#	s >300ns	0 <sup>+</sup>	12	1999 $\beta^- ?$		
<sup>192</sup> Re	-31590	70	15.4 s	0.5	(0 <sup>-</sup> )	12 20Wa.A TJ	1965 $\beta^- = 100$		
<sup>192</sup> Re <sup>m</sup>	-31430	70	159	1	88 $\mu$ s	8 12 11St21 ETD	2005 IT=100		
<sup>192</sup> Re <sup>n</sup>	-31320	70	267	10	< 500 ms	12 20Wa.A IT	2012 $\beta^- ?; IT ?$		
<sup>192</sup> Os	-35882.3	2.3	STABLE	>53Ey	0 <sup>+</sup>	12 13Be07 T	1931 IS=40.78 32; 2 $\beta^- ?; \alpha ?$		
<sup>192</sup> Os <sup>m</sup>	-33866.9	2.3	2015.40	0.11	5.94 s	0.09 10 <sup>-</sup>	12 13Dr05 J	1965 IT $\approx$ 100; $\beta^- ?$	
<sup>192</sup> Os <sup>n</sup>	-31302.0	2.5	4580.3	1.0	205 ns	7 (20 <sup>+</sup> )	12 13Dr05 ETJ	2004 IT=100	
<sup>192</sup> Ir	-34835.6	1.3	73.820	d	0.014	4 <sup>+</sup> *	12 FGK209 T	1937 $\beta^- = 95.24$ 4; $\epsilon = 4.76$ 4	
<sup>192</sup> Ir <sup>m</sup>	-34778.9	1.3	56.720	0.005	1.45 m	0.05 1 <sup>-</sup>	12 1937	IT $\approx$ 100; $\beta^- = 0.0175$	
<sup>192</sup> Ir <sup>n</sup>	-34667.5	1.3	168.14	0.12	241 y	9 (11 <sup>-</sup> )	12 1959	IT=100	
<sup>192</sup> Pt	-36288.5	2.6	STABLE	>60Py	0 <sup>+</sup>	12 11Be08 T	1935 IS=0.782 24; $\alpha ?$		
<sup>192</sup> Pt <sup>m</sup>	-34116.1	2.6	2172.37	0.13	272 ns	23 10 <sup>-</sup>	12 1976	IT=100	
<sup>192</sup> Au	-32772	16	4.94	h	0.09	1 <sup>-</sup> *	12 1948	$\beta^+ = 100$	
<sup>192</sup> Au <sup>m</sup>	-32637	16	135.41	0.25	29 ms	5 <sup>+</sup>	12 1976	IT=100	
<sup>192</sup> Au <sup>n</sup>	-32340	16	431.6	0.5	160 ms	20 11 <sup>-</sup>	12 1976	IT=100	
<sup>192</sup> Hg	-32011	16	4.85	h	0.20	0 <sup>+</sup>	12 1952	$\epsilon = 100; \alpha ?$	
<sup>192</sup> Tl	-25870	30	9.6	m	0.4	2 <sup>-</sup> *	12 13Ba41 J	1961 $\beta^+ = 100$	
<sup>192</sup> Tl <sup>m</sup>	-25670	30	196	7	10.8 m	0.2 7 <sup>+</sup> *	12 13Ba41 J	1961 $\beta^+ = 100$	
<sup>192</sup> Tl <sup>n</sup>	-25420	30	447	7	296 ns	5 (8 <sup>-</sup> )	12 1980	IT=100	
<sup>192</sup> Tl <sup>p</sup>	-25695	25	180	40	AD	(3 <sup>+</sup> )	12 91Va04 E	1991 $\alpha = 100$	
<sup>192</sup> Pb	-22552	6	3.5	m	0.1	0 <sup>+</sup>	12 1974	$\beta^+ \approx 100; \alpha = 0.0059$ 7	
<sup>192</sup> Pb <sup>m</sup>	-19971	6	2581.1	0.4	166 ns	6 10 <sup>+</sup>	12 07Io03 J	1985 IT=100	
<sup>192</sup> Pb <sup>n</sup>	-19927	6	2625.1	1.1	1.09 $\mu$ s	0.04 12 <sup>+</sup>	12 07Io03 J	1979 IT=100	
<sup>192</sup> Pb <sup>p</sup>	-19809	6	2743.5	0.4	756 ns	14 11 <sup>-</sup>	12 07Io03 J	1991 IT=100	
<sup>192</sup> Bi	-13530	30	34.6	s	0.9	(3 <sup>+</sup> )*	12 1971	$\beta^+ = 88$ 5; $\alpha = 12$ 5	
<sup>192</sup> Bi <sup>m</sup>	-13398	9	140	30	MD	39.6 s	0.4 10 <sup>-</sup> *	12 1966	$\beta^+ = 90$ 3; $\alpha = 10$ 3
<sup>192</sup> Po	-8066	11	32.2	ms	0.3	0 <sup>+</sup>	12 1977	$\alpha \approx 100; \beta^+ ?$	
<sup>192</sup> Po <sup>m</sup>	-5771	11	2294.6	1.0	580 ns	100 11 <sup>-</sup>	12 1999	IT=100	
<sup>192</sup> At	2926	28	11.5	ms	0.6	3 <sup>+</sup> #	12 13An03 D	2006 $\alpha = 100; \beta^+ ?; \beta^+ SF < 0.51$	
<sup>192</sup> At <sup>m</sup>	2926	28	0	40	AD*&	88 ms	6 (9 <sup>-</sup> , 10 <sup>-</sup> )	12 13An03 DT	2006 $\alpha = 100; \beta^+ ?; \beta^+ SF < 0.51$
* <sup>192</sup> Re	T : average 20Wa.A=15.1(0.6) 12Al05=16(2) 79Ka.B=16(1)						**		
* <sup>192</sup> Re <sup>m</sup>	T : average 11St21=85(10) 09Al30=93(15); other 05Ca02=120(+210-50)us						**		
* <sup>192</sup> Re <sup>n</sup>	E : 159.3 keV gamma and X rays seen only in 11St21						**		
* <sup>192</sup> Re <sup>n</sup>	T : not observed in 20Wa.A, based on the extraction time of the isotope						**		
* <sup>192</sup> Re <sup>n</sup>	T : separation system; other 12Re19=61(+40-20) s for q=75+ (bare ions)						**		
* <sup>192</sup> Os	T : lower limit is for 2 $\beta^-$ ; T1/2( $\alpha, 0^+ \rightarrow 2^+$ ) 20Be23 >5.8Ey						**		
* <sup>192</sup> Os <sup>m</sup>	T : average 79KaYT=5.9(0.1) 73Pa21=6.1(0.2); other 15Ak02=10.5(+1.0-0.9) s						**		
* <sup>192</sup> Os <sup>m</sup>	T : from $\tau=15.1(+1.5-1.3)$ s for q=75+ (H-like)						**		
* <sup>192</sup> Os <sup>n</sup>	T : from 13Dr05 $\tau=295(10)$ ns						**		
* <sup>192</sup> Pt <sup>m</sup>	J : E2 to 8- band member						**		

Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)

Nuclide	Mass excess (keV)	Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>192</sup> Au <sup>m</sup>				J : E3 to 2-; M2 to 3-				**
* <sup>192</sup> Au <sup>n</sup>				J : E3 to 8+				**
* <sup>192</sup> Tl				J : also 92Me07=2				**
* <sup>192</sup> Tl <sup>m</sup>				J : also 92Me07=7				**
* <sup>192</sup> Tl <sup>m</sup>				E : from 91Va04=168+x keV, 15 keV <x< 40 keV				**
* <sup>192</sup> Tl <sup>n</sup>				E : 250.6(0.2) keV above <sup>192</sup> Tl <sup>m</sup>				**
* <sup>192</sup> Bi				J : 17Ba12=(3)				**
* <sup>192</sup> Bi <sup>m</sup>				J : 17Ba12=(10); favored $\alpha$ decay to <sup>188</sup> Tl (J=10-)				**
* <sup>192</sup> Po <sup>m</sup>				J : E1 to 10+				**
* <sup>192</sup> Po <sup>m</sup>				E : uncertainty estimated by the evaluator				**
* <sup>192</sup> At				D : % $\beta^+$ SF 13An03=0.42(0.09) for both isomers				**
* <sup>192</sup> At <sup>m</sup>				T : other 13An03=110(+26-18)				**
<sup>193</sup> Ta	-20810#	400#		220# ms >300ns	7/2 <sup>+</sup> #	17 12Ku26 I	2012	$\beta^-$ ?; $\beta^-$ -n ?
<sup>193</sup> W	-26190#	200#		30# s >300ns	1/2 <sup>-</sup> #	17 09St16 I	2009	$\beta^-$ ?
<sup>193</sup> Re	-30230	40		3# m >300ns	5/2 <sup>+</sup> #	17 99Be63 I	1999	$\beta^-$ ?
<sup>193</sup> Re <sup>m</sup>	-30080	40	146.0	69 $\mu$ s 6	(9/2 <sup>-</sup> )	17 11St21 ETJ	2005	IT=100
<sup>193</sup> Os	-33394.4	2.3		29.830 h 0.018	3/2 <sup>-</sup>	17	1940	$\beta^-$ =100
<sup>193</sup> Os <sup>m</sup>	-33078.8	2.3	315.6	121 ns 28	(9/2 <sup>-</sup> )	17	2011	IT=100
<sup>193</sup> Ir	-34536.3	1.3		STABLE	3/2 <sup>+</sup> *	17	1935	IS=62.7 2
<sup>193</sup> Ir <sup>m</sup>	-34456.1	1.3	80.238	10.53 d 0.04	11/2 <sup>-</sup>	17	1957	IT=100
<sup>193</sup> Ir <sup>n</sup>	-32257.4	1.4	2278.9	124.8 $\mu$ s 2.1	31/2 <sup>+</sup>	17	2012	IT=100
<sup>193</sup> Pt	-34479.7	1.4		50 y 6	1/2 <sup>-</sup> *	17	1948	$\epsilon$ =100
<sup>193</sup> Pt <sup>m</sup>	-34329.9	1.4	149.78	4.33 d 0.03	13/2 <sup>+</sup> *	17 86Sc04 J	1949	IT=100
<sup>193</sup> Au	-33405	9		17.65 h 0.15	3/2 <sup>+</sup> *	17	1948	$\beta^+$ =100; $\alpha$ ?
<sup>193</sup> Au <sup>m</sup>	-33115	9	290.20	3.9 s 0.3	11/2 <sup>-</sup> *	17 20Ba17 J	1955	IT $\approx$ 100; $\beta^+$ $\approx$ 0.03
<sup>193</sup> Au <sup>n</sup>	-30918	9	2486.7	150 ns 50	31/2 <sup>+</sup>	17 07Ok05 J	1985	IT=100
<sup>193</sup> Hg	-31062	16		3.80 h 0.15	3/2 <sup>-</sup> *	17	1952	$\beta^+$ =100
<sup>193</sup> Hg <sup>m</sup>	-30921	16	140.76	11.8 h 0.2	13/2 <sup>+</sup> *	17	1973	$\beta^+$ =92.8 5; IT=7.2 5
<sup>193</sup> Tl	-27477	7		21.6 m 0.8	1/2 <sup>+</sup> *	17	1960	$\beta^+$ =100
<sup>193</sup> Tl <sup>m</sup>	-27105	8	372	2.11 m 0.15	9/2 <sup>-</sup> *	17	1963	IT $\approx$ 75; $\beta^+$ $\approx$ 25
<sup>193</sup> Pb	-22229	10		4# m	3/2 <sup>-</sup> #	17	1974	$\beta^+$ =?
<sup>193</sup> Pb <sup>m</sup>	-22137	7	93	5.8 m 0.2	13/2 <sup>+</sup> *	17 17A134 E	1974	$\beta^+$ =100
<sup>193</sup> Pb <sup>n</sup>	-19522	16	2707	180 ns 15	33/2 <sup>+</sup>	17 04Io01 J	1991	IT=100
<sup>193</sup> Bi	-15885	8		63.6 s 3.0	9/2 <sup>-</sup> *	17	1971	$\beta^+$ =96.5 15; $\alpha$ =3.5 15
<sup>193</sup> Bi <sup>m</sup>	-15580	9	305	3.20 s 0.14	1/2 <sup>+</sup> *	17 15He27 T	1970	$\alpha$ =84 16; $\beta^+$ ?
<sup>193</sup> Bi <sup>n</sup>	-15279	8	605.53	153 ns 10	13/2 <sup>+</sup>	17	2004	IT=100
<sup>193</sup> Bi <sup>p</sup>	-13535	8	2349.6	85 $\mu$ s 3	29/2 <sup>+</sup>	17	2004	IT=100
<sup>193</sup> Bi <sup>q</sup>	-13480	8	2405.1	3.02 $\mu$ s 0.08	(29/2 <sup>-</sup> )	17	2004	IT=100
<sup>193</sup> Po	-8325	15		399 ms 34	3/2 <sup>-</sup> *	17	1967	$\alpha$ $\approx$ 100; $\beta^+$ ?
<sup>193</sup> Po <sup>m</sup>	-8225	15	100	245 ms 11	13/2 <sup>+</sup> *	17	1981	$\alpha$ $\approx$ 100; $\beta^+$ ?
<sup>193</sup> At	-67	22		29 ms 5	1/2 <sup>+</sup>	17 03Ke08 T	2003	$\alpha$ $\approx$ 100
<sup>193</sup> At <sup>m</sup>	-59	21	8	21 ms 5	7/2 <sup>-</sup>	17	1995	$\alpha$ $\approx$ 100
<sup>193</sup> At <sup>n</sup>	-25	21	42	28 ms 4	13/2 <sup>+</sup>	17 03Ke08 T	2003	IT=76 10; $\alpha$ =24 10
<sup>193</sup> Rn	9043	25		1.15 ms 0.27	(3/2 <sup>-</sup> )	07	2006	$\alpha$ $\approx$ 100
* <sup>193</sup> Re <sup>m</sup>				E : average 05Ca02=146.1(0.3) 11St21=145.2(0.5) 09A130=146.1(0.2) keV				**
* <sup>193</sup> Re <sup>m</sup>				T : average 11St21=65(9) 09A130=72(8); other 05Ca02=75(+450-40)				**
* <sup>193</sup> Pt				J : 92Hi07=1/2				**
* <sup>193</sup> Tl <sup>m</sup>				E : 76Ha25<13 keV above 365.2-keV level due to negligible L Xray yield				**
* <sup>193</sup> Tl <sup>m</sup>				J : 13Ba41, 12Ba32=9/2				**
* <sup>193</sup> Pb				T : 4.0 m reported in the Karlsruhe charts 1981 and 1995; not traceable				**
* <sup>193</sup> Pb <sup>m</sup>				J : 91Du07=13/2				**
* <sup>193</sup> Pb <sup>n</sup>				E : 2612.5(0.5) keV above <sup>193</sup> Pb <sup>m</sup>				**
* <sup>193</sup> Bi				J : 16Ba42=9/2				**
* <sup>193</sup> Bi <sup>m</sup>				J : 16Ba42=1/2				**
* <sup>193</sup> Po				J : 13Se03, 14Se07=(3/2); favored $\alpha$ decay to <sup>189</sup> Pb (J=3/2-)				**
* <sup>193</sup> Po <sup>m</sup>				J : 13Se03, 14Se07=(13/2); favored $\alpha$ decay to				**
* <sup>193</sup> Po <sup>m</sup>				J : <sup>189</sup> Pb <sup>m</sup> (J=13/2+)				**
* <sup>193</sup> At				T : symmetrized from 03Ke08=28(+5-4)				**
* <sup>193</sup> At				J : favored $\alpha$ decay to <sup>189</sup> Bi <sup>m</sup> (J=1/2+)				**
* <sup>193</sup> At <sup>m</sup>				J : favored $\alpha$ decay to <sup>189</sup> Bi (J=7/2-)				**
* <sup>193</sup> At <sup>n</sup>				T : symmetrized from 03Ke08=27(+4-3)				**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>193</sup> At <sup>n</sup>	J : favored $\alpha$ decay to <sup>189</sup> Bi <sup>n</sup> (J=13/2+)						**
* <sup>193</sup> Rn	J : favored $\alpha$ decay to <sup>189</sup> Po [J=(3/2-)]						**
<sup>194</sup> Ta	-17130# 500#		2# s >300ns		13 12Ku26 I	2012	$\beta^-$ ?; $\beta^-$ n ?
<sup>194</sup> W	-24410# 300#		20# s >300ns	0 <sup>+</sup>	11	2008	$\beta^-$ ?
<sup>194</sup> Re	-27260# 200#		5 s 1	1 <sup>-</sup> #	14 12Al05 T	1999	$\beta^-$ =100
<sup>194</sup> Re <sup>m</sup>	-27110# 210#	150# 50#	45 $\mu$ s 18	4 <sup>-</sup> #	14 11St21 TD	2011	IT=100
<sup>194</sup> Re <sup>n</sup>	-26980# 200#	285 40	25 s 8	11 <sup>-</sup> #	14 12Re19 E	2012	$\beta^-$ =100
<sup>194</sup> Re <sup>p</sup>	-26430# 200#	833 33	100 s 10		14 12Re19 E	2012	$\beta^-$ =100
<sup>194</sup> Os	-32435.2 2.4		6.0 y 0.2	0 <sup>+</sup>	06	1951	$\beta^-$ =100
<sup>194</sup> Ir	-32531.8 1.3		19.35 h 0.07	1 <sup>-</sup> *	06 16Kr06 T	1937	$\beta^-$ =100
<sup>194</sup> Ir <sup>m</sup>	-32384.7 1.3	147.072 0.002	31.85 ms 0.24	4 <sup>+</sup>	06	1959	IT=100
<sup>194</sup> Ir <sup>n</sup>	-32160 70	370 70 BD	171 d 11	(11 <sup>-</sup> )	06	1968	$\beta^-$ =100
<sup>194</sup> Pt	-34760.1 0.5		STABLE	0 <sup>+</sup>	06	1935	IS=32.864 410
<sup>194</sup> Au	-32211.9 2.1		38.02 h 0.10	1 <sup>-</sup> *	06	1948	$\beta^+$ =100
<sup>194</sup> Au <sup>m</sup>	-32104.5 2.2	107.4 0.5	600 ms 8	5 <sup>+</sup>	06	1975	IT=100
<sup>194</sup> Au <sup>n</sup>	-31736.1 2.2	475.8 0.6	420 ms 10	11 <sup>-</sup>	06	1953	IT=100
<sup>194</sup> Hg	-32184.0 2.9		447 y 28	0 <sup>+</sup>	06 15Do01 T	1962	$\epsilon$ =100
<sup>194</sup> Tl	-26937 14		33.0 m 0.5	2 <sup>-</sup> *	06 13Ba41 J	1960	$\beta^+$ =100; $\alpha$ ?
<sup>194</sup> Tl <sup>m</sup>	-26677 4	260 14 MD	32.8 m 0.2	7 <sup>+</sup> *	06 13Ba41 J	1960	$\beta^+$ =100
<sup>194</sup> Pb	-24208 17		10.7 m 0.6	0 <sup>+</sup>	06	1960	$\beta^+$ =100; $\alpha$ =7.3e-6 29
<sup>194</sup> Pb <sup>m</sup>	-21580 17	2628.1 0.4	370 ns 13	12 <sup>+</sup>	06 FGK128 J	1972	IT=100
<sup>194</sup> Pb <sup>n</sup>	-21275 17	2933.0 0.4	133 ns 7	11 <sup>-</sup>	06	1986	IT=100
<sup>194</sup> Bi	-16023 5		95 s 3	3 <sup>+</sup> *	06	1971	$\beta^+$ $\approx$ 100; $\alpha$ =0.46 25
<sup>194</sup> Bi <sup>m</sup>	-15880 50	150 50 MD	125 s 2	(6 <sup>+</sup> , 7 <sup>+</sup> )	06	1976	$\beta^+$ $\approx$ 100; $\alpha$ ?
<sup>194</sup> Bi <sup>n</sup>	-15860 5	163 4 AD	115 s 4	10 <sup>-</sup> *	06	1988	$\beta^+$ $\approx$ 100; $\alpha$ =0.20 7
<sup>194</sup> Po	-11005 13		392 ms 4	0 <sup>+</sup>	06	1967	$\alpha$ $\approx$ 100; $\beta^+$ ?
<sup>194</sup> Po <sup>m</sup>	-8692 13	2313.4 0.3	12.9 $\mu$ s 0.5	(10 <sup>-</sup> )	06 16An10 TJE	1999	IT=100
<sup>194</sup> At	-716 24		286 ms 7	(5 <sup>-</sup> )	06 13An03 TD	2009	$\alpha$ $\approx$ 100; $\beta^+$ =8.3#; $\beta^+$ SF=0.032#
<sup>194</sup> At <sup>m</sup>	-740 30	-20 40 AD*	323 ms 7	10 <sup>-</sup>	06 13An03 T	1984	$\alpha$ $\approx$ 100; $\beta^+$ =8.3#; $\beta^+$ SF=0.032#
<sup>194</sup> Rn	5725 17		780 $\mu$ s 160	0 <sup>+</sup>	07	2006	$\alpha$ $\approx$ 100; $\beta^+$ ?
* <sup>194</sup> Re	T : other 09Ku28=1.0(0.5) withdrawn by authors in 14Ku23						**
* <sup>194</sup> Re <sup>m</sup>	E : only 86.3 keV gamma is seen in 11St21						**
* <sup>194</sup> Re <sup>m</sup>	I : assignment from 11St21; similar experiment, but with less statistics,						**
* <sup>194</sup> Re <sup>m</sup>	I : in 05Ca02 also reports a us isomer with 464, 148, 128 gammas						**
* <sup>194</sup> Re <sup>m</sup>	I : labeled in a singles spectrum, among others, on the top of high						**
* <sup>194</sup> Re <sup>m</sup>	I : background; the assignment of these gammas to <sup>194</sup> Re is ambiguous						**
* <sup>194</sup> Re <sup>n</sup>	T : associated with 194,349 and 554 keV gammas following $\beta^-$ decay and						**
* <sup>194</sup> Re <sup>n</sup>	T : placed in 12Al05 in the high-spin part of <sup>194</sup> Os level scheme						**
* <sup>194</sup> Ir	T : average 16Kr06=19.20(0.02) 72Ge10=19.15(0.03) 72Em01=19.41(0.01);						**
* <sup>194</sup> Ir	T : Birge ratio=8.27						**
* <sup>194</sup> Ir <sup>n</sup>	J : direct $\beta^-$ feeding to J=10+ and no feeding to 8+ and 9-; systematics						**
* <sup>194</sup> Au <sup>m</sup>	J : M2 to 3- member of K=1- gs band						**
* <sup>194</sup> Au <sup>n</sup>	J : E3 to 8+						**
* <sup>194</sup> Hg	T : average 81Ho18=477(32) 79Pr15=358(55), values corrected in 15Do01 for						**
* <sup>194</sup> Hg	T : the new branching intensity of the 328.5 keV gamma ray.						**
* <sup>194</sup> Tl	J : also 92Me07=2						**
* <sup>194</sup> Tl <sup>m</sup>	J : also 92Me07=7						**
* <sup>194</sup> Pb <sup>m</sup>	J : E2 to 10+; magnetic moment						**
* <sup>194</sup> Pb <sup>n</sup>	J : E2 to 9-; magnetic moment						**
* <sup>194</sup> Bi	J : 17Ba12=(3); favored $\alpha$ decay to <sup>190</sup> Tl (J=3+)						**
* <sup>194</sup> Bi <sup>n</sup>	J : 17Ba12=(10); favored $\alpha$ decay from <sup>198</sup> At <sup>m</sup> (J=10-)						**
* <sup>194</sup> At	T : 13An03, supersedes 09An11=253(10)						**
* <sup>194</sup> At	D : % $\beta^+$ SF 13An03=0.065(0.008) for both isomers						**
* <sup>194</sup> At	J : favored $\alpha$ decay to <sup>190</sup> Bi <sup>n</sup> [J=(5-)]						**
* <sup>194</sup> At <sup>m</sup>	T : 13An03=323(7), supersedes 09An11=310(8); other 13Ny01=300(+50-40)						**
<sup>195</sup> W	-20740# 300#		30# s >160ns	3/2 <sup>-</sup> #	16 12Ku26 I	2012	$\beta^-$ ?
<sup>195</sup> Re	-25560# 300#		6 s 1	5/2 <sup>+</sup> #	14	2008	$\beta^-$ =100



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{195}\text{Os}$	-29510	60			6.5 m 1.1	(3/2 <sup>-</sup> )	14 13Bi14	TD 2004	$\beta^-$ =100	*
$^{195}\text{Os}^m$	-29080	60	427.8	0.3	47 s 3	(13/2 <sup>+</sup> )	14 20Wa12	ETJ 2012	IT $\approx$ 100; $\beta^-$ ?	*
$^{195}\text{Ir}$	-31692.3	1.3			2.29 h 0.17	3/2 <sup>+</sup>	14 13Bi14	TD 1952	$\beta^-$ =100	*
$^{195}\text{Ir}^m$	-31592	5	100	5	3.74 h 0.07	11/2 <sup>-</sup>	14	1968	$\beta^- \approx 100$ ; IT ?	*
$^{195}\text{Ir}^n$	-29338	6	2354	6	4.4 $\mu$ s 0.6	(27/2 <sup>+</sup> )	11St21	ETJ 2011	IT=100	*
$^{195}\text{Pt}$	-32793.9	0.5			STABLE >6.3Ey	1/2 <sup>-</sup> *	14 11Be08	T 1935	IS=33.775 240; $\alpha$ ?	*
$^{195}\text{Pt}^m$	-32534.8	0.5	259.077	0.023	4.010 d 0.005	13/2 <sup>+</sup> *	14	1960	IT=100	*
$^{195}\text{Au}$	-32567.1	1.1			186.01 d 0.06	3/2 <sup>+</sup> *	14 14Un01	T 1948	$\varepsilon$ =100	*
$^{195}\text{Au}^m$	-32248.5	1.1	318.58	0.04	30.5 s 0.2	11/2 <sup>-</sup> *	14 20Ba17	J 1952	IT=100	*
$^{195}\text{Au}^n$	-30066#	20#	2501#	20#	12.89 $\mu$ s 0.21	31/2(-)	14 13Dr01	ET 2013	IT=100	*
$^{195}\text{Hg}$	-31013	23			10.69 h 0.16	1/2 <sup>-</sup> *	14 15Do01	T 1952	$\beta^+$ =100	*
$^{195}\text{Hg}^m$	-30837	23	176.07	0.04	41.60 h 0.19	13/2 <sup>+</sup> *	14 15Do01	T 1951	IT=54.2 20; $\beta^+$ =45.8 20	*
$^{195}\text{Tl}$	-28155	11			1.16 h 0.05	1/2 <sup>+</sup> *	14	1955	$\beta^+$ =100	*
$^{195}\text{Tl}^m$	-27672	11	482.63	0.17	3.6 s 0.4	9/2 <sup>-</sup> *	14	1957	IT=100	*
$^{195}\text{Pb}$	-23738	5			15.0 m 1.4	3/2 <sup>-</sup>	14	1957	$\beta^+$ =100	*
$^{195}\text{Pb}^m$	-23535	5	202.9	0.7	IT 15.0 m 1.2	13/2 <sup>+</sup>	14 91Gr12	E 1957	$\beta^+$ =100; IT ?	*
$^{195}\text{Pb}^n$	-21979	5	1759.0	0.7	10.0 $\mu$ s 0.7	21/2 <sup>-</sup>	14	1976	IT=100	*
$^{195}\text{Pb}^p$	-20836	5	2901.7	0.8	95 ns 20	33/2 <sup>+</sup>	14	1982	IT=100	*
$^{195}\text{Bi}$	-18026	5			183 s 4	9/2 <sup>-</sup> *	14	1971	$\beta^+ \approx 100$ ; $\alpha=0.030$ 12	*
$^{195}\text{Bi}^m$	-17626	8	399	6	AD 87 s 1	1/2 <sup>+</sup> *	14	1974	$\beta^+$ =67 17; $\alpha=33$ 17	*
$^{195}\text{Bi}^n$	-15645	5	2381.0	0.5	614 ns 5	(29/2 <sup>-</sup> )	14 17He12	EJT 2003	IT=100	*
$^{195}\text{Bi}^p$	-15410	5	2615.9	0.5	1.49 $\mu$ s 0.01	29/2 <sup>+</sup>	15 17He12	ETJ 2018	IT=100	*
$^{195}\text{Po}$	-11117	6			4.64 s 0.09	3/2 <sup>-</sup> *	15	1967	$\alpha=94$ 4; $\beta^+$ ?	*
$^{195}\text{Po}^m$	-10968	7	148	9	MD 1.92 s 0.02	13/2 <sup>+</sup> *	15 17Al34	EJ 1967	$\alpha \approx 100$ ; $\beta^+$ ?; IT ?	*
$^{195}\text{At}$	-3470	10			290 ms 20	1/2 <sup>+</sup> *	14	1999	$\alpha \approx 100$ ; $\beta^+$ ?	*
$^{195}\text{At}^m$	-3441	8	29	7	AD 143 ms 3	7/2 <sup>-</sup> *	14	1995	$\alpha=88$ 4; IT=12 4; $\beta^+$ ?	*
$^{195}\text{At}^p$	-3370#	40#	100#	40#		(13/2 <sup>+</sup> )	13Uu01	J	IT ?	*
$^{195}\text{Rn}$	5050	50			*	3/2 <sup>-</sup>	14	2001	$\alpha=100$	*
$^{195}\text{Rn}^m$	5131	17	80	50	AD* 6 ms 3	13/2 <sup>+</sup>	14	2001	$\alpha=100$	*
$^{195}\text{Os}$	J : E3 from (13/2+) and subsequent E2 to the ground state in 21Wa.B									
$^{195}\text{Os}^m$	T : other 12Re19=32(+154-16) m for q=76+ (bare ion)									
$^{195}\text{Os}^n$	E : from 21Wa.B,20Wa12=427.8(0.3); other 12Re19=454(10) keV									
$^{195}\text{Ir}^m$	T : average 68Ja06,73Ja10=3.67(0.08) 68Ho01=4.00(0.15)									
$^{195}\text{Ir}^n$	E : from 78Ya03,83Ci01=100(5) keV in $^{196}\text{Pt}(t, \alpha)$ ; other									
$^{195}\text{Ir}^m$	E : 73Ja10=120(36) keV from $\beta^-$ decay end-point energies									
$^{195}\text{Ir}^n$	E : 268.4,404.4,476.4,537.8,566.7 gammas in a cascade to $^{195}\text{Ir}^m$									
$^{195}\text{Pt}$	J : 92Hi07=1/2									
$^{195}\text{Au}^n$	E : 13Dr01=2460.9 + x; x=40#(20#) estimated by Nubase									
$^{195}\text{Hg}$	T : average 15Do01=10.84(0.03) 01Li17=10.53(0.03); Birge ratio B=7.3									
$^{195}\text{Hg}^m$	T : average 15Do01=41.6(0.2) 73Vi09=41.6(0.8)									
$^{195}\text{Tl}^m$	J : 13Ba41,12Ba32=9/2									
$^{195}\text{Pb}$	T : from 82Hi04, determined to be within 1.2 m of the $^{195}\text{Pb}^m$									
$^{195}\text{Pb}$	T : half-life									
$^{195}\text{Bi}$	J : 16Ba42=9/2									
$^{195}\text{Bi}$	D : % $\alpha$ from 85Co06=0.01-0.05									
$^{195}\text{Bi}^m$	J : 16Ba42=1/2									
$^{195}\text{Bi}^n$	E : uncertainty estimated by Nubase; other Esndf14=2395.5(0.5)									
$^{195}\text{Bi}^p$	E : uncertainty estimated by Nubase									
$^{195}\text{Po}$	J : 13Se03,14Se07,17Al34=(3/2); favored $\alpha$ decay to $^{191}\text{Pb}$ (J=3/2-)									
$^{195}\text{Po}^m$	J : 13Se03,14Se07,17Al34=(13/2); favored $\alpha$ decay to $^{191}\text{Pb}^m$									
$^{195}\text{Po}^n$	J : (J=13/2+)									
$^{195}\text{At}$	J : 18Cu02=(1/2); favored $\alpha$ decay to $^{191}\text{Bi}^m$ (J=1/2+)									
$^{195}\text{At}^m$	E : Esndf14=33.0(1.0) is erroneous									
$^{195}\text{At}^n$	J : 18Cu02=(7/2); favored $\alpha$ decay to $^{191}\text{Bi}$ (J=7/2-)									
$^{195}\text{At}^p$	E : estimated 70#(40#) above $^{195}\text{At}^m$ ; 13Ny01<130 keV									
$^{195}\text{Rn}$	T : symmetrized from 01Uu01=6(+3-2)									
$^{195}\text{Rn}^m$	T : symmetrized from 01Uu01=5(+3-2)									
$^{196}\text{W}$	-18740#	400#			25# s >300ns	0 <sup>+</sup>	13 12Ku26	I 2012	$\beta^-$ ?	*
$^{196}\text{Re}$	-22360#	300#			2.4 s 1.5		13	2008	$\beta^-$ ?	*
$^{196}\text{Re}^m$	-22240#	300#	120#	40#	3.6 $\mu$ s 0.6		11St21	T 2009	IT=100	*
$^{196}\text{Os}$	-28280	40			34.9 m 0.2	0 <sup>+</sup>	17 77Ha32	T 1977	$\beta^-$ =100	*





Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
<sup>198</sup> Tl <sup>m</sup>	-26985	8	543.6	0.4	1.87 h	0.03		7 <sup>+</sup> * 16 1949 $\beta^+$ =55.9 23;IT=44.1 23	
<sup>198</sup> Tl <sup>n</sup>	-26842	8	686.8	0.5	150 ns	40		(5) <sup>+</sup> 16 77Kr04 EJT 1977 IT=100	
<sup>198</sup> Tl <sup>p</sup>	-26787	8	742.4	0.4	32.1 ms	1.0		10 <sup>-</sup> 16 FGK128 J 1975 IT=100 *	
<sup>198</sup> Pb	-26067	9			2.4 h	0.1		0 <sup>+</sup> 16 1955 $\beta^+$ =100	
<sup>198</sup> Pb <sup>m</sup>	-23926	9	2141.4	0.4	4.12 $\mu$ s	0.07		7 <sup>-</sup> 16 FGK128 J 1972 IT=100 *	
<sup>198</sup> Pb <sup>n</sup>	-23836	9	2231.4	0.5	137 ns	10		9 <sup>-</sup> 16 FGK128 J 1989 IT=100 *	
<sup>198</sup> Pb <sup>p</sup>	-23245	9	2821.7	0.6	212 ns	4		12 <sup>+</sup> 16 FGK128 J 1973 IT=100 *	
<sup>198</sup> Bi	-19374	28			10.3 m	0.3		3 <sup>+</sup> * 16 16Ly01 J 1950 $\beta^+$ =100	
<sup>198</sup> Bi <sup>m</sup>	-19085	28	290	40 MD	11.6 m	0.3		7 <sup>+</sup> * 16 16Ly01 J 1992 $\beta^+$ =100	
<sup>198</sup> Bi <sup>n</sup>	-18837	28	540	40 MD	7.7 s	0.5		10 <sup>-</sup> * 16 1972 IT=100 *	
<sup>198</sup> Po	-15473	17			1.760 m	0.024		0 <sup>+</sup> 16 1965 $\alpha$ =57 2; $\beta^+$ =43 2	
<sup>198</sup> Po <sup>m</sup>	-12907	17	2565.92	0.20	200 ns	20		11 <sup>-</sup> 16 1990 IT=100	
<sup>198</sup> Po <sup>n</sup>	-12730#	50#	2740#	50#	750 ns	50		12 <sup>+</sup> 16 90Ma14 T 1990 IT=100 *	
<sup>198</sup> At	-6709	5			4.47 s	0.05		3 <sup>+</sup> * 16 19Gh11 T 1967 $\alpha$ ≈97.0 17; $\beta^+$ ? *	
<sup>198</sup> At <sup>m</sup>	-6442	5	266.6	2.7 IT	1.23 s	0.05		10 <sup>-</sup> * 16 19Gh11 E 1967 $\alpha$ =93 4; $\beta^+$ ? *	
<sup>198</sup> Rn	-1230	13			64.4 ms	1.6		0 <sup>+</sup> 16 95Bi17 T 1984 $\alpha$ =93 7; $\beta^+$ ? *	
<sup>198</sup> Fr	9580	30		*	15 ms	3		3 <sup>+</sup> # 16 13Ka16 TD 2013 $\alpha$ ≈100	
<sup>198</sup> Fr <sup>m</sup>	9580	40	0	50 AD*	1.1 ms	0.7		(10 <sup>-</sup> ) 16 13Ka16 TD 2013 $\alpha$ ≈100 *	
* <sup>198</sup> Re	I : other 12Ku26>300 ns **								
* <sup>198</sup> Ir	T : average 20Mu16=8.9(0.4) 14Ku23=8(2) 72ScYY=8(1) 73S03=8(3); others **								
* <sup>198</sup> Ir	T : 18Hi07,18Mu.1=9.1(0.8), superseded by 18Hi07, 14Mo15=8(3), **								
* <sup>198</sup> Ir	T : same as 14Ku23 **								
* <sup>198</sup> Pt	T : Onu-BB 52Fr23>320 Ty; $\alpha$ 11Be08>470Py **								
* <sup>198</sup> Au <sup>n</sup>	J : M4 to 8+; magnetic moment **								
* <sup>198</sup> Tl <sup>p</sup>	J : E3 to 7+ **								
* <sup>198</sup> Pb <sup>m</sup>	J : E2 to 5-; magnetic moment **								
* <sup>198</sup> Pb <sup>m</sup>	T : average 87Ca23=4.19(0.10) 18La03=4.05(0.10); others (not used) **								
* <sup>198</sup> Pb <sup>m</sup>	T : 72Is01=3.7(0.3) 73Dj01=4 92Wa20 5.3 **								
* <sup>198</sup> Pb <sup>n</sup>	J : E2 to 7- **								
* <sup>198</sup> Pb <sup>p</sup>	J : E2 to 10+; magnetic moment **								
* <sup>198</sup> Pb <sup>p</sup>	T : average 87Ca23=212(4) 83St15=211(10) 18La03=212(10); others (not used) **								
* <sup>198</sup> Pb <sup>p</sup>	T : 73Pa03=221(30) 86Ho03=240(20) **								
* <sup>198</sup> Bi <sup>n</sup>	E : from 92Hu04=248.5(0.5) keV above <sup>198</sup> Bi <sup>m</sup> **								
* <sup>198</sup> Bi <sup>n</sup>	J : 17Ba12=(10); E3 to 7+ **								
* <sup>198</sup> Po <sup>n</sup>	E : 2691.86(0.20) + x keV; x=50#(50#) by Nubase **								
* <sup>198</sup> At	J : 18Cu02=(3); favored $\alpha$ decay to <sup>194</sup> Bi (J=3+) **								
* <sup>198</sup> At	D : % $\alpha$ from 95Bi.A>94 **								
* <sup>198</sup> At	T : others 14Ka23=3.0(0.1) 12Fo09=4.2(2.0) 05Uu02=3.8(0.4) 92Hu04=4.2(0.3) **								
* <sup>198</sup> At	T : 67Tr06=4.9(0.5) **								
* <sup>198</sup> At <sup>m</sup>	J : 18Cu02=(10); favored $\alpha$ decay from <sup>202</sup> Fr <sup>m</sup> (J=10-) **								
* <sup>198</sup> At <sup>m</sup>	T : average 19Gh11=1.28(0.10) 14Ka23=1.24(0.06) 05Uu02=1.04(0.15); others **								
* <sup>198</sup> At <sup>m</sup>	T : 92Hu04=1.0(0.2) 67Tr06=1.5(0.3) **								
* <sup>198</sup> At <sup>m</sup>	D : % $\alpha$ from 95Bi.A>86 **								
* <sup>198</sup> Rn	T : average 95Bi17=64(2) 90Ta30=66(+3-2) 84Ca32=50(9); others (not used) **								
* <sup>198</sup> Rn	T : 14Ka23=34(+11-7) 05Uu02=22(+110-10) **								
* <sup>198</sup> Rn	D : % $\alpha$ value quoted in 93Wa04 from a PhD thesis of M. Leino (1983) **								
* <sup>198</sup> Fr <sup>m</sup>	J : favored $\alpha$ decay to <sup>194</sup> At <sup>m</sup> (J=10-) **								
<sup>199</sup> Re	-14730#	400#			250# ms	>300ns		5/2 <sup>+</sup> # 13 12Ku26 I 2012 $\beta^-$ ?	
<sup>199</sup> Os	-20270#	200#			6 s	3		5/2 <sup>-</sup> # 07 14Ku23 T 2008 $\beta^-$ =100 *	
<sup>199</sup> Ir	-24400	40			7 s	5		3/2 <sup>+</sup> # 07 14Ku23 T 1993 $\beta^-$ =100 *	
<sup>199</sup> Pt	-27388.7	2.2			30.80 m	0.21		5/2 <sup>-</sup> * 07 1937 $\beta^-$ =100 *	
<sup>199</sup> Pt <sup>m</sup>	-26964.7	3.0	424	2	13.48 s	0.16		13/2 <sup>+</sup> * 07 18Mu.1 T 1959 IT=100 *	
<sup>199</sup> Au	-29093.8	0.5			3.139 d	0.007		3/2 <sup>+</sup> 07 1937 $\beta^-$ =100	
<sup>199</sup> Au <sup>m</sup>	-28544.9	0.5	548.9405	0.0021	440 $\mu$ s	30		11/2 <sup>-</sup> 07 1968 IT=100 *	
<sup>199</sup> Hg	-29546.1	0.5			STABLE			1/2 <sup>-</sup> * 07 1925 IS=16.94 12	
<sup>199</sup> Hg <sup>m</sup>	-29013.6	0.5	532.48	0.10	42.67 m	0.09		13/2 <sup>+</sup> * 07 1948 IT=100	
<sup>199</sup> Tl	-28059	28			7.42 h	0.08		1/2 <sup>+</sup> * 07 1949 $\beta^+$ =100	
<sup>199</sup> Tl <sup>m</sup>	-27310	28	748.87	0.06	28.4 ms	0.2		9/2 <sup>-</sup> * 07 1963 IT=100	
<sup>199</sup> Pb	-25232	7			90 m	10		3/2 <sup>-</sup> 07 1950 $\beta^+$ =100	
<sup>199</sup> Pb <sup>m</sup>	-24803	8	429.5	2.7	12.2 m	0.3		(13/2 <sup>+</sup> ) 07 1955 IT≈100; $\beta^+$ = ? *	
<sup>199</sup> Pb <sup>n</sup>	-22668	8	2563.8	2.7	10.1 $\mu$ s	0.2		(29/2 <sup>-</sup> ) 07 1981 IT=100 *	





**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)		
$^{201}\text{Rn}$	-4107	10				7.0 s 0.4	$3/2^-$	07	1967	$\alpha=?; \beta^+ ?$	*	
$^{201}\text{Rn}^m$	-3863	7	245	12	AD	3.8 s 0.1	$13/2^+$	07 17A134	E 1967	$\alpha=?; \beta^+ ?$	*	
$^{201}\text{Fr}$	3589	9				62.8 ms 1.9	$9/2^-$	07 14Ka23	TD 1980	$\alpha \approx 100; \beta^+ ?$	*	
$^{201}\text{Fr}^m$	3718	10	129	10	AD	24 ms 6	$1/2^+$	07 20Au01	T 2005	$\alpha=100$	*	
$^{201}\text{Fr}^n$	3879	9	289.5	0.4		720 ns 40	$13/2^+$	20Au01	ETJ 2014	$IT=100$	*	
$^{201}\text{Ra}$	11937	20				20 ms 30	$(3/2^-)$	14Ka23	TJ 2005	$\alpha=100$	*	
$^{201}\text{Ra}^m$	12200	26	263	26	AD	6 ms 5	$13/2^+$	07 05Uu02	T 2005	$\alpha=100$	*	
* $^{201}\text{Ir}$	J: $13\text{Mo}20=(1/2+, 3/2+, 5/2+)$ , but $3/2+$ agrees with systematics at Z=77											**
* $^{201}\text{Pt}^m$	I: floating high-spin level populated in decay of high-spin isomer											**
* $^{201}\text{Pt}^m$	I: in $11\text{St}21$ ; systematics of similar isomers in neighboring odd-N Pt											**
* $^{201}\text{Pt}^m$	I: nuclei											**
* $^{201}\text{Au}^m$	T: symmetrized from $11\text{St}21=340(+900-290)$											**
* $^{201}\text{Au}^m$	J: $l(t, \alpha)$ in 81Fl05											**
* $^{201}\text{Au}^n$	E: 378.2 keV + 638.0 keV gammas above $^{201}\text{Au}^m$											**
* $^{201}\text{Tl}^m$	J: E3 to $3/2+$											**
* $^{201}\text{Pb}^n$	E: 2917.6(0.9) + x keV; x < 70 keV in 81He07											**
* $^{201}\text{Bi}^n$	E: 1933.3(0.4) + x keV; x < 80 keV in 85Pi05											**
* $^{201}\text{Bi}^p$	E: 1972.3(0.4) + x keV; x < 80 keV in 85Pi05											**
* $^{201}\text{Bi}^q$	E: 2741.0(0.3) + x keV; x < 80 keV in 85Pi05											**
* $^{201}\text{Po}$	J: other $13\text{Se}03, 14\text{Se}07=3/2$											**
* $^{201}\text{Po}^m$	J: $13\text{Se}03, 14\text{Se}07=13/2$											**
* $^{201}\text{At}$	J: $18\text{Cu}02=(9/2)$ ; favored $\alpha$ decay to $^{197}\text{Bi}$ (J=9/2-)											**
* $^{201}\text{Rn}$	J: favored $\alpha$ decay to $^{197}\text{Pb}$ (J=3/2-)											**
* $^{201}\text{Rn}^m$	T: other $10\text{He}25=3.24(+3.24-1.08)$ ms											**
* $^{201}\text{Rn}^m$	J: favored $\alpha$ decay to $^{197}\text{Pb}^m$ (J=13/2+)											**
* $^{201}\text{Fr}$	T: average $14\text{Ka}23=64(3)$ $05\text{Uu}02=53(4)$ $05\text{De}01=67(3)$ ; others (not used)											**
* $^{201}\text{Fr}$	T: $96\text{En}01=69(+16-11)$ $80\text{Ew}03=48(15)$											**
* $^{201}\text{Fr}$	J: favored $\alpha$ decay to $^{197}\text{At}$ (J=9/2-)											**
* $^{201}\text{Fr}^m$	T: average $20\text{Au}01=37(+14-8)$ $14\text{Ka}23=8(+12-3)$ $05\text{Uu}02=19(+19-6)$											**
* $^{201}\text{Fr}^m$	J: favored $\alpha$ decay to $^{197}\text{At}^m$ (J=1/2+)											**
* $^{201}\text{Fr}^n$	T: other $14\text{Ka}23=700(+500-200)$											**
* $^{201}\text{Ra}$	T: symmetrized from $14\text{Ka}23=8(+40-4)$											**
* $^{201}\text{Ra}^m$	T: symmetrized from $05\text{Uu}02=1.6(+7.7-0.7)$											**
* $^{201}\text{Ra}^m$	J: favored $\alpha$ decay to $^{197}\text{Po}^m$ (J=13/2+)											**
$^{202}\text{Os}$	-12530#	400#				2# s >300ns	$0^+$	13	2009	$\beta^- ?$		
$^{202}\text{Ir}$	-16640#	300#				11 s 3	$(2^-)$	08 14Ku23	T 2008	$\beta^-=100$	*	
$^{202}\text{Ir}^m$	-14040#	420#	2600#	300#		3.4 $\mu\text{s}$ 0.6		11St21	TD 2011	$IT=100$	*	
$^{202}\text{Pt}$	-22692	25				44 h 15	$0^+$	08	1992	$\beta^-=100$		
$^{202}\text{Pt}^m$	-20904	25	1788.5	0.4		141 $\mu\text{s}$ 7	$(7^-)$	08 11St21	T 2005	$IT=100$		
$^{202}\text{Au}$	-24353	23				28.4 s 1.2	$(1^-)$	08	1967	$\beta^-=100$		
$^{202}\text{Hg}$	-27345.3	0.7						STABLE	08	1920	$IS=29.74$ 13	
$^{202}\text{Tl}$	-25980.4	1.8				12.31 d 0.08	$2^-*$	08	1940	$\epsilon=100$		
$^{202}\text{Tl}^m$	-25030.2	1.8	950.19	0.10		591 $\mu\text{s}$ 3	$7^+$	08	1958	$IT=100$		
$^{202}\text{Pb}$	-25941	4				52.5 ky 2.8	$0^+$	08	1954	$\epsilon=100$		
$^{202}\text{Pb}^m$	-23771	4	2169.85	0.08		3.54 h 0.02	$9^-$	08	1954	$IT=90.5$ $5; \beta^+=9.5$ 5		
$^{202}\text{Pb}^n$	-21800#	50#	4140#	50#		100 ns 3	$16^+$	08 19Ro12	T 1986	$IT=100$	*	
$^{202}\text{Pb}^p$	-20640#	50#	5300#	50#		108 ns 3	$19^-$	08 19Ro12	T 1987	$IT=100$	*	
$^{202}\text{Bi}$	-20751	14				1.72 h 0.05	$5^+*$	08	1951	$\beta^+=100; \alpha < 1e-5$		
$^{202}\text{Bi}^m$	-20126	18	625	12		3.04 $\mu\text{s}$ 0.06	$10^- \#$	08	1981	$IT=100$	*	
$^{202}\text{Bi}^n$	-18134	18	2617	12		310 ns 50	$(17^+)$	08	1981	$IT=100$	*	
$^{202}\text{Po}$	-17942	9				44.6 m 0.4	$0^+$	08	1951	$\beta^+=98.08$ 7; $\alpha=1.92$ 7		
$^{202}\text{Po}^m$	-16230	15	1712	12		110 ns 15	$8^+$	08	1971	$IT=100$	*	
$^{202}\text{At}$	-10595	28				184 s 1	$3^+*$	08 16Ly01	JD 1961	$\beta^+=88$ 7; $\alpha=12$ 7	*	
$^{202}\text{At}^m$	-10401	28	190	40	MD	182 s 2	$7^+*$	08 16Ly01	JD 1992	$\beta^+=91.5$ 15; $\alpha=8.5$ 15; $IT ?$	*	
$^{202}\text{At}^n$	-10010	28	590	40	MD	460 ms 50	$10^-$	08 16Ly01	J 1992	$IT=99.904$ 11; $\alpha=0.096$ 11; $\beta^+ ?$	*	
$^{202}\text{Rn}$	-6275	18				9.7 s 0.1	$0^+$	08	1967	$\alpha=78$ 8; $\beta^+ ?$		
$^{202}\text{Rn}^m$	-3970#	50#	2310#	50#		2.22 $\mu\text{s}$ 0.07	$11^- \#$	02Do19	T 2002	$IT=100$		
$^{202}\text{Fr}$	3102	6				372 ms 12	$3^+*$	08 14Ka23	T 1980	$\alpha \approx 100; \beta^+ ?$	*	
$^{202}\text{Fr}^m$	3359	6	257	6	AD	286 ms 13	$10^-*$	08 14Ka23	T 1980	$\alpha \approx 100; IT=?; \beta^+ ?$	*	
$^{202}\text{Ra}$	9075	15				4.1 ms 1.1	$0^+$	08 14Ka23	T 2005	$\alpha=100$	*	







**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>204</sup> At	T : other 10Ka29=9.6(2)						**
* <sup>204</sup> At	J : 18Cu02=(7)						**
* <sup>204</sup> Fr	T : average 05Uu02=1.9(0.5) 92Hu04=1.7(0.3)						**
* <sup>204</sup> Fr	J : 15Vo05,14Ly01,13Vo10=3						**
* <sup>204</sup> Fr <sup>m</sup>	T : average 13Ja06=2.6(0.3) 05Uu02=1.6(+0.5-0.3) 92Hu04=2.6(0.3)						**
* <sup>204</sup> Fr <sup>m</sup>	J : 15Vo05,14Ly01=7						**
* <sup>204</sup> Fr <sup>n</sup>	E : 276.1 keV above <sup>204</sup> Fr <sup>m</sup> from 95Bi.A						**
* <sup>204</sup> Fr <sup>n</sup>	T : 13Ja06=1.65(0.15) supersedes 05Uu02=0.8(0.2) (same group)						**
* <sup>204</sup> Fr <sup>n</sup>	J : 15Vo05=10,14Ly01=(10)						**
* <sup>204</sup> Fr <sup>n</sup>	D : % $\alpha$ from 14Ly01; other (not used) 94Le05=1.4(+0.8-0.4)						**
* <sup>204</sup> Ra	T : average 05Uu02=54(+19-11) 96Le09=59(+12-9); other 10He25=44(+44-15)						**
* <sup>204</sup> Ra	T : 95Le04=45(+55-21)						**
<sup>205</sup> Ir	-5600#	500#	1# s >300ns	3/2 <sup>+</sup> #	20 12Ku26 I	2012	$\beta^- ?; \beta^- n ?$
<sup>205</sup> Pt	-12820#	300#	2# s >300ns	9/2 <sup>+</sup> #	20	2009	$\beta^- ?$
<sup>205</sup> Au	-18570#	200#	32.0 s 1.4	3/2 <sup>+</sup> #	20	1994	$\beta^- =100$
<sup>205</sup> Au <sup>m</sup>	-17660#	200#	6 s 2	11/2 <sup>-</sup> #	20	2009	IT=?; $\beta^- =?$
<sup>205</sup> Au <sup>n</sup>	-15720#	200#	2849.7	0.4	163 ns 5	19/2 <sup>+</sup> #	20 2011 IT=100
<sup>205</sup> Hg	-22288	4	5.14 m 0.09	1/2 <sup>-</sup> *	20	1940	$\beta^- =100$ *
<sup>205</sup> Hg <sup>m</sup>	-20732	4	1556.4	0.3	1.09 ms 0.04	13/2 <sup>+</sup>	20 1985 IT=100
<sup>205</sup> Hg <sup>n</sup>	-18971	4	3316.6	0.8	5.89 $\mu$ s 0.18	(23/2 <sup>-</sup> )	20 2011 IT=100
<sup>205</sup> Tl	-23820.8	1.2	STABLE	1/2 <sup>+</sup> *	20	1931	IS=70.485 44 *
<sup>205</sup> Tl <sup>m</sup>	-20530.2	1.2	3290.61	0.17	2.6 $\mu$ s 0.2	25/2 <sup>-</sup>	20 1976 IT=100
<sup>205</sup> Tl <sup>n</sup>	-18985.2	1.9	4835.6	1.5	235 ns 10	(35/2 <sup>-</sup> )	20 2004 IT=100
<sup>205</sup> Pb	-23770.2	1.1	17.0 My0.9	5/2 <sup>-</sup>	20	1954	$\epsilon =100$
<sup>205</sup> Pb <sup>m</sup>	-23767.9	1.1	2.329	0.007	24.2 $\mu$ s 0.4	1/2 <sup>-</sup>	20 1994 IT=100
<sup>205</sup> Pb <sup>n</sup>	-22756.4	1.1	1013.85	0.03	5.55 ms 0.02	13/2 <sup>+</sup>	20 1960 IT=100
<sup>205</sup> Pb <sup>p</sup>	-20574.4	1.3	3195.8	0.6	217 ns 5	25/2 <sup>-</sup>	20 1973 IT=100
<sup>205</sup> Bi	-21066	5	14.91 d 0.07	9/2 <sup>-</sup> *	20	1951	$\beta^+ =100$
<sup>205</sup> Bi <sup>m</sup>	-19569	5	1497.17	0.09	7.9 $\mu$ s 0.7	1/2 <sup>+</sup>	20 1972 IT=100
<sup>205</sup> Bi <sup>n</sup>	-19001	5	2064.7	0.4	100 ns 6	21/2 <sup>+</sup>	20 1978 IT=100
<sup>205</sup> Bi <sup>p</sup>	-18927	5	2139.0	0.7	220 ns 25	25/2 <sup>+</sup>	20 1978 IT=100
<sup>205</sup> Po	-17521	10	1.74 h 0.08	5/2 <sup>-</sup> *	20	1951	$\beta^+ =99.960 12; \alpha =0.040 12$
<sup>205</sup> Po <sup>m</sup>	-17378	10	143.166	0.015	310 ns 60	1/2 <sup>-</sup>	20 1960 IT=100
<sup>205</sup> Po <sup>n</sup>	-16641	10	880.31	0.04	645 $\mu$ s 20	13/2 <sup>+</sup>	20 1962 IT=100
<sup>205</sup> Po <sup>p</sup>	-16060	10	1461.21	0.21	57.4 ms 0.9	19/2 <sup>-</sup>	20 1973 IT=100
<sup>205</sup> Po <sup>q</sup>	-14434	10	3087.2	0.4	115 ns 10	29/2 <sup>-</sup>	20 1985 IT=100
<sup>205</sup> At	-12985	12	26.9 m 0.8	9/2 <sup>-</sup> *	20	1951	$\beta^+ =90 2; \alpha =10 2$ *
<sup>205</sup> At <sup>m</sup>	-10645	12	2339.64	0.23	7.76 $\mu$ s 0.14	29/2 <sup>+</sup>	20 1982 IT=100
<sup>205</sup> Rn	-7710	5	170 s 4	5/2 <sup>-</sup> *	20	1967	$\beta^+ =75.4 9; \alpha =24.6 9$
<sup>205</sup> Rn <sup>m</sup>	-7053	5	> 10 s	13/2 <sup>+</sup> #	20	2010	IT $\approx$ 100; $\alpha ?; \beta^+ ?$
<sup>205</sup> Fr	-1310	8	3.90 s 0.07	9/2 <sup>-</sup> *	20	1964	$\alpha =98.5 4; \beta^+ =1.5 4$ *
<sup>205</sup> Fr <sup>m</sup>	-766	8	544.0	1.0	80 ns 20	13/2 <sup>+</sup>	20 2012 IT=100
<sup>205</sup> Fr <sup>n</sup>	-701	10	609	6	1.15 ms 0.04	(1/2 <sup>+</sup> )	20 2012 IT=100
<sup>205</sup> Ra	5804	23	220 ms 50	3/2 <sup>-</sup>	20 96Le09 T	1987	$\alpha \approx 100; \beta^+ ?$ *
<sup>205</sup> Ra <sup>m</sup>	6067	11	263	25	180 ms 50	13/2 <sup>+</sup>	20 17A134 E 1995 $\alpha \approx 100; IT ?; \beta^+ ?$ *
<sup>205</sup> Ac	14110	60	80 ms 60	9/2 <sup>-</sup>	20 14Zh03 T	2014	$\alpha \approx 100; \beta^+ ?$ *
* <sup>205</sup> Hg	T : other 10Ku02=5.61(0.38) for q=80+ (bare ion)						**
* <sup>205</sup> Tl	J : other 13Ba41,12Ba32=1/2						**
* <sup>205</sup> At	J : 18Cu02=9/2						**
* <sup>205</sup> Fr	J : 14Ly01,13Vo10,15Vo05,13Fi09=9/2						**
* <sup>205</sup> Ra	T : symmetrized from 96Le09=210(+60-40)						**
* <sup>205</sup> Ra	J : favored $\alpha$ decay to <sup>201</sup> Rn (J=3/2-)						**
* <sup>205</sup> Ra <sup>m</sup>	T : symmetrized from 96Le09=170(+60-40); other 10He25=68(+68-23)						**
* <sup>205</sup> Ra <sup>m</sup>	J : favored $\alpha$ decay to <sup>201</sup> Rn <sup>m</sup> (J=13/2+)						**
* <sup>205</sup> Ac	T : symmetrized from 14Zh03=20(+97-9)						**
* <sup>205</sup> Ac	J : favored $\alpha$ decay to <sup>201</sup> Fr (J=9/2-)						**
<sup>206</sup> Pt	-9240#	300#	500# ms >300ns	0 <sup>+</sup>	13 12Ku26 I	2012	$\beta^- ?; \beta^- n ?$
<sup>206</sup> Au	-14190#	300#	47 s 11	6 <sup>+</sup> #	16 17Ca12 TJ	2009	$\beta^- =100$ *
<sup>206</sup> Hg	-20946	20	8.32 m 0.07	0 <sup>+</sup>	08	1961	$\beta^- =100$







**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^{\pi}$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>210</sup> Rn <sup>n</sup>	E : 2147.4(0.2) keV above the 8+ level at 1664.6(0.1)								**
* <sup>210</sup> Rn <sup>p</sup>	E : 4803.7(0.4) keV above the 8+ level at 1664.6(0.1)								**
* <sup>210</sup> Fr <sup>m</sup>	E : uncertainty estimated by evaluator								**
* <sup>210</sup> Ra	T : others 07Le14=2.5(+1.4-0.7) and 3.5(+4.8-1.3) 68Lo15=3.6(0.2)								**
* <sup>210</sup> Ra	T : 67Va22=3.8(0.2)								**
* <sup>210</sup> Ra <sup>m</sup>	T : average 13Ba29=2.1(0.1) 06Ha17=2.28(0.08) 04Re04=2.1(0.1)								**
* <sup>210</sup> Ra <sup>m</sup>	T : 04He25=2.36(0.04); other 99Co13=2.24								**
* <sup>210</sup> Ac	T : average 00He17=335(+64-46) 68Va04=350(50)								**
<sup>211</sup> Hg	-390#	200#			26.4 s 8.1	9/2+#	13 17Ca12	TD 2010	$\beta^- = 100; \beta^-_n = 6.3\ 63$
<sup>211</sup> Tl	-6080	40			81 s 16	1/2+	13 14Mo02	TJ 1998	$\beta^- = 100; \beta^-_n = 2.2\ 22$
<sup>211</sup> Tl <sup>m</sup>	-4840#	110#	1244#	100#	580 ns 80	17/2+#	19Go10	ETJ 2019	IT=100
<sup>211</sup> Pb	-10493.0	2.3			36.1628 m 0.0025	9/2+*	13 17Lo.1	T 1904	$\beta^- = 100$
<sup>211</sup> Pb <sup>m</sup>	-8774	23	1719	23	159 ns 28	(27/2+)	13 05La01	JT 2005	IT=100
<sup>211</sup> Bi	-11859	5			2.14 m 0.02	9/2-*	13 18Ba03	J 1905	$\alpha \approx 100; \beta^- = 0.276\ 4$
<sup>211</sup> Bi <sup>m</sup>	-10602	11	1257	10	1.4 $\mu$ s 0.3	(25/2-)	13	1998	IT=100
<sup>211</sup> Po	-12432.5	1.3			516 ms 3	9/2+*	15	1913	$\alpha = 100$
<sup>211</sup> Po <sup>m</sup>	-10970	5	1462	5 AD	25.2 s 0.6	(25/2+)	15	1954	$\alpha = 99.984\ 4; IT = 0.016\ 4$
<sup>211</sup> Po <sup>n</sup>	-10298	5	2135	5	243 ns 21	(31/2-)	15	1998	IT $\approx$ 100; $\alpha$ ?
<sup>211</sup> Po <sup>p</sup>	-7561	6	4872	6	2.8 $\mu$ s 0.7	(43/2+)	15	1998	IT $\approx$ 100; $\alpha$ ?
<sup>211</sup> At	-11647.2	2.7			7.214 h 0.007	9/2-*	13	1940	$\epsilon = 58.20\ 8; \alpha = 41.80\ 8$
<sup>211</sup> At <sup>m</sup>	-6832.7	2.7	4814.5	0.5	4.23 $\mu$ s 0.07	(39/2-)	13	1971	IT=100
<sup>211</sup> Rn	-8755	7			14.6 h 0.2	1/2-*	13	1952	$\beta^+ = 72.6\ 17; \alpha = 27.4\ 17$
<sup>211</sup> Rn <sup>m</sup>	-7152#	16#	1603#	14#	596 ns 28	17/2-	13 81Po08	EJT 1981	IT=100
<sup>211</sup> Rn <sup>n</sup>	150#	21#	8905#	20#	201 ns 4	63/2-	13 85Po06	EJT 1981	IT=100
<sup>211</sup> Fr	-4140	12			3.10 m 0.02	9/2-*	13 05Ku06	D 1964	$\alpha = 87\ 3; \beta^+ = 13\ 3$
<sup>211</sup> Fr <sup>m</sup>	-1717	12	2423.16	0.24	146 ns 14	29/2+	13 86By01	ETJ 1986	IT=100
<sup>211</sup> Fr <sup>n</sup>	517	12	4657.3	0.4	124.5 ns 1.2	45/2-	13 16Ma41	T 1986	IT=100
<sup>211</sup> Ra	832	5			12.6 s 1.2	5/2-*	13 19Zh54	T 1967	$\alpha \approx 100; \beta^+ ?$
<sup>211</sup> Ra <sup>m</sup>	2030	5	1198.1	0.8	9.5 $\mu$ s 0.3	13/2+	13 13Ba29	T 2004	IT=100
<sup>211</sup> Ac	7140	50			213 ms 25	9/2-	13 00He17	T 1968	$\alpha \approx 100; \beta^+ ?$
<sup>211</sup> Th	13880	90			48 ms 20	5/2-#	13	1995	$\alpha \approx 100; \beta^+ ?$
<sup>211</sup> Pa	22050	70			6 ms 3	9/2-	13 20Au04	TD 2006	$\alpha \approx 100; \beta^+ ?; p ?$
* <sup>211</sup> Tl	T : average 17Ca12=77(18) 14Mo02,12Be28=88(+46-29)								**
* <sup>211</sup> Tl <sup>m</sup>	E : 144 keV + x keV; x=1100#(100#) keV by Nubase from interpretation in								**
* <sup>211</sup> Tl <sup>m</sup>	E : 19Go10 and similarity with <sup>209</sup> Tl <sup>m</sup>								**
* <sup>211</sup> Pb	T : others 16Ai01=36.164(0.013) 15Ko09=36.165(0.037)								**
* <sup>211</sup> Pb <sup>m</sup>	E : 1679.1 + x keV; x<80 keV estimated by Nubase								**
* <sup>211</sup> Po	J : 13Se03,14Se07,15Fi07=9/2								**
* <sup>211</sup> At	J : 18Cu02=9/2								**
* <sup>211</sup> Rn	J : other 83Ah03=5/2 (same group)								**
* <sup>211</sup> Rn <sup>m</sup>	E : 1577.8 + x keV; x<50 keV from 81Po08								**
* <sup>211</sup> Rn <sup>n</sup>	E : from 8880#(14#) + y keV; y<50 keV from 85Po06								**
* <sup>211</sup> Fr	J : other 14Ly01=9/2								**
* <sup>211</sup> Fr <sup>p</sup>	J : from 86By01								**
* <sup>211</sup> Ra	T : average 19Zh54=10(3) 07Le14=9(5) 68Lo15=12(2) 67Va22=15(2)								**
* <sup>211</sup> Ra	D : % $\alpha$ estimated by Nubase								**
* <sup>211</sup> Ra <sup>m</sup>	T : average 13Ba29=9.4(0.4) 06Ha17=9.7(0.6); other 04He25=4.0(0.5)								**
* <sup>211</sup> Ac	T : average 00He17=200(29) 68Va04=250(50)								**
* <sup>211</sup> Th	T : symmetrized from 95Uu01=37(+28-11); other 15Ya13=20.8(+37.9-8.2) (2 evts)								**
* <sup>211</sup> Pa	T : symmetrized from 20Au04=3.8(+4.6-1.4)								**
* <sup>211</sup> Pa	J : favored $\alpha$ decay to <sup>207</sup> Ac (J=9/2-)								**
<sup>212</sup> Hg	3020#	300#			30# s >300ns	0+	20 10Ai24	I 2010	$\beta^- ?; \beta^-_n ?$
<sup>212</sup> Tl	-1550#	200#			31 s 8	(5+)	20 17Ca12	TD 1998	$\beta^- = 100; \beta^-_n = 1.8\ 18$
<sup>212</sup> Pb	-7548.9	1.8			10.627 h 0.006	0+	20 17Ko16	T 1905	$\beta^- = 100$
<sup>212</sup> Pb <sup>m</sup>	-6213.9	2.7	1335	2	6.0 $\mu$ s 0.8	8+#	20 12Re.B	E 1998	IT=100
<sup>212</sup> Bi	-8117.9	1.9			60.55 m 0.06	1-*	20 89Ha.A	D 1905	$\beta^- = 64.06\ 6; \alpha = 35.94\ 6;$ $\beta^- \alpha \approx 0.014$
<sup>212</sup> Bi <sup>m</sup>	-7870	30	250	30 MD	25.0 m 0.2	(8-, 9-)	20	1978	$\alpha = 67\ 1; \beta^- = 33\ 1; \beta^- \alpha = 30\ 1$
<sup>212</sup> Bi <sup>n</sup>	-6639	30	1479	30 MD	7.0 m 0.3	(18-)	20 13Ch12	D 1978	$\beta^- = ?; IT ?$



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>213</sup> Tl <sup>n</sup>	E : 19Go10 and similarity with <sup>209</sup> Tl <sup>m</sup>								**	
<sup>213</sup> Bi	T : average 21Ta01=45.60(0.09) 13Ma13=45.62(0.06) 73Po16=45.59(0.06)								**	
<sup>213</sup> Bi <sup>m</sup>	E : from 12Ch19								**	
<sup>213</sup> Po	T : average 20Ko06=3.709(12) 18Al32=3.705(1) 13Su13=3.708(8); other								**	
<sup>213</sup> Po	T : 18Sa45=3.5(0.5)								**	
<sup>213</sup> At <sup>m</sup>	E : 1318.1(0.6) + x keV; 20<x<100 keV from 80Sj01								**	
<sup>213</sup> At <sup>n</sup>	E : 1615(2) + y keV above <sup>213</sup> At <sup>m</sup> ; y~50 keV in 03LaZZ								**	
<sup>213</sup> Rn	T : other 19Mi08=16(1)								**	
<sup>213</sup> Rn	J : other 83Ah03=1/2, inconsistent with the excited structures								**	
<sup>213</sup> Rn <sup>m</sup>	E : 1664.0(1.0) + x keV; x<35 keV from 88St10								**	
<sup>213</sup> Rn <sup>n</sup>	E : 522.7 keV above <sup>213</sup> Rn <sup>m</sup> from 88St10								**	
<sup>213</sup> Rn <sup>p</sup>	E : 4265 + y keV above <sup>213</sup> Rn <sup>m</sup> ; y<35 keV from 88St10								**	
<sup>213</sup> Fr	T : others: 16Pr08=33.2(2.0) and 28.4(3.5) 19Mi08=20(+48-8)								**	
<sup>213</sup> Fr	D : %β <sup>+</sup> other 17Lo13=0.25(0.15), discrepant likely due to Rn diffusion								**	
<sup>213</sup> Ac	J : 17Gr18,16Fe11=(9/2); favored α decay to <sup>209</sup> Fr (J=9/2-)								**	
<sup>213</sup> Th	J : favored α decay to <sup>209</sup> Ra (J=5/2-)								**	
<sup>213</sup> Th <sup>m</sup>	E : from 381(1) keV and 799(1) keV gammas in cascade; uncertainties								**	
<sup>213</sup> Th <sup>n</sup>	E : in gamma-ray energies were estimated by Nubase								**	
<sup>213</sup> Pa	T : average 20Au04=4.9(+5.9-1.8) 95Ni05=5.3(+4.0-1.6)								**	
<sup>213</sup> Pa	J : favored α decay to <sup>209</sup> Ac (J=9/2-)								**	
<sup>214</sup> Hg	11770#	400#			8# s >300ns	0 <sup>+</sup>	11 10Al24	I 2010	β <sup>-</sup> ?;β <sup>-</sup> n?	
<sup>214</sup> Tl	6470#	200#			11.0 s 2.4	5 <sup>+</sup> #	11 17Ca12	TD 2010	β <sup>-</sup> =100;β <sup>-</sup> n=34 12	
<sup>214</sup> Pb	-183.0	2.0			27.06 m 0.07	0 <sup>+</sup>	15	1904	β <sup>-</sup> =100	
<sup>214</sup> Pb <sup>m</sup>	1237	20	1420	20	6.2 μs 0.3	8 <sup>+</sup> #	15	2012	IT=100 *	
<sup>214</sup> Bi	-1201	11			19.9 m 0.4	1 <sup>-</sup>	09 89Ha.A	D 1904	β <sup>-</sup> =99.979 1;α=0.021 1; β <sup>-</sup> α≈0.003	
<sup>214</sup> Bi <sup>m</sup>	-660	30	539	30	> 93 s	8 <sup>-</sup> #	08Ch.A	TE 2008	β <sup>+</sup> ?; IT?	
<sup>214</sup> Po	-4470.0	1.4			163.47 μs 0.03	0 <sup>+</sup>	09 16Al28	T 1912	α=100 *	
<sup>214</sup> At	-3379	4			558 ns 10	1 <sup>-</sup>	09	1949	α=100	
<sup>214</sup> At <sup>m</sup>	-3321	8	59	9	AD	265 ns 30	09	1982	α≈100;IT?	
<sup>214</sup> At <sup>n</sup>	-3147	5	232	5	AD	760 ns 15	09	1982	α≈100;IT?	
<sup>214</sup> Rn	-4320	9			259 ns 3	0 <sup>+</sup>	09 19Pa45	T 1970	α=100 *	
<sup>214</sup> Rn <sup>m</sup>	275	9	4595.4	1.8		245 ns 30	(22 <sup>+</sup> )	09	1983	IT=100
<sup>214</sup> Fr	-958	9			5.51 ms 0.13	(1 <sup>-</sup> )*	09 19Mi06	T 1967	α=100 *	
<sup>214</sup> Fr <sup>m</sup>	-837	8	121	5	AD	3.35 ms 0.05	(8 <sup>-</sup> )	09	1962	α=100
<sup>214</sup> Fr <sup>n</sup>	-320	10	638	5		103 ns 4	(11 <sup>+</sup> )	09	1993	IT=100 *
<sup>214</sup> Fr <sup>p</sup>	5620#	100#	6577#	100#		108 ns 7	(33 <sup>+</sup> )	09	1994	IT=100 *
<sup>214</sup> Ra	93	5			2.437 s 0.016	0 <sup>+</sup>	09 15Kh09	T 1967	α=99.941 4;β <sup>+</sup> =0.059 4 *	
<sup>214</sup> Ra <sup>m</sup>	1913	5	1819.7	1.8		118 ns 7	6 <sup>+</sup>	09	2004	IT=100
<sup>214</sup> Ra <sup>n</sup>	1958	5	1865.2	1.8		67.3 μs 1.5	8 <sup>+</sup>	09	1971	IT=99.91 7;α=0.09 7
<sup>214</sup> Ra <sup>p</sup>	2776	5	2683.2	1.8		295 ns 7	11 <sup>-</sup>	09	1979	IT=100
<sup>214</sup> Ra <sup>q</sup>	3571	5	3478.4	1.8		279 ns 4	14 <sup>+</sup>	09	1979	IT=100
<sup>214</sup> Ra <sup>r</sup>	4240	5	4146.8	1.8		225 ns 4	17 <sup>-</sup>	09	1979	IT=100
<sup>214</sup> Ra <sup>s</sup>	6670	5	6577.0	1.8		128 ns 4	(25 <sup>-</sup> )	09	1992	IT=100
<sup>214</sup> Ac	6433	14			8.2 s 0.2	5 <sup>+</sup> *	09	1968	α=93 4;β <sup>+</sup> =7 4 *	
<sup>214</sup> Th	10695	11			87 ms 10	0 <sup>+</sup>	09	1968	α≈100;β <sup>+</sup> ? *	
<sup>214</sup> Th <sup>m</sup>	12876	11	2181.0	2.7		1.24 μs 0.12	8 <sup>+</sup> #	09	2007	IT=100
<sup>214</sup> Pa	19460	80			17 ms 3	7 <sup>+</sup> #	09 95Ni05	D 1995	α≈100	
<sup>214</sup> Pb <sup>m</sup>	E : 1365 + x keV; x=20-90 keV from 12Go19								**	
<sup>214</sup> Po	T : average 16Al28=163.47(0.03) 13Be31=163.6(0.3) 12Su11=164.2(0.6)								**	
<sup>214</sup> Rn	T : from 19Pa45; other 70Va13=270(20)								**	
<sup>214</sup> Fr	T : average 19Mi08=6.0(0.2) 15Kh09=5.9(0.4) 05Li17=4.6(0.7)								**	
<sup>214</sup> Fr	T : 68To10=5.0(0.2) 68Va18=5.5(0.5)								**	
<sup>214</sup> Fr	J : 16Fa11=(1)								**	
<sup>214</sup> Fr <sup>n</sup>	E : 516.6(6) keV above <sup>214</sup> Fr <sup>m</sup>								**	
<sup>214</sup> Fr <sup>p</sup>	E : 6477 + y keV; y=100#(100#) keV estimated by Nubase								**	
<sup>214</sup> Ra	T : average 15Kh09=2.36(0.06) 12No08=2.435(0.020) 73Be33=2.46(0.03)								**	
<sup>214</sup> Ac	J : 17Gr18,16Fe11=5								**	
<sup>214</sup> Ac	D : %β <sup>+</sup> from 68Va04<14 %								**	





**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>216</sup> Pb <sup>m</sup>	E : 1459 + x keV; x=20-90 keV from 12Go19						**
* <sup>216</sup> Bi	T : average 14Mo02=2.21(25) 00Ku06=2.25(0.05) 96Ry.B=2.17(0.05);						**
* <sup>216</sup> Bi	T : other 90Ru02=3.6(0.4)						**
* <sup>216</sup> Po	T : others 19Mi18=145(11) 18Ba44=136(6) 03Da24=144(8) 63Di05=145(2)						**
* <sup>216</sup> Rn	T : other 61Ru06=45(5)						**
* <sup>216</sup> Ra	T : average 19Pa45=161(11) 17Su18=167(53) 73No09=182(10)						**
* <sup>216</sup> Ac <sup>m</sup>	E : 322 + x keV, x=100#(100#) keV by Nubase						**
* <sup>216</sup> Th	T : average 19Zh54=26.3(0.5) 05Ku31=26.0(0.2) 01Ha46=25.4(0.8)						**
* <sup>216</sup> Th	T : 00He17=27.0(0.3) 68Va18=28(2); others 14Ya19=29(+13-7)						**
* <sup>216</sup> Th	T : 05Li17=30(9) 00He17=30(3)						**
* <sup>216</sup> Th <sup>m</sup>	T : average 19Zh54=126(14) 05Ku31=135(4) 01Ha46=128(8) 00He17=140(5)						**
* <sup>216</sup> Th <sup>m</sup>	J : favored $\alpha$ decay to <sup>214</sup> Rn <sup>m</sup> (J=8+)						**
* <sup>216</sup> Th <sup>n</sup>	E : 05Ku31=606.8(0.1) keV above <sup>216</sup> Th <sup>m</sup>						**
* <sup>216</sup> Th <sup>n</sup>	T : average 05Ku31=570(30) 01Ha46=615(55)						**
* <sup>216</sup> Th <sup>p</sup>	E : 05Ku31=1641.4(0.7) keV above <sup>216</sup> Th <sup>m</sup>						**
* <sup>216</sup> Pa	T : average 19Zh23=92(+50-24) 96An21=105(12); others 98Ik01=150(70-40),						**
* <sup>216</sup> Pa	T : 140(50-30) 79Sc09=170(100-40) 71Su14=200(40)						**
* <sup>216</sup> U	T : average 15Ma37=4.72(+4.72-1.57) 15De22=3.8(+8.8-3.2)						**
* <sup>216</sup> U <sup>m</sup>	T : symmetrized from 15Ma37=0.74(+1.34-0.29)						**
<sup>217</sup> Tl	18660#	400#		2# s >300ns	1/2 <sup>+</sup> #	18 10Al24 I 2010	$\beta^-$ ?; $\beta^-$ n ?
<sup>217</sup> Pb	12260#	300#		19.9 s 5.3	9/2 <sup>+</sup> #	18 17Ca12 TD 2010	$\beta^-$ =100
<sup>217</sup> Bi	8730	18		98.5 s 1.3	9/2 <sup>-</sup> #	18 1998	$\beta^-$ =100
<sup>217</sup> Bi <sup>m</sup>	10221	27	1491 20	3.0 $\mu$ s 0.2	25/2 <sup>-</sup> #	18 2012	IT=100
<sup>217</sup> Po	5883	7		1.53 s 0.05	(9/2 <sup>+</sup> )*	18 04Li28 TJ 1956	$\alpha$ =97.5 14; $\beta^-$ =2.5 14
<sup>217</sup> At	4395	5		32.6 ms 0.3	9/2 <sup>-</sup> *	18 19Ba22 J 1947	$\alpha$ =99.992 2; $\beta^-$ =0.008 2
<sup>217</sup> Rn	3659	4		593 $\mu$ s 38	9/2 <sup>+</sup>	18 18Sa45 T 1949	$\alpha$ =100
<sup>217</sup> Fr	4315	7		22 $\mu$ s 5	9/2 <sup>-</sup>	18 1968	$\alpha$ =100
<sup>217</sup> Ra	5890	7		1.95 $\mu$ s 0.12	(9/2 <sup>+</sup> )	18 19Mi08 T 1970	$\alpha$ =100
<sup>217</sup> Ac	8702	11		69 ns 4	9/2 <sup>-</sup>	18 1972	$\alpha$ $\approx$ 100; $\beta^+$ ?
<sup>217</sup> Ac <sup>m</sup>	10715	18	2012 20	740 ns 40	29/2 <sup>+</sup>	18 85De14 DJT 1973	IT=95.49 18 10; $\alpha$ =4.51 18
<sup>217</sup> Th	12206	11		248 $\mu$ s 4	9/2 <sup>+</sup> #	18 19Zh54 T 1968	$\alpha$ =100
<sup>217</sup> Th <sup>m</sup>	12879	11	673.3 0.1	141 ns 50	(15/2 <sup>-</sup> )	18 1989	IT=100
<sup>217</sup> Th <sup>n</sup>	14510	30	2307 32	71 $\mu$ s 14	(25/2 <sup>+</sup> )	05Ku31 ETJ 2002	IT=100
<sup>217</sup> Pa	17055	12		3.8 ms 0.2	9/2 <sup>-</sup>	18 1968	$\alpha$ =100; $\beta^-$ ?
<sup>217</sup> Pa <sup>m</sup>	18915	13	1860 7 AD	1.08 ms 0.03	(23/2 <sup>-</sup> )	18 1979	$\alpha$ =73 4; IT ?
<sup>217</sup> U	22970#	80#		850 $\mu$ s 710	1/2 <sup>-</sup> #	18 05Le42 T 2000	$\alpha$ $\approx$ 100; $\beta^-$ ?
* <sup>217</sup> Bi <sup>m</sup>	E : 1436 + x keV; x=20-90 keV from 12Go19						**
* <sup>217</sup> Po	T : average 03Ku25=1.53(0.03) 96Ry.B=1.47(0.05); other 04Li28=1.6(0.2)						**
* <sup>217</sup> Po	J : 15Fi07=(9/2,11/2)						**
* <sup>217</sup> Po	D : % $\beta^-$ from 77Vy02<5						**
* <sup>217</sup> At	D : % $\beta^-$ average 97Ch53=0.0067(24)% 69Le.A=0.012(4)%						**
* <sup>217</sup> Rn	T : average 18Sa45=670(60) 61Ru06=540(50)						**
* <sup>217</sup> Ra	T : average 19Mi08=2.5(0.2) 19Ya04=1.4(+0.4-0.3) 90An19=1.7(0.3)						**
* <sup>217</sup> Ra	T : 70Va13=1.6(0.2) 70To07=4(2)						**
* <sup>217</sup> Ac	T : others 19Mi08=150(+370-60) 82GoZU=75(3) 73No09=111(7)						**
* <sup>217</sup> Th	T : average 19Zh54=249(11) 15Kh09=259(12) 05Ku31=257(2) 02He29=237(2)						**
* <sup>217</sup> Th	T : 00He17=248(3) 00Ni02=261(+22-18) 73Ha32=252(7); Birge ratio=2.95;						**
* <sup>217</sup> Th	T : other 05Li17=310(70)						**
* <sup>217</sup> Th <sup>n</sup>	T : symmetrized from 05Ku31=67(+17-11); other 02Mu.A=20(5)						**
* <sup>217</sup> Th <sup>n</sup>	E : 2251.9 + x keV; x<110 keV in 05Ku31, due to the observed weak Kx rays						**
* <sup>217</sup> Pa	J : favored $\alpha$ decay to <sup>213</sup> Ac (J=9/2-)						**
* <sup>217</sup> U	T : symmetrized from 05Le42=0.19(+1.13-0.10) ms; other 00Ma65=15.6(+21.3-5.7)						**
<sup>218</sup> Tl	23710#	400#		1# s	6 <sup>+</sup> #		$\beta^-$ ?; $\beta^-$ n ?
<sup>218</sup> Pb	15630#	300#		14.8 s 6.8	0 <sup>+</sup>	19 17Ca12 TD 2009	$\beta^-$ =100
<sup>218</sup> Bi	13216	27		33 s 1	8 <sup>-</sup> #	19 04De16 JT 1998	$\beta^-$ =100
<sup>218</sup> Po	8356.7	2.0		3.097 m 0.012	0 <sup>+</sup>	19 1904	$\alpha$ =99.980 2; $\beta^-$ =0.020 2
<sup>218</sup> At	8100	12		1.28 s 0.06	(2 <sup>-</sup> , 3 <sup>-</sup> )*	19 19Ba22 J 1943	$\alpha$ $\approx$ 100; $\beta^-$ ?
<sup>218</sup> Rn	5217.4	2.3		33.75 ms 0.15	0 <sup>+</sup>	19 1948	$\alpha$ =100
<sup>218</sup> Fr	7059	4		1.4 ms 0.5	1 <sup>-</sup>	19 82Ew01 T 1949	$\alpha$ =100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{218}\text{Fr}^m$	7147	5	87	4	AD	21.9 ms 0.5	(8 <sup>-</sup> )	19 99Sh03 J 1982	$\alpha \approx 100$ ; IT ?	
$^{218}\text{Ra}$	6646	10				25.91 $\mu\text{s}$ 0.14	0 <sup>+</sup>	19	1970	$\alpha = 100$
$^{218}\text{Ac}$	10850	60			*	1.00 $\mu\text{s}$ 0.04	(1 <sup>-</sup> )	19 19Ya04 T	1970	$\alpha = 100$
$^{218}\text{Ac}^m$	10900	90	50	70	*	> 100# ns	8 <sup>-</sup> #			IT ?; $\alpha$ ?; $\beta^+$ ?
$^{218}\text{Ac}^n$	11460#	110#	607#	86#		103 ns 11	(11 <sup>+</sup> )	19	1994	IT=100
$^{218}\text{Th}$	12367	11				122 ns 5	0 <sup>+</sup>	19	1973	$\alpha = 100$
$^{218}\text{Pa}$	18650	18				108 $\mu\text{s}$ 5	8 <sup>-</sup> #	19 20Zh01 TDJ	1979	$\alpha = 100$
$^{218}\text{Pa}^m$	18731	20	81	19	AD	150 $\mu\text{s}$ 50	1 <sup>-</sup> #	19 20Zh01 TDJ	1979	$\alpha = 100$
$^{218}\text{U}$	21895	14				354 $\mu\text{s}$ 91	0 <sup>+</sup>	19 18Ya01 T	1992	$\alpha = 100$
$^{218}\text{U}^m$	24004	18	2109	17	AD	408 $\mu\text{s}$ 125	8 <sup>+</sup> #	19 18Ya01 T	2005	$\alpha \approx 100$ ; IT=?
* $^{218}\text{Bi}$	T : others 17Ca12=38.5(21.6) 14Mo02=33(6)									**
* $^{218}\text{Fr}$	T : symmetrized from 82Ew01=1.3(+0.5-0.4)									**
* $^{218}\text{Ac}$	T : average 19Ya04=1.04(0.12) 15Kh09=0.96(0.05) 89Mi17=1.06(0.09)									**
* $^{218}\text{Ac}$	T : 83Sc23=1.12(0.11) 17Su18=0.98(0.12). others 19Mi08=1.5(0.1), 1.8(0.1)									**
* $^{218}\text{Ac}^n$	E : 507.0(0.3) + x keV above $^{218}\text{Ac}^m$ ; x=50#(50#) keV by Nubase									**
* $^{218}\text{Pa}$	T : average 20Zh01=107(5) 00He17=113(10), supersedes 96An21=110(20)									**
* $^{218}\text{Pa}^m$	T : symmetrized from 20Zh01=135(+62-32)									**
* $^{218}\text{U}$	T : average 18Ya01=131(+179-48) 05Le42=510(+170-100)									**
* $^{218}\text{U}^m$	T : average 18Ya01=134(+244-53) 05Le42=560(+260-140); other (not used)									**
* $^{218}\text{U}^m$	T : 15Ma37=280(+1300-120)									**
$^{219}\text{Pb}$	20620#	400#				3# s >300ns	11/2 <sup>+</sup> #	11 10AI24 I	2009	$\beta^-$ ?
$^{219}\text{Bi}$	16320#	200#				8.7 s 2.9	9/2 <sup>-</sup> #	12 17Ca12 T	2009	$\beta^- = 100$ ; $\beta^-_n$ ?
$^{219}\text{Po}$	12681	16				10.3 m 1.0	9/2 <sup>+</sup> #	15 15Fi07 T	1998	$\beta^- = 71.8$ 20; $\alpha = 28.2$ 20
$^{219}\text{At}$	10396	3				56 s 3	(9/2 <sup>-</sup> )*	16 19Ba22 J	1953	$\alpha = 93.6$ 10; $\beta^- = 6.4$ 10
$^{219}\text{Rn}$	8829.3	2.1				3.96 s 0.01	5/2 <sup>+</sup> *	01	1903	$\alpha = 100$
$^{219}\text{Fr}$	8617	7				22.5 ms 1.7	9/2 <sup>-</sup> *	01 18Sa45 T	1948	$\alpha = 100$
$^{219}\text{Ra}$	9394	7				9 ms 2	(7/2) <sup>+</sup>	01 18Sa45 TJD	1952	$\alpha = 100$
$^{219}\text{Ra}^m$	9411	7	16.7	0.8		10 ms 3	(11/2) <sup>+</sup>	01 18Sa45 TJD	2018	$\alpha \approx 100$ ; IT ?
$^{219}\text{Ac}$	11570	50				9.4 $\mu\text{s}$ 1.0	9/2 <sup>-</sup>	01 19Mi08 T	1970	$\alpha = 100$ ; $\beta^+$ ?
$^{219}\text{Th}$	14460	60				1.023 $\mu\text{s}$ 0.018	9/2 <sup>+</sup> #	12 19Ya04 T	1973	$\alpha = 100$ ; $\beta^+$ ?
$^{219}\text{Pa}$	18580	70				56 ns 9	9/2 <sup>-</sup>	01 17Su18 TD	2005	$\alpha = 100$ ; $\beta^+$ ?
$^{219}\text{U}$	23296	13				60 $\mu\text{s}$ 7	9/2 <sup>+</sup> #	01 19Zh54 T	1993	$\alpha = 100$ ; $\beta^+$ ?
$^{219}\text{Np}$	29440	90				570 $\mu\text{s}$ 450	9/2 <sup>-</sup> #	16 18Ya01 T	2015	$\alpha = 100$
* $^{219}\text{Bi}$	T : other 12Be28=22(7)									**
* $^{219}\text{Po}$	T : from 15Fi07=620(59) s									**
* $^{219}\text{At}$	J : 19Ba22=(9/2)									**
* $^{219}\text{At}$	D : % $\alpha$ from 15Fi07									**
* $^{219}\text{Fr}$	T : average 18Sa45=28(3) 51Me10=20(2)									**
* $^{219}\text{Fr}$	J : 15De28, 20Ba29=9/2									**
* $^{219}\text{Ra}$	T : from 18Sa45=8(2) and 10(3) for E( $\alpha$ )=7.98 MeV and 7.66 MeV									**
* $^{219}\text{Ra}^m$	T : from 18Sa45 for E( $\alpha$ )=7.68 MeV									**
* $^{219}\text{Ac}$	T : average 19Mi08=7.6(+2.1-1.4) 89Mi17=11.8(1.5) 70Bo13=7(2)									**
* $^{219}\text{Th}$	T : average 19Ya04=1.03(0.03) 18Br13=0.94(0.08) 17Su18=1.09(0.08)									**
* $^{219}\text{Th}$	T : 15Kh09=0.97(0.04) 73Ha32=1.05(0.03); others 20Su02=0.94(+0.21-0.15)									**
* $^{219}\text{Th}$	T : 19Zh54=1.24(0.68-0.32)									**
* $^{219}\text{Pa}$	T : average 17Su18=60(+28-15) 87Fa.A=53(10)									**
* $^{219}\text{U}$	T : others 93An07=42(+34-13) 05Le42=80(+100-30)									**
* $^{219}\text{Np}$	T : symmetrized from 18Ya01=150(+720-70)									**
$^{220}\text{Pb}$	24130#	400#				1# s >300ns	0 <sup>+</sup>	11 10AI24 I	2010	$\beta^-$ ?
$^{220}\text{Bi}$	20960#	300#				9.5 s 5.7	1 <sup>-</sup> #	11 17Ca12 TD	2010	$\beta^- = 100$ ; $\beta^-_n$ ?
$^{220}\text{Po}$	15263	18				10# s >300ns	0 <sup>+</sup>	11 98Pf02 I	1998	$\beta^-$ ?
$^{220}\text{At}$	14376	14				3.71 m 0.04	3 <sup>-</sup> #	11	1989	$\beta^- = 92$ 2; $\alpha = 8$ 2
$^{220}\text{Rn}$	10612.0	1.8				55.6 s 0.1	0 <sup>+</sup>	11	1900	$\alpha = 100$ ; $2\beta^-$ ?
$^{220}\text{Fr}$	11482	4				27.4 s 0.3	1 <sup>+</sup> *	11	1948	$\alpha \approx 100$ ; $\beta^- = 0.35$ 5
$^{220}\text{Ra}$	10272	8				18.1 ms 1.2	0 <sup>+</sup>	11 18Sa45 T	1949	$\alpha = 100$
$^{220}\text{Ac}$	13744	6				26.36 ms 0.19	(3 <sup>-</sup> )	11 97Sh09 J	1970	$\alpha = 100$ ; $\beta^+$ ?
$^{220}\text{Th}$	14690	14				10.2 $\mu\text{s}$ 0.3	0 <sup>+</sup>	11 19Pa45 T	1973	$\alpha = 100$ ; $\epsilon$ ?
$^{220}\text{Pa}$	20278	15			*	0.85 $\mu\text{s}$ 0.06	1 <sup>-</sup> #	11 20Ma27 T	2005	$\alpha = 100$ ; $\beta^+$ ?
$^{220}\text{Pa}^m$	20304	22	26	23	AD*	410 ns 180		18Hu13 ET	2018	$\alpha = 100$

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{220}\text{Pa}^n$	20570	50	290	50	AD	260 ns 210		18Hu13 ET	2018	$\alpha=100$	*
$^{220}\text{U}$	23010#	100#				60# ns	$0^+$			$\alpha ?; \beta^+ ?$	
$^{220}\text{Np}$	30480	30				29 ns 11	$1^- \#$	19Zh23 TD	2019	$\alpha=100$	*
* $^{220}\text{Rn}$	T : other 18Ba44=58(4)										**
* $^{220}\text{Fr}$	J : other 14Ly01=1										**
* $^{220}\text{Ra}$	T : average 18Sa45=19(3) 00He17=18(2) 90An19=17(2) 61Ru06=23(5)										**
* $^{220}\text{Ac}$	T : average 90An19=26.4(0.2) 70Bo13=26.1(0.5)										**
* $^{220}\text{Th}$	T : average 19Pa45=10.4(0.4) 73Ha32=9.7(0.6)										**
* $^{220}\text{Pa}$	T : average 20Ma27=0.73(0.11) 19Ya04=0.91(0.10) 17Hu08=0.90(0.13);										**
* $^{220}\text{Pa}$	T : others 19Zh54=0.98(+0.40-0.22) 87Fa.A=0.780(0.16)										**
* $^{220}\text{Pa}^m$	T : symmetrized from 18Hu13=308(+250-95)										**
* $^{220}\text{Pa}^n$	T : symmetrized from 18Hu13=69(+330-30)										**
* $^{220}\text{Np}$	T : symmetrized 19Zh23=25(+14-7)										**
$^{221}\text{Bi}$	24200#	300#				2# s >300ns	$9/2^- \#$	11 10Al24 I	2009	$\beta^- ?; \beta^- n ?$	
$^{221}\text{Po}$	19774	20				2.2 m 0.7	$9/2^+ \#$	13	2010	$\beta^- =100$	*
$^{221}\text{At}$	16783	14				2.3 m 0.2	$3/2^- \#$	07	1989	$\beta^- =100$	
$^{221}\text{Rn}$	14471	6				25.7 m 0.5	$7/2^+ *$	07 97Li23 T	1956	$\beta^- =78 1; \alpha=22 1$	*
$^{221}\text{Fr}$	13277	5				4.801 m 0.005	$5/2^- *$	07 13Su13 T	1947	$\alpha \approx 100; \beta^- =0.0048 15; 14C=8.8e-11 11$	*
$^{221}\text{Ra}$	12964	5				25 s 4	$5/2^+ *$	07 18Sa45 T	1949	$\alpha=100; 14C=1.2e-10 9$	*
$^{221}\text{Ac}$	14530	60				52 ms 2	$9/2^- \#$	07	1968	$\alpha=100$	
$^{221}\text{Th}$	16940	8				1.75 ms 0.02	$7/2^+ \#$	07 14Lo10 T	1970	$\alpha=100$	*
$^{221}\text{Pa}$	20370	60				5.9 $\mu$ s 1.7	$9/2^-$	07	1983	$\alpha=100$	*
$^{221}\text{U}$	24520	70				660 ns 140	$9/2^+ \#$	15	2015	$\alpha \approx 100; \beta^+ ?$	
$^{221}\text{Np}$	29910#	200#				30# ns	$9/2^- \#$			$\alpha ?$	
$^{221}\text{Pu}$	35930#	300#				100# $\mu$ s	$9/2^+ \#$			$\alpha ?; SF ?$	
* $^{221}\text{Po}$	T : symmetrized from 10Ch19=112(+58-28) s										**
* $^{221}\text{Rn}$	J : other 83Ah03=5/2										**
* $^{221}\text{Fr}$	D : % $\beta^-$ from 97Ch53; % $^{14}\text{C}$ from 94Bo28										**
* $^{221}\text{Fr}$	T : average 13Su13=4.806(0.006) 10Wa42=4.768(0.017) 07Je07=4.79(0.02)										**
* $^{221}\text{Fr}$	J : other 14Ly01, 15De28=5/2										**
* $^{221}\text{Ra}$	T : average 18Sa45=16(2) 58To25=28(2) 51Me10=30(2); Birge ratio=3.79										**
* $^{221}\text{Ra}$	D : % $^{14}\text{C}$ from 94Bo28										**
* $^{221}\text{Th}$	T : average 14Lo10=1.78(0.03) 01Ko07=1.73(0.03) 70To07=1.68(0.06); others										**
* $^{221}\text{Th}$	T : 19Mi08=1.0(0.2) 19Ya04=2.28(+0.70-0.43) 05Li17=2.3(0.4)										**
* $^{221}\text{Th}$	T : 00He17=2.0(+0.3-0.2)										**
* $^{221}\text{Pa}$	T : other 19Mi08=3.5(+8.5-1.4)										**
$^{222}\text{Bi}$	28950#	300#				3# s >300ns	$1^- \#$	10Al24 I	2009	$\beta^- ?; \beta^- n ?$	
$^{222}\text{Po}$	22490	40				9.1 m 7.2	$0^+$	11	2010	$\beta^- =100$	*
$^{222}\text{At}$	20953	16				54 s 10		11	1989	$\beta^- =100$	
$^{222}\text{Rn}$	16372.0	1.9				3.8215 d 0.0002	$0^+$	11 15Be07 T	1899	$\alpha=100$	*
$^{222}\text{Fr}$	16378	7				14.2 m 0.3	$2^- *$	11	1975	$\beta^- =100$	
$^{222}\text{Ra}$	14320	4				33.6 s 0.4	$0^+$	11 12Po13 T	1948	$\alpha=100; 14C=3.0e-8 10$	*
$^{222}\text{Ac}$	16622	5			*	5.0 s 0.5	$1^-$	11	1949	$\alpha=99 1; \beta^+=1 1$	
$^{222}\text{Ac}^m$	16700	21	78	21	AD*	1.05 m 0.05	$5^+ \#$	11 72Es03 DTJ	1972	$\alpha \approx 98.6 4; \beta^+ \approx 1.4 4; IT ?$	*
$^{222}\text{Th}$	17203	10				2.24 ms 0.03	$0^+$	11	1970	$\alpha=100; \epsilon ?$	
$^{222}\text{Pa}$	22060	90				3.8 ms 0.2	$1^- \#$	11 19Mi08 T	1970	$\alpha=100$	*
$^{222}\text{U}$	24270	50				4.7 $\mu$ s 0.7	$0^+$	15	1983	$\alpha=100; \beta^+ ?$	
$^{222}\text{Np}$	31270	40				480 ns 190	$1^- \#$	20Ma27 TD	2020	$\alpha=100$	*
$^{222}\text{Pu}$	35060#	300#				10# $\mu$ s	$0^+$			$\alpha ?; SF ?$	
* $^{222}\text{Po}$	T : symmetrized from 10Ch19=145(+694-66) s										**
* $^{222}\text{Rn}$	T : rounded from 15Be07=3.82146(16stat,4syst)=3.82146(0.00016)										**
* $^{222}\text{Ra}$	T : others (not used) 95Ko54=36.17(0.10) 82Bo04=43(4)										**
* $^{222}\text{Ac}^m$	D : % $\beta^+$ from 0.7 < 72Es03 < 2										**
* $^{222}\text{Pa}$	T : average 19Mi08=4.5(0.3) 95Ni.A=3.3(0.3) 79Sc09=2.9(+0.6-0.4); other										**
* $^{222}\text{Pa}$	T : 70Bo13=5.7(0.5) conflicting (not used)										**
* $^{222}\text{Np}$	T : symmetrized from 20Ma27=380(+260-110)										**



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>225</sup> Ac	J : also favored $\alpha$ decay to J=3/2- level at 552.1 keV in <sup>221</sup> Fr						**
* <sup>225</sup> Th	J : favored $\alpha$ decay to J=3/2+ level at 321.4 keV in <sup>221</sup> Ra						**
* <sup>225</sup> Th	D : % $\alpha$ from 51Me10						**
* <sup>225</sup> Th	T : from 87Mi10; other 51Me10=8.0(0.5)						**
* <sup>225</sup> Pa	T : average 70Bo13=1.8(0.3) 78IbZZ=1.7(0.1)						**
* <sup>225</sup> U	T : average 19Mi08=63(7) 00He17=59(+5-2); others 03Ni10=135(+93-39)						**
* <sup>225</sup> U	T : 01Ku07=84(4) 94An02=68(+45-20) 92To02=95(15) 89He13=80(+40-10)						**
* <sup>225</sup> Np	T : average 15De22=3.8(+7.6-2.7)( <sup>233</sup> Bk decay) 3.3(+7.6-0.7)						**
* <sup>225</sup> Np	T : ( <sup>229</sup> Am decay); other: 19Mi08=0.31(+0.75-0.13)						**
<sup>226</sup> Po	37550#	400#	1# m >300ns	0 <sup>+</sup>	11 10A124 I	2010	$\beta^-$ ?
<sup>226</sup> At	34660#	300#	7# m >300ns	2 <sup>+</sup> #	11 10A124 I	2010	$\beta^-$ ?; $\beta^-n$ ?
<sup>226</sup> Rn	28747	10	7.4 m 0.1	0 <sup>+</sup>	96	1969	$\beta^-$ =100
<sup>226</sup> Fr	27521	6	48.5 s 0.7	1 <sup>-</sup> *	96 86Bo35 T	1969	$\beta^-$ =100
<sup>226</sup> Ra	23667.6	1.9	1.600 ky 0.007	0 <sup>+</sup>	96 90We01 D	1898	$\alpha$ =100;14C=2.6e-9 6;2 $\beta^-$ ?
<sup>226</sup> Ac	24309	3	29.37 h 0.12	(1 <sup>-</sup> )	96	1950	$\beta^-$ =83 3; $\epsilon$ =17 3; $\alpha$ =0.006 2
<sup>226</sup> Th	23198	4	30.70 m 0.03	0 <sup>+</sup>	96 01Bo11 D	1948	$\alpha$ =100;18O<3.2e-12
<sup>226</sup> Pa	26034	11	1.8 m 0.2	1 <sup>-</sup> #	96	1949	$\alpha$ =74 5; $\beta^+$ =26 5
<sup>226</sup> U	27329	11	269 ms 6	0 <sup>+</sup>	14 01Ca.B T	1973	$\alpha$ =100
<sup>226</sup> Np	32820	100	35 ms 10		96 19Mi08 T	1990	$\alpha$ =100; $\beta^+$ ?
<sup>226</sup> Pu	35630#	200#	10# ms	0 <sup>+</sup>			$\alpha$ ?;SF ?
<sup>226</sup> Am	42970#	300#	100# $\mu$ s				$\alpha$ ?;SF ?
* <sup>226</sup> Fr	T : average 75Ra03=48(1) 86Bo35=49(1)						**
* <sup>226</sup> Ra	D : % <sup>14</sup> C average 90We01=2.3(0.8)e-9% 86Ba26=2.9(1.0)e-9%						**
* <sup>226</sup> Ra	D : 85Ho21=3.2(1.6)e-9%						**
* <sup>226</sup> Th	T : from 12Po13; others 87Mi10=30.57(0.10) 95Ko54=30.83(0.01)						**
* <sup>226</sup> U	T : average 01Ca.B=258(13) 00He17=281(9) 99Gr28=260(10); other						**
* <sup>226</sup> U	T : 18Mi11=400(100)						**
* <sup>226</sup> Np	T : average 19Mi08=48(5) 90Ni05=35(10); other 95Le15=58(+70-20)						**
<sup>227</sup> Po	42280#	400#	2# s >300ns	5/2 <sup>+</sup> #	16	2010	$\beta^-$ ?
<sup>227</sup> At	37430#	300#	5# s >300ns	1/2 <sup>+</sup> #	16	2010	$\beta^-$ ?; $\beta^-n$ ?
<sup>227</sup> Rn	32886	14	20.2 s 0.4	(3/2 <sup>+</sup> )*	16 83Ah03 J	1986	$\beta^-$ =100
<sup>227</sup> Fr	29682	6	2.47 m 0.03	1/2 <sup>+</sup> *	16	1972	$\beta^-$ =100
<sup>227</sup> Ra	27177.5	1.9	42.2 m 0.5	3/2 <sup>+</sup> *	16	1953	$\beta^-$ =100
<sup>227</sup> Ac	25849.5	1.9	21.772 y 0.003	3/2 <sup>-</sup> *	16	1851	$\beta^-$ =98.62 36; $\alpha$ =1.38 36
<sup>227</sup> Th	25804.8	2.1	18.693 d 0.004	(1/2 <sup>+</sup> )	16 19Ko06 T	1906	$\alpha$ =100
<sup>227</sup> Pa	26830	7	38.3 m 0.3	(5/2 <sup>-</sup> )	16	1948	$\alpha$ =85 2; $\epsilon$ =15 2
<sup>227</sup> U	29045	9	1.1 m 0.1	(3/2 <sup>+</sup> )	16	1952	$\alpha$ =100; $\beta^+$ ?
<sup>227</sup> Np	32580	80	510 ms 60	5/2 <sup>+</sup> #	16	1990	$\alpha$ $\approx$ 100; $\beta^+$ ?
<sup>227</sup> Pu	36770#	100#	2# s	5/2 <sup>+</sup> #			$\alpha$ ?
<sup>227</sup> Am	42180#	200#	20# ms	9/2 <sup>-</sup> #			$\alpha$ ?;SF ?
* <sup>227</sup> Ra	J : 18Ly01,88Ah02,83Ah03,87We03=3/2						**
* <sup>227</sup> Ac	J : other 17Gr18,16Fe11=3/2						**
* <sup>227</sup> Th	T : average 19Ko06=18.681(0.009) 15Co11=18.695(0.004)						**
<sup>228</sup> At	41880#	400#	1# m >300ns	3 <sup>+</sup> #	14 10A124 I	2010	$\beta^-$ ?; $\beta^-n$ ?
<sup>228</sup> Rn	35243	18	65 s 2	0 <sup>+</sup>	14	1989	$\beta^-$ =100
<sup>228</sup> Fr	33384	7	38 s 1	2 <sup>-</sup> *	14	1972	$\beta^-$ =100
<sup>228</sup> Fr <sup>m</sup>	34390	30	1004 30	180 s 110	08Ch.A TIE	2008	IT ?; $\beta^-$ ?
<sup>228</sup> Ra	28940.2	2.0	5.75 y 0.03	0 <sup>+</sup>	14	1907	$\beta^-$ =100
<sup>228</sup> Ac	28894.7	2.1	6.15 h 0.02	3 <sup>+</sup> *	14	1908	$\beta^-$ =100
<sup>228</sup> Ac <sup>m</sup>	29430	30	539 30	180 s 70	08Ch.A TIE	2008	IT ?; $\beta^-$ ?
<sup>228</sup> Th	26770.9	1.8	1.9125 y 0.0007	0 <sup>+</sup>	14 93Bo20 D	1905	$\alpha$ =100;20O=1.13e-11 22
<sup>228</sup> Pa	28924	4	22 h 1	3 <sup>+</sup>	14	1948	$\beta^+$ =98.15 17; $\alpha$ =1.85 17
<sup>228</sup> U	29220	13	9.1 m 0.2	0 <sup>+</sup>	14 61Ru05 TD	1949	$\alpha$ =97.5 1.4; $\epsilon$ =2.5 14
<sup>228</sup> Np	33830#	100#	61.4 s 1.4	4 <sup>+</sup> #	14 94Kr13 D	1994	$\epsilon$ =59 7; $\alpha$ =41 7; $\beta^+$ SF=0.012 6
<sup>228</sup> Pu	36108	23	2.1 s 1.3	0 <sup>+</sup>	14 03Ni10 T	1994	$\alpha$ $\approx$ 100; $\beta^+$ ?

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
<sup>228</sup> Am	42850# 200#		100# ms				$\alpha$ ?;SF ?
* <sup>228</sup> Fr <sup>m</sup>	T : symmetrized from 08Ch.A=94(+170-29) s						
* <sup>228</sup> Ac <sup>m</sup>	T : symmetrized from 08Ch.A=149(+95-42) s						
* <sup>228</sup> Th	T : average 14Un01=698.4(0.4) 71Jo14=698.77(0.32) 56Ki16=697.6(0.7)						
* <sup>228</sup> U	D : % $\epsilon$ from 61Ru05<5%						
* <sup>228</sup> Np	D : % $\beta^+$ +SF from 94Kr13=0.020(9) relative to % $\epsilon$ =59(7); % $\alpha$ =40(+8-6)						
* <sup>228</sup> Np	D : % $\beta^+$ =60(+6-8), derived from $\beta^+/\alpha=1.5(4)$ in 94Kr13						
* <sup>228</sup> Pu	T : symmetrized from 03Ni10=1.1(+2.0-0.5)						
<sup>229</sup> At	44890# 400#		1# s >300ns	1/2 <sup>+</sup> #	11 10A124 I	2010	$\beta^-$ ?; $\beta^-$ n ?
<sup>229</sup> Rn	39362 13		11.9 s 1.3	(5/2 <sup>+</sup> )*	09 83Ah03 J	2009	$\beta^-$ =100
<sup>229</sup> Fr	35668 5		50.2 s 0.4	(1/2 <sup>+</sup> )	08 14Bu06 J	1975	$\beta^-$ =100
<sup>229</sup> Ra	32562 15		4.0 m 0.2	5/2 <sup>+</sup> *	08 15Ah04 EJD	1982	$\beta^-$ =100
<sup>229</sup> Ac	30690 12		62.7 m 0.5	3/2 <sup>+</sup>	08 77Th04 J	1952	$\beta^-$ =100
<sup>229</sup> Th	29585.5 2.4		7.916 ky 0.017	5/2 <sup>+</sup> *	08 18Es07 T	1947	$\alpha$ =100
<sup>229</sup> Th <sup>m</sup>	29585.5 2.4	0.0082 0.0001	7 $\mu$ s 1	(3/2 <sup>+</sup> )	08 19Ya18 E	1994	IT=100
<sup>229</sup> Pa	29897 3		1.55 d 0.04	5/2 <sup>+</sup>	08 18Gr09 TD	1949	$\epsilon$ =99.51 5; $\alpha$ =0.49 5
<sup>229</sup> Pa <sup>m</sup>	29909 3	12.20 0.04	420 ns 30	3/2 <sup>-</sup>	08 15Ah04 EJD	1982	IT=100
<sup>229</sup> U	31211 6		57.8 m 0.5	3/2 <sup>+</sup>	08 15Ah04 T	1949	$\beta^+$ $\approx$ 80; $\alpha$ $\approx$ 20
<sup>229</sup> Np	33800 100		4.00 m 0.18	5/2 <sup>+</sup> #	08 04Sa05 TD	1968	$\alpha$ =68 11; $\beta^+$ ?
<sup>229</sup> Np <sup>p</sup>	33960# 110#	160# 50#		5/2 <sup>-</sup> #			
<sup>229</sup> Pu	37390 60		91 s 26	3/2 <sup>+</sup> #	08 10Kh06 TD	1994	$\alpha$ $\approx$ 50 20; $\beta^+$ $\approx$ 50 20;SF<7
<sup>229</sup> Am	42180 110		1.8 s 1.5	5/2 <sup>-</sup> #	15	2015	$\alpha$ $\approx$ 100; $\beta^+$ ?
<sup>229</sup> Am <sup>p</sup>	42440# 230#	260# 200#					
* <sup>229</sup> Rn	T : symmetrized from 09Ne03=12.0(+1.2-1.3)						
* <sup>229</sup> Fr	T : 92Bo05=50.2(0.4); Ensdf2008=50.2(2.0) is misprint						
* <sup>229</sup> Fr	J : strong $\beta^-$ feeding to 1/2+ (142.8 keV) and 1/2- (479.2 keV) in 92Bo05						
* <sup>229</sup> Ra	J : other 18Ly01=5/2						
* <sup>229</sup> Th	T : average 18Es07=7.825(0.087) 14Va04=7.917(0.024) 11Ki16=7.932(0.028)						
* <sup>229</sup> Th	T : 89Go19=7.880(0.060); other 50Ha52=7.340(0.080)						
* <sup>229</sup> Th <sup>m</sup>	T : 17Se01=7(1) us from internal conversion vs time (nickel alloy surface);						
* <sup>229</sup> Th <sup>m</sup>	T : others 19Sh38=10(8) (oxide or hydroxide source) 16We07>60 s						
* <sup>229</sup> Th <sup>m</sup>	T : (for 2+ charge state) 09In01(1 m<T1/2<3 m) 09Ki14<2 h						
* <sup>229</sup> Th <sup>m</sup>	T : 03Mi02 (same group as 09Ki14)=13.9(3.0)h 01Br20(T1/2<6 h or T1/2>20 d)						
* <sup>229</sup> Th <sup>m</sup>	T : 94He08=70(50)h. 19Ve05 (metallic host) excludes 4 us<T1/2<50 us isomer						
* <sup>229</sup> Th <sup>m</sup>	E : rounded from 8.15(0.10) eV, average 20Si22=8.10(0.17) eV						
* <sup>229</sup> Th <sup>m</sup>	E : 20Ge.A=8.09(+0.14-0.19) eV 19Se13=8.28(0.03,stat)(0.16,syst) eV						
* <sup>229</sup> Th <sup>m</sup>	E : 19Ya18=8.30(0.45,stat)(0.81,syst) eV 07Be16=8.1(0.5) eV, recalibrated						
* <sup>229</sup> Th <sup>m</sup>	E : in 19Ye18 from 7.5(0.5) eV; others 16We07 (6.3<E<18.3) eV						
* <sup>229</sup> Th <sup>m</sup>	E : 94He08=3.5(1.0) eV						
* <sup>229</sup> Pa	T : average 87Ah05=1.50(0.05) 18Gr09=1.67(0.08), determined by evaluator						
* <sup>229</sup> Pa	T : as unweighted average (Birge ratio=4.88) from 8 values in 18Gr09						
* <sup>229</sup> Pa	D : % $\alpha$ average 87Ah05=0.48(0.05) 18Gr09=0.53(0.10)						
* <sup>229</sup> Pa <sup>m</sup>	D : from 98Le15						
* <sup>229</sup> Pa <sup>m</sup>	T : from 82Ah08, time-difference between Pa K x rays and 80-400 eV electrons						
* <sup>229</sup> U	J : favored $\alpha$ decay to <sup>225</sup> Th (J=3/2+)						
* <sup>229</sup> Np	T : average 04Sa05=4.0(0.4) 68Ha14=4.0(0.2)						
* <sup>229</sup> Pu	T : average 10Kh06=67(+41-19) 01Ca.B=90(+71-27)						
* <sup>229</sup> Am	T : symmetrized from 15De22=0.9(+2.1-0.7); also 15De22=6.4(+14.9-5.4)						
<sup>230</sup> Rn	42170# 200#		24# s >300ns	0 <sup>+</sup>	12 10A124 I	2010	$\beta^-$ ?
<sup>230</sup> Fr	39487 7		19.1 s 0.5	2 <sup>+</sup> #	12	1987	$\beta^-$ =100
<sup>230</sup> Ra	34516 10		93 m 2	0 <sup>+</sup>	12	1978	$\beta^-$ =100
<sup>230</sup> Ac	33838 16		122 s 3	(1 <sup>+</sup> )	12	1973	$\beta^-$ =100; $\beta^-$ -SF=1.2e-6 4
<sup>230</sup> Th	30862.5 1.2		75.4 ky 0.3	0 <sup>+</sup>	12	1907	IS=0.02 2; $\alpha$ =100;SF<4e-12; 24Ne=5.8e-11 13
<sup>230</sup> Pa	32174 3		17.4 d 0.5	2 <sup>-</sup>	14	1948	$\beta^+$ =92.2 7; $\beta^-$ =7.8 7; $\alpha$ =0.0032 1
<sup>230</sup> U	31615 5		20.23 d 0.02	0 <sup>+</sup>	12 12Po12 T	1948	$\alpha$ =100;22Ne=4.8e-12 20; SF ?
<sup>230</sup> Np	35240 60		4.6 m 0.3	4 <sup>+</sup> #	12	1968	$\beta^+$ <97; $\alpha$ >3

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>230</sup> Pu	36932	14			105 s 10	0 <sup>+</sup>	12 18Mi11 T	1990	$\alpha \approx 100; \beta^+ ?$	
<sup>230</sup> Am	42870#	140#			40 s 9	1 <sup>-</sup> #	12 17Wi13 TD	2003	$\beta^+ \approx 100; \beta^+ SF > 30$	
* <sup>230</sup> Pu	T : average 18Mi11=200(+77-43) 01Ca.B=102(10)									
* <sup>230</sup> Am	T : average 17Wi13=36(+15-8) 16Ka13td=32(+22-9)									
<sup>231</sup> Rn	46550#	300#			2# s >300ns	1/2 <sup>+</sup> #	13 10AI24 I	2010	$\beta^- ?$	
<sup>231</sup> Fr	42081	8			17.6 s 0.6	(1/2 <sup>+</sup> )	13 14Bu06 J	1985	$\beta^- = 100$	
<sup>231</sup> Ra	38216	11			104 s 1	(5/2 <sup>+</sup> )*	13 06Bo33 T	1983	$\beta^- = 100$	
<sup>231</sup> Ra <sup>m</sup>	38282	11	66.21	0.09	~ 53 $\mu$ s	(1/2 <sup>+</sup> )	13	2001	IT=100	
<sup>231</sup> Ac	35763	13			7.5 m 0.1	1/2 <sup>+</sup>	13	1973	$\beta^- = 100$	
<sup>231</sup> Th	33815.8	1.2			25.52 h 0.01	5/2 <sup>+</sup>	13	1911	$\beta^- = 100$	
<sup>231</sup> Pa	33424.3	1.8			32.65 ky 0.20	3/2 <sup>-</sup> *	13 20Je01 T	1918	IS=100; $\alpha$ =100;SF<3e-10; 24Ne=13.4e-10 17;...	
<sup>231</sup> U	33806.0	2.7			4.2 d 0.1	5/2 <sup>+</sup> #	13	1949	$\epsilon \approx 100; \alpha = 0.004$ 1	
<sup>231</sup> Np	35620	50			48.8 m 0.2	5/2 <sup>+</sup> #	13	1950	$\beta^+ = 98$ 1; $\alpha = 2$ 1	
<sup>231</sup> Pu	38309	22			8.6 m 0.5	(3/2 <sup>+</sup> )	13	1999	$\beta^+ ?; \alpha = 13$ 5	
<sup>231</sup> Am	42410#	300#			1# m	5/2 <sup>-</sup> #			$\beta^+ ?; \alpha ?$	
<sup>231</sup> Cm	47270#	300#			20# s	3/2 <sup>+</sup> #			$\beta^+ ?; \alpha ?$	
* <sup>231</sup> Ra	J : 18Ly01=(5/2)									
* <sup>231</sup> Pa	T : average 20Je01=32.57(0.13) 69Ro33=32.765(0.110) 68Br04=32.340(0.115)									
* <sup>231</sup> Pa	T : 61Ki05=32.643(260) 49Va02=34.3(0.3); Birge ratio=3.12									
* <sup>231</sup> Pu	D : % $\alpha$ symmetrized from 99La14=10(+7-3)%; $\beta^+$ not observed directly									
<sup>232</sup> Fr	46073	14			5.5 s 0.6	(5)	06 04Pe17 J	1990	$\beta^- = 100; \beta^- SF ?$	
<sup>232</sup> Ra	40497	9			4.0 m 0.3	0 <sup>+</sup>	06 08Ch.A T	1983	$\beta^- = 100$	
<sup>232</sup> Ac	39154	13			1.98 m 0.08	(1 <sup>+</sup> )	06	1986	$\beta^- = 100$	
<sup>232</sup> Th	35446.7	1.4			14.0 Gy 0.1	0 <sup>+</sup>	06 95Bo18 D	1898	IS=99.98 2; $\alpha$ =100; SF=1.1e-9 4; 24Ne <sup>+</sup> 26Ne<2.78e-10;2 $\beta^- ?$	
<sup>232</sup> Pa	35947	8			1.32 d 0.02	(2 <sup>-</sup> )	06	1949	$\beta^- \approx 100; \epsilon ?$	
<sup>232</sup> U	34609.4	1.8			68.9 y 0.4	0 <sup>+</sup>	06	1949	$\alpha = 100; 24Ne = 8.9e-10$ 7; SF=2.7e-12 6;28Mg<5e-12	
<sup>232</sup> Np	37360#	100#			14.7 m 0.3	(5 <sup>-</sup> )	06	1950	$\beta^+ \approx 100; \alpha ?$	
<sup>232</sup> Pu	38361	17			33.7 m 0.5	0 <sup>+</sup>	06	1973	$\epsilon = ?; \alpha < 20$	
<sup>232</sup> Am	43420#	300#			1.31 m 0.04	1 <sup>-</sup> #	06 90Ha28 D	1967	$\beta^+ \approx 97; \alpha ?; \beta^+ SF = 0.069$ 10	
<sup>232</sup> Cm	46330#	200#			10# s	0 <sup>+</sup>			$\beta^+ ?; \alpha ?$	
* <sup>232</sup> Ra	T : average 08Ch.A=4.00(0.33) 86Gi08=4.2(0.8)									
* <sup>232</sup> Th	D : % <sup>24</sup> Ne+ <sup>26</sup> Ne from 95Bo18; %SF from 00Ho27									
* <sup>232</sup> Np	J : favored $\alpha$ decay from <sup>236</sup> Am (J=5-)									
* <sup>232</sup> Pu	T : average 00La25=33.1(0.8) 73Ja06=34.1(0.7)									
<sup>233</sup> Fr	48920	20			900 ms 100	1/2 <sup>+</sup> #	20	2010	$\beta^- = 100; \beta^- n ?$	
<sup>233</sup> Ra	44334	9			30 s 5	1/2 <sup>+</sup> #	20	1990	$\beta^- = 100$	
<sup>233</sup> Ac	41308	13			143 s 10	(1/2 <sup>+</sup> )	20	1983	$\beta^- = 100$	
<sup>233</sup> Th	38731.6	1.4			21.83 m 0.04	1/2 <sup>+</sup>	20	1935	$\beta^- = 100$	
<sup>233</sup> Th <sup>m</sup>	38737.7	1.4	6.06	0.02	2# s	7/2 <sup>-</sup> #			IT ?; $\beta^- ?$	
<sup>233</sup> Pa	37489.4	1.3			26.975 d 0.013	3/2 <sup>-</sup> *	20	1938	$\beta^- = 100$	
<sup>233</sup> U	36919.1	2.3			159.19 ky 0.15	5/2 <sup>+</sup> *	20	1947	$\alpha = 100; SF < 6e-11$ ; 24Ne=7.2e-11 9; 28Mg<1.3e-13	
<sup>233</sup> Np	37950	50			36.2 m 0.1	5/2 <sup>+</sup> #	20 50Ma14 D	1950	$\beta^+ \approx 100; \alpha \approx 0.0007$	
<sup>233</sup> Np <sup>p</sup>	38000#	60#	50#	30#		(5/2 <sup>-</sup> )				
<sup>233</sup> Pu	40050	50			20.9 m 0.4	5/2 <sup>+</sup> #	20	1957	$\beta^+ \approx 100; \alpha = 0.12$ 5	
<sup>233</sup> Am	43290#	110#			3.2 m 0.8	5/2 <sup>-</sup> #	20 00Sa52 TD	2000	$\beta^+ ?; \alpha = 4.5$ 9	
<sup>233</sup> Cm	47290	80			27 s 10	3/2 <sup>+</sup> #	20 10Kh06 TD	2001	$\alpha = 20$ 10; $\beta^+ = 80$ 10	
<sup>233</sup> Bk	52770#	230#			40 s 30	3/2 <sup>-</sup> #	20 15De22 TD	2015	$\alpha \approx 82; \beta^+ ?$	
* <sup>233</sup> Th <sup>m</sup>	J : from expected conf=n7/2[743]									
* <sup>233</sup> Th <sup>m</sup>	T : from B(E3:7/2- ->1/2+)( <sup>233</sup> Th)=B(E3: 1/2+ -> 7/2-)( <sup>235</sup> U)/4 with									
* <sup>233</sup> Th <sup>m</sup>	T : T1/2( <sup>235</sup> U)=25.7(0.1) m and $\alpha_T$ ( <sup>235</sup> U)=2.79(0.05)e20									
* <sup>233</sup> Th <sup>m</sup>	T : and $\alpha_T$ ( <sup>233</sup> Th)=3.36(0.09)e9									



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>233</sup> Np	D : % $\alpha$ observed in 50Ma14 with $\beta^+/\alpha=1.5e5$								**
* <sup>233</sup> Am	D : % $\alpha$ combining 10Kh06<6 and 00Sa52>3								**
* <sup>233</sup> Cm	T : symmetrized from 10Kh06=23(+13-6)								**
* <sup>233</sup> Bk	T : symmetrized from 15De22=21(+48-17)								**
<sup>234</sup> Ra	46931	8			30 s 10	0 <sup>+</sup>	07	1990	$\beta^- = 100; \beta^- \text{ SF ?}$
<sup>234</sup> Ac	44841	14			45 s 2	1 <sup>+</sup> #	07 08Ch.A T	1986	$\beta^- = 100$
<sup>234</sup> Ac <sup>m</sup>	44980	30	140	30	> 93 s		08Ch.A TE	2008	$\beta^- ?; IT ?$
<sup>234</sup> Ac <sup>n</sup>	45460	30	620	30	180 s 70		08Ch.A TE	2008	$\beta^- ?; IT ?$
<sup>234</sup> Th	40613.0	2.6			24.107 d 0.024	0 <sup>+</sup>	07 18Pa45 T	1900	$\beta^- = 100; \alpha ?$
<sup>234</sup> Pa	40339	4			6.70 h 0.05	4 <sup>+</sup>	07	1913	$\beta^- = 100$
<sup>234</sup> Pa <sup>m</sup>	40417.9	2.8	79	3	IT	(0 <sup>-</sup> )	07 73Go40 E	1951	$\beta^+ \approx 100; IT=0.16 4$
<sup>234</sup> U	38145.0	1.1			245.5 ky 0.6	0 <sup>+</sup>	07	1912	IS=0.0054 5; $\alpha=100$ ; SF=1.64e-9 22; 28Mg=1.4e-11 3; 24Ne <sup>+</sup> 26Ne=9e-12 7
<sup>234</sup> U <sup>m</sup>	39566.3	1.1	1421.257	0.017	33.5 $\mu$ s 2.0	6 <sup>-</sup>	07	1963	IT=100
<sup>234</sup> Np	39955	8			4.4 d 0.1	(0 <sup>+</sup> )	07	1949	$\beta^+ = 100$
<sup>234</sup> Pu	40350	7			8.8 h 0.1	0 <sup>+</sup>	07	1949	$\epsilon \approx 94; \alpha \approx 6$
<sup>234</sup> Am	44460#	160#			2.32 m 0.08	0 <sup>-</sup> #	07 90Ha02 D	1967	$\beta^+ \approx 100; \alpha=0.039 12$ ; $\beta^+ \text{ SF}=0.0066 18$
<sup>234</sup> Cm	46722	17			52 s 9	0 <sup>+</sup>	07 10Kh06 D	2001	$\beta^+ \approx 71; \alpha \approx 27; \text{SF} \approx 2$
<sup>234</sup> Bk	53400#	150#			20 s 5	3 <sup>-</sup> #	07 16Ka13 T	2003	$\alpha > 80; \beta^+ < 20$
* <sup>234</sup> Ac	I : 08Ch.A reports two isomers with T1/2>93 s and T1/2=149(+95-42) s								**
* <sup>234</sup> Ac <sup>n</sup>	T : symmetrized from 08Ch.A=145(+95-42)								**
* <sup>234</sup> Th	T : average 18Pa45=24.157(0.073) 48Kn23=24.101(0.025) 39Sa11=24.1(0.2)								**
* <sup>234</sup> Pa <sup>m</sup>	E : from 73Go40<10 keV above (3+) level at 73.92(0.02)								**
* <sup>234</sup> Am	T : also 04Sa05=3.5(1.3), not used								**
* <sup>234</sup> Cm	T : average 16Ka13=49(+15-9) 01Ca.B=51(12)								**
* <sup>234</sup> Bk	T : symmetrized from 16Ka13=19(+6-4)								**
<sup>235</sup> Ra	51130#	300#			5# s	5/2 <sup>+</sup> #			$\beta^- ?$
<sup>235</sup> Ac	47357	14			62 s 4	1/2 <sup>+</sup> #	14 08Ch.A T	2006	$\beta^- = 100$
<sup>235</sup> Th	44018	13			7.2 m 0.1	1/2 <sup>+</sup> #	14	1969	$\beta^- = 100$
<sup>235</sup> Pa	42289	14			24.4 m 0.2	3/2 <sup>-</sup>	14 77Th04 J	1950	$\beta^- = 100$
<sup>235</sup> U	40918.8	1.1			704 My1	7/2 <sup>-</sup> *	14	1935	IS=0.7204 6; $\alpha=100$ ; SF=7e-9 2; 20Ne=8e-10 4; 25Ne $\approx$ 8e-10; 28Mg=8e-10
<sup>235</sup> U <sup>m</sup>	40918.9	1.1	0.0767	0.0001	25.7 m 0.1	1/2 <sup>+</sup>	14 18Po07 E	1966	IT=100
<sup>235</sup> U <sup>n</sup>	43420	300	2500	300	3.6 ms 1.8		14	2007	SF $\approx$ 100; IT ?
<sup>235</sup> Np	41043.0	1.4			396.1 d 1.2	5/2 <sup>+</sup>	14	1949	$\epsilon=99.99740 13$ ;; $\alpha=0.00260 13$
<sup>235</sup> Pu	42182	21			25.3 m 0.5	(5/2 <sup>+</sup> )	14	1957	$\beta^+ = 99.9972 7; \alpha=0.0028 7$
<sup>235</sup> Am	44620	50			10.3 m 0.6	5/2 <sup>-</sup> #	14	1996	$\beta^+ = 99.60 5; \alpha=0.40 5$
<sup>235</sup> Cm	48010#	100#			7 m 3	5/2 <sup>+</sup> #	14 20Kh10 TD	1981	$\beta^+ ?; \alpha=4 3$
<sup>235</sup> Bk	52770#	400#			1# m	3/2 <sup>-</sup> #			$\beta^+ ?; \alpha ?$
* <sup>235</sup> U <sup>m</sup>	E : rounded from 18Po07=0.076737 (0.000018) keV								**
* <sup>235</sup> U <sup>n</sup>	T : from 16Ch11; value depends on the chemical environment								**
* <sup>235</sup> Cm	T : symmetrized from 20Kh10=300(+250-100)s								**
* <sup>235</sup> Cm	D : % $\alpha$ determined from 0<% $\alpha$ <8 in 20Kh10								**
<sup>236</sup> Ac	51220	40			4.5 m 3.6	3 <sup>+</sup> #	15 10Ch19 T	2010	$\beta^- = 100$
<sup>236</sup> Th	46255	14			37.3 m 1.5	0 <sup>+</sup>	15	1973	$\beta^- = 100$
<sup>236</sup> Pa	45334	14			9.1 m 0.1	1 <sup>(-)</sup>	06	1963	$\beta^- = 100; \beta^- \text{ SF}=6e-8 4$
<sup>236</sup> U	42444.6	1.1			23.42 My0.04	0 <sup>+</sup>	06	1951	$\alpha=100; \text{SF}=9.4e-8 4$
<sup>236</sup> U <sup>m</sup>	43497.1	1.3	1052.5	0.6	100 ns 4	4 <sup>-</sup>	06	1973	IT=100
<sup>236</sup> U <sup>n</sup>	45195	3	2750	3	120 ns 2	(0 <sup>+</sup> )	06	1969	IT=87 6; SF=13 6; $\alpha ?$
<sup>236</sup> Np	43380	50			153 ky 5	(6 <sup>-</sup> )	06	1949	$\epsilon=86.3 8; \beta^- = 13.5 8$ ; $\alpha=0.16 4$
<sup>236</sup> Np <sup>m</sup>	43438	7	60	50	IT*	(1 <sup>-</sup> )	06	1949	$\epsilon=50 3; \beta^- = 50 3$

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
<sup>236</sup> Np <sup>p</sup>	43616	14	240	50	AD		(3 <sup>-</sup> )	06		
<sup>236</sup> Pu	42901.5	1.8				2.858 y 0.008	0 <sup>+</sup>	06 90Og01 D	1949	$\alpha=100$ ;SF=1.9e-7 4; 28Mg=2e-12;2 $\beta^+$ ?
<sup>236</sup> Pu <sup>m</sup>	44087.0	1.8	1185.45	0.15		1.2 $\mu$ s 0.3	5 <sup>-</sup>	06	2005	IT=100
<sup>236</sup> Am	46040#	120#				3.6 m 0.1	5 <sup>-</sup>	06 04Sa05 D	1998	$\beta^+ \approx 100$ ; $\alpha=4.0e-3$ 1
<sup>236</sup> Am <sup>m</sup>	46090#	130#	50#	50#		2.9 m 0.2	(1 <sup>-</sup> )	06	2004	$\beta^+ \approx 100$ ; $\alpha$ ?
<sup>236</sup> Cm	47853	18				6.8 m 0.8	0 <sup>+</sup>	06 10Kh06 TD	2010	$\beta^+=82$ 2; $\alpha=18$ 2; SF ?
<sup>236</sup> Bk	53540#	360#				26 s 10	4 <sup>+</sup> #	16 17Ko02 TD	2017	$\beta^+ \approx 100$ ; $\alpha$ ?; $\beta^+$ SF=0.04 2
* <sup>236</sup> Ac	T : symmetrized from 10Ch19=72(+345-33) s									
* <sup>236</sup> Pa	D : $\beta^-$ SF decay questioned in 90Ha02									
* <sup>236</sup> Bk	T : symmetrized from 17Ko02=22(+13-6); other 20Po07~19									
<sup>237</sup> Ac	54020#	400#				23# s	1/2 <sup>+</sup> #			$\beta^-$ ?
<sup>237</sup> Th	49955	16				4.8 m 0.5	5/2 <sup>+</sup> #	06	1993	$\beta^- =100$
<sup>237</sup> Pa	47528	13				8.7 m 0.2	1/2 <sup>+</sup>	06 77Th04 J	1954	$\beta^- =100$
<sup>237</sup> U	45390.1	1.2				6.752 d 0.002	1/2 <sup>+</sup>	06	1940	$\beta^- =100$
<sup>237</sup> U <sup>m</sup>	45664.1	1.6	274.0	1.0		155 ns 6	7/2 <sup>-</sup>	06	1968	IT=100
<sup>237</sup> Np	44871.6	1.1				2.144 My0.007	5/2 <sup>+</sup> *	06 89Pr.A D	1948	$\alpha=100$ ;SF<2e-10; 30Mg<4e-12
<sup>237</sup> Np <sup>m</sup>	45816.8	1.1	945.20	0.10		710 ns 40	13/2 <sup>-</sup>	06 90St29 JED	1990	IT=100
<sup>237</sup> Pu	45091.7	1.7				45.64 d 0.04	7/2 <sup>-</sup>	06	1949	$\epsilon=99.9958$ 4; $\alpha=0.0042$ 4
<sup>237</sup> Pu <sup>m</sup>	45237.2	1.7	145.543	0.008		180 ms 20	1/2 <sup>+</sup>	06	1972	IT=100
<sup>237</sup> Pu <sup>m</sup>	47990	250	2900	250		1.1 $\mu$ s 0.1		06	1970	SF $\approx$ 100; IT ?
<sup>237</sup> Am	46570#	60#				73.6 m 0.8	5/2 <sup>-</sup>	06	1970	$\beta^+=99.975$ 3; $\alpha=0.025$ 3
<sup>237</sup> Cm	49250	70				> 10# m	5/2 <sup>+</sup> #	06 06As03 DT	2002	$\beta^+$ ?; $\alpha$ =?
<sup>237</sup> Cm <sup>p</sup>	49450#	170#	200#	150#			7/2 <sup>-</sup> #	20Kh10 E		
<sup>237</sup> Bk	53210#	230#				2# m	3/2 <sup>-</sup> #			$\beta^+$ ?; $\alpha$ ?
<sup>237</sup> Cf	57940	100				0.8 s 0.2	5/2 <sup>+</sup> #	06 10Kh06 TD	1995	$\alpha=70$ 10;SF=30 10; $\beta^+$ ?
* <sup>237</sup> U <sup>m</sup>	J : E1 to 5/2+									
* <sup>237</sup> Np	D : also cluster (Z=10-14) emission 92Mo03<1.8e-12%									
* <sup>237</sup> Np <sup>m</sup>	J : multiple decay branches in 90St29 agree with J=11/2,13/2-, but the									
* <sup>237</sup> Np <sup>m</sup>	J : absence of gamma rays to J=7/2- and 9/2+ argues against J=11/2									
* <sup>237</sup> Cm	T : partial T1/2( $\alpha$ ) 06As03=1100 m and by assuming % $\alpha$ =1									
* <sup>237</sup> Cm <sup>p</sup>	E : 50(1) keV E1 gamma above the <sup>237</sup> Cm gs in 20Kh10									
* <sup>237</sup> Cf	T : other (not used) 95La09=2.1(0.3)									
<sup>238</sup> Th	52530#	280#				9.4 m 2.0	0 <sup>+</sup>	15	1999	$\beta^- =100$
<sup>238</sup> Pa	50894	16				2.28 m 0.09	3 <sup>-</sup> #	15 85Ba57 D	1968	$\beta^- =100$ ; $\beta^-$ SF<2.6e-6
<sup>238</sup> U	47307.7	1.5				4.463 Gy 0.003	0 <sup>+</sup>	15 18Pa45 T	1896	IS=99.2742 10; $\alpha=100$ ; SF=5.44e-5 7; 2 $\beta^- =2.2e-10$ 3
<sup>238</sup> U <sup>m</sup>	49865.6	1.6	2557.9	0.5		280 ns 6	0 <sup>+</sup>	15	1979	IT=97.4 4;SF=2.6 4
<sup>238</sup> Np	47454.6	1.1				2.099 d 0.002	2 <sup>+</sup> *	15	1949	$\beta^- =100$
<sup>238</sup> Np <sup>m</sup>	49760#	200#	2300#	200#		112 ns 39		15	1970	SF $\approx$ 100;IT ?
<sup>238</sup> Pu	46163.1	1.1				87.7 y 0.1	0 <sup>+</sup>	15 89Wa10 D	1949	$\alpha=100$ ;SF=1.9e-7 1; 32Si $\approx$ 1.4e-14; 28Mg+30Mg $\approx$ 6e-15
<sup>238</sup> Am	48420	60				98 m 3	1 <sup>+</sup>	15 72Ah04 TD	1950	$\beta^+=100$ ; $\alpha=1.0e-4$ 4
<sup>238</sup> Am <sup>m</sup>	50920#	210#	2500#	200#		35 $\mu$ s 18		15	1967	SF $\approx$ 100;IT ?
<sup>238</sup> Cm	49445	12				2.2 h 0.4	0 <sup>+</sup>	15	1994	$\epsilon$ ?; $\alpha=3.84$ 18; SF=0.048 2
<sup>238</sup> Bk	54220#	260#				2.40 m 0.08	1#	15 94Kr03 TD	1994	$\beta^+ \approx 100$ ; $\alpha$ ?; $\beta^+$ SF=0.048 2
<sup>238</sup> Cf	57280#	300#				21.1 ms 1.3	0 <sup>+</sup>	15 10Kh06 D	1995	SF=97.5 14; $\alpha=2.5$ 14
* <sup>238</sup> U	T : average 18Pa45=4.456(0.021),									
* <sup>238</sup> U	T : 4.468(0.005), adjusted in 04Sc03 from 71Ja07=4.4683(0.0024),									
* <sup>238</sup> U	T : 4.457(0.004), adjusted in 04Sc03 from 59St45=4.460(0.005),									
* <sup>238</sup> U	T : 4.51(0.02), adjusted in 04Sc03 from 55Ko13=4.507(0.009),									
* <sup>238</sup> U	T : 4.495(0.018), adjusted in 04Sc03 from 49Ki26=4.490(0.005).									
* <sup>238</sup> U	D : %2 $\beta^- =2.2(3)e-10$ %, derived from T1/2(2v- $\beta$ $\beta$ )=2.0(0.6) Zy in 91Tu02;									
* <sup>238</sup> U	D : %SF=5.44(0.07)e-5%, derived from T1/2(SF)=8.2(0.1) Py in 00Ho27									
* <sup>238</sup> Cf	D : % $\alpha$ from 10Kh06<5%									

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
<sup>239</sup> Th	56500#	400#			1# m	7/2 <sup>+</sup> #			$\beta^-$ ?
<sup>239</sup> Pa	53340#	200#			1.8 h 0.5	1/2 <sup>+</sup> #	14	1995	$\beta^-$ =100
<sup>239</sup> U	50572.7	1.5			23.45 m 0.02	5/2 <sup>+</sup>	14	1937	$\beta^-$ =100
<sup>239</sup> U <sup>m</sup>	50706.5	1.5	133.7991	0.0010	780 ns 40	1/2 <sup>+</sup>	14	1975	IT=100
<sup>239</sup> U <sup>n</sup>	53070#	900#	2500#	900#	> 250 ns		14	94Ob0	IT 1994
<sup>239</sup> Np	49311.0	1.3			2.356 d 0.003	5/2 <sup>+</sup> *	14	1940	$\beta^-$ =100; $\alpha$ ?
<sup>239</sup> Pu	48588.2	1.1			24.11 ky 0.03	1/2 <sup>+</sup> *	14	1946	$\alpha$ =100;SF=3.1e-10 6
<sup>239</sup> Pu <sup>m</sup>	48979.8	1.1	391.584	0.003	193 ns 4	7/2 <sup>-</sup>	14	1955	IT=100
<sup>239</sup> Pu <sup>n</sup>	51690	200	3100	200	7.5 $\mu$ s 1.0	(5/2 <sup>+</sup> )	14	1970	SF $\approx$ 100;IT ?
<sup>239</sup> Am	49390.4	2.0			11.9 h 0.1	5/2 <sup>-</sup>	14	1949	$\epsilon$ =99.990 1; $\alpha$ =0.010 1
<sup>239</sup> Am <sup>m</sup>	51890	200	2500	200	163 ns 12	(7/2 <sup>+</sup> )	14	1969	SF $\approx$ 100;IT ?
<sup>239</sup> Cm	51150	150			2.5 h 0.4	7/2 <sup>-</sup> #	14	02Sh.C	TD 1952
<sup>239</sup> Cm <sup>p</sup>	51390#	180#	240#	100#	> 100# ns	1/2 <sup>+</sup>			IT ?; $\beta^+$
<sup>239</sup> Bk	54250#	210#		*	100# s	(7/2 <sup>+</sup> )	14	10An08	TD 1989
<sup>239</sup> Bk <sup>p</sup>	54290#	210#	41	11	> 30# $\mu$ s	(3/2 <sup>-</sup> )	89Ha27	J 1989	IT ?; $\beta^+$ ?
<sup>239</sup> Cf	58200#	120#			28 s 2	5/2 <sup>+</sup> #	14	20Kh10	TD 1981
<sup>239</sup> Es	63630#	300#			1# s	3/2 <sup>-</sup> #			$\alpha$ ?; $\beta^+$ ?;SF ?
* <sup>239</sup> U <sup>n</sup>	T : other 94Ob01<0.3 ns is less likely								
* <sup>239</sup> Am	J : favored $\alpha$ decay to J=5/2 <sup>-</sup> level at 49.10 keV in <sup>235</sup> Np								
* <sup>239</sup> Cm <sup>p</sup>	E : 146 keV in <sup>237</sup> Pu, N=143 isotope								
* <sup>239</sup> Bk	J : from 89Ha27								
* <sup>239</sup> Cf	T : other 81Mu12=39(+37-12)								
<sup>240</sup> Pa	57010#	200#			20# s	3 <sup>+</sup> #			$\beta^-$ ?
<sup>240</sup> U	52715.5	2.6			14.1 h 0.1	0 <sup>+</sup>	08	1953	$\beta^-$ =100; $\alpha$ ?
<sup>240</sup> Np	52316	17		*	61.9 m 0.2	(5 <sup>+</sup> )	08	1953	$\beta^-$ =100
<sup>240</sup> Np <sup>m</sup>	52334	13	18	14	IT*	(1 <sup>+</sup> )	08	81Hs02	E 1948
<sup>240</sup> Pu	50125.3	1.1			6.561 ky 0.007	0 <sup>+</sup>	08	18Be29	D 1949
<sup>240</sup> Pu <sup>m</sup>	51434.0	1.1	1308.74	0.05	165 ns 10	5 <sup>-</sup>	08	1967	IT=100
<sup>240</sup> Am	51510	14			50.8 h 0.3	(3 <sup>-</sup> )	08	1949	$\beta^+$ =100; $\alpha$ $\approx$ 1.9e-4 7
<sup>240</sup> Am <sup>m</sup>	54510	200	3000	200	940 $\mu$ s 40		08	1967	SF $\approx$ 100;IT ?
<sup>240</sup> Cm	51724.2	1.9			30.4 d 3.7	0 <sup>+</sup>	08	08Qi03	T 1949
<sup>240</sup> Bk	55660#	150#			4.8 m 0.8	7 <sup>-</sup> #	08	83Ga05	D 1980
<sup>240</sup> Bk <sup>p</sup>	55900#	180#	240#	100#		am			$\beta^+$ ?; $\alpha$ ?; $\beta^+$ SF=0.0020 13
<sup>240</sup> Cf	57989	18			40.3 s 0.9	0 <sup>+</sup>	08	10As.A	T 1970
<sup>240</sup> Es	64230#	370#			6.0 s 1.7	4 <sup>-</sup> #		17Ko02	TD 2017
* <sup>240</sup> Pu	D : also %SF=5.632(0.062)e-6 from T1/2(SF)=116.5(1.3) Gy in 13Sa65								
* <sup>240</sup> Pu <sup>m</sup>	J : M1 to 4- and 6-								
* <sup>240</sup> Cm	T : from 08Qi03; other Ensdf2009=27(1), based on 49Se01=26.8 and 67Ba42=28								
* <sup>240</sup> Cm	T : values that are reported without uncertainties								
* <sup>240</sup> Bk	D : % $\beta^+$ SF symmetrized from 83Ga05=0.0013(+18-7)%								
* <sup>240</sup> Cf	D : $\alpha$ , %SF from 10Kh06; other $\alpha$ $\sim$ 9, %SF $\sim$ 2 in 95La09								
* <sup>240</sup> Es	T : average 20Po07=4.7(+3.8-1.4) 17Ko02=6(2); other 20Kh08=8(+6-2)								
<sup>241</sup> Pa	59740#	300#			28# m	1/2 <sup>+</sup> #			$\beta^-$ ?
<sup>241</sup> U	56200#	200#			4# m	7/2 <sup>+</sup> #	15		$\beta^-$ ?
<sup>241</sup> Np	54320	100			13.9 m 0.2	(5/2 <sup>+</sup> )	15	1959	$\beta^-$ =100; $\alpha$ ?
<sup>241</sup> Pu	52955.1	1.1			14.329 y 0.029	5/2 <sup>+</sup> *	15	1949	$\beta^-$ $\approx$ 100; $\alpha$ =0.00245 8;
<sup>241</sup> Pu <sup>m</sup>	53116.8	1.1	161.6853	0.0009	880 ns 50	1/2 <sup>+</sup>	15	1975	SF<2.4e-14
<sup>241</sup> Pu <sup>n</sup>	55160	200	2200	200	20.5 $\mu$ s 2.2		15	1970	IT=100
<sup>241</sup> Am	52934.3	1.1			432.6 y 0.6	5/2 <sup>-</sup> *	15	1949	$\alpha$ =100;SF=3.6e-10 9
<sup>241</sup> Am <sup>m</sup>	55130	200	2200	200	1.2 $\mu$ s 0.3		15	71Br39	E 1969
<sup>241</sup> Cm	53701.8	1.6			32.8 d 0.2	1/2 <sup>+</sup>	15	1952	$\epsilon$ =99.0 1; $\alpha$ =1.0 1
<sup>241</sup> Bk	55980#	170#			4.6 m 0.4	(7/2 <sup>+</sup> )	15	2003	$\beta^+$ =?; $\alpha$ ?
<sup>241</sup> Bk <sup>p</sup>	56030#	170#	51	3	AD	(3/2 <sup>-</sup> )	15		IT ?
<sup>241</sup> Cf	59330#	170#			> 25# $\mu$ s		15	20Kh10	D 1970
					2.35 m 0.18	7/2 <sup>-</sup> #	15	20Kh10	D 1970
									$\beta^+$ ?; $\alpha$ =15 1

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>241</sup> Cf <sup>p</sup>	59480#	200#	150#	100#	Nm		1/2 <sup>+</sup> #	15			
<sup>241</sup> Es	63890#	230#				5.1 s 0.8	3/2 <sup>-</sup> #	15 20Kh08	TD 1996	$\alpha \approx 100; \beta^+ ?$	*
<sup>241</sup> Es <sup>p</sup>	64120#	250#	230#	100#			<i>am</i>				
<sup>241</sup> Fm	69220#	300#				730 $\mu$ s 60	5/2 <sup>+</sup> #	15 08Kh10	TD 2008	SF=?; $\alpha < 14; \beta^+ < 12$	
* <sup>241</sup> Pu	D : % $\alpha$ from $\beta^-/\alpha=2.45(0.08)e-5$ in 68Ah01										**
* <sup>241</sup> Cf	T : from 10As.A=141(11) s; other 70Si19=3.78(0.70) m										**
* <sup>241</sup> Es	T : symmetrized from 20Kh08=4.3(+2.4-1.2); other 96Ni09=8(+6-4)										**
<sup>242</sup> U	58620#	200#				16.8 m 0.5	0 <sup>+</sup>	02	1979	$\beta^- = 100$	
<sup>242</sup> Np	57420	200			*	2.2 m 0.2	(1 <sup>+</sup> )	02	1979	$\beta^- = 100$	
<sup>242</sup> Np <sup>m</sup>	57470#	210#	50#	50#	*	5.5 m 0.1	(6 <sup>+</sup> )	02	1981	$\beta^- = 100$	
<sup>242</sup> Pu	54716.9	1.2				375 ky 2	0 <sup>+</sup>	02 18Be29	D 1950	$\alpha=100; SF=5.510e-4 41$	*
<sup>242</sup> Am	55468.0	1.1				16.02 h 0.02	1 <sup>-</sup> *	02	1949	$\beta^- = 82.7 3; \epsilon = 17.3 3$	
<sup>242</sup> Am <sup>m</sup>	55516.6	1.1	48.60	0.05		141 y 2	5 <sup>-</sup>	02	1950	IT=99.55 2; $\alpha=0.45 2$ ; SF<4.7e-9	
<sup>242</sup> Am <sup>n</sup>	57670	80	2200	80		14.0 ms 1.0	(2 <sup>+</sup> , 3 <sup>-</sup> )	02	1962	SF $\approx$ 100; IT=?	
<sup>242</sup> Cm	54803.7	1.1				162.8 d 0.2	0 <sup>+</sup>	02	1949	$\alpha=100; SF=6.2e-6 3$ ; 34Si=1.1e-14 4 2; $\beta^+ ?$	*
<sup>242</sup> Cm <sup>m</sup>	57600	100	2800	100		180 ns 70		02	1971	SF ?; IT ?	
<sup>242</sup> Bk	57750#	140#				7.0 m 1.3	3 <sup>+</sup> #	02 80Ga07	D 1972	$\beta^+ \approx 100; \beta^+ SF < 3e-5; \alpha ?$	
<sup>242</sup> Bk <sup>m</sup>	59750#	240#	2000#	200#		600 ns 100		02	1972	SF $\approx$ 100; IT ?	
<sup>242</sup> Bk <sup>p</sup>	57900	90	150#	100#			4 <sup>-</sup>				
<sup>242</sup> Cf	59387	13				3.49 m 0.15	0 <sup>+</sup>	02 70Si19	T 1967	$\alpha=61 3; \beta^+=39 3; SF < 0.014$	*
<sup>242</sup> Es	64800#	260#				17.8 s 1.6	2 <sup>+</sup> #	02 10An08	TD 1994	$\alpha=57 3; \beta^+=43 3$ ; $\beta^+ SF=0.6 2$	*
<sup>242</sup> Fm	68400#	400#				800 $\mu$ s 200	0 <sup>+</sup>	02	1975	SF $\approx$ 100; $\alpha ?$	*
* <sup>242</sup> Pu	D : %SF other 13Sa65=5.564(0.072)e-4 from T1/2(SF) 13Sa65=67.4(0.9) Gy										**
* <sup>242</sup> Cm	D : % <sup>34</sup> Si symmetrized from 1.0(+4-3)e-14										**
* <sup>242</sup> Cf	T : average 70Si19=3.68(0.44) 67Si07=3.4(0.2) 67Fi04=3.2(0.5)										**
* <sup>242</sup> Cf	T : 67I101=3.7(0.3)										**
* <sup>242</sup> Cf	D : % $\alpha$ from 11Ve03; other 81Mu12=80(20)										**
* <sup>242</sup> Es	T : others 00Sh10=11(3) 96Ni09=16(+6-4)										**
* <sup>242</sup> Es	D : % $\beta^+ SF$ from 00Sh10; other 10An08=1.3(+1.2-0.7)										**
* <sup>242</sup> Fm	T : 08Kh10 excludes 4 us-1s (conflicting)										**
<sup>243</sup> U	62480#	300#				16# m	9/2 <sup>-</sup> #			$\beta^- ?$	
<sup>243</sup> Np	59810#	30#				1.85 m 0.15	5/2 <sup>+</sup> #	14	1979	$\beta^- = 100$	
<sup>243</sup> Np <sup>p</sup>	59926	10	120#	30#	Nm		5/2 <sup>-</sup> #				
<sup>243</sup> Pu	57754.6	2.5				4.9553 h 0.0025	7/2 <sup>+</sup>	14 19Le09	T 1951	$\beta^- = 100$	*
<sup>243</sup> Pu <sup>m</sup>	58138.2	2.5	383.64	0.25		330 ns 30	(1/2 <sup>+</sup> )	14	1975	IT=100	
<sup>243</sup> Am	57175.0	1.4				7.350 ky 0.009	5/2 <sup>-</sup> *	14 20Ma.A	T 1950	$\alpha=100; SF=3.7e-9 9$	*
<sup>243</sup> Am <sup>m</sup>	59480	200	2300	200		5.5 $\mu$ s 0.5		14	1970	SF $\approx$ 100; IT ?	
<sup>243</sup> Cm	57181.9	1.5				29.1 y 0.1	5/2 <sup>+</sup> *	14	1950	$\alpha \approx 100; \epsilon = 0.29 3$ ; SF=5.3e-9 9	
<sup>243</sup> Cm <sup>m</sup>	57269.3	1.5	87.4	0.1		1.08 $\mu$ s 0.03	1/2 <sup>+</sup>	14	1971	IT=100	
<sup>243</sup> Cm <sup>p</sup>	57285	15	103	15	AD		(7/2 <sup>+</sup> )	14	1984	IT ?	
<sup>243</sup> Bk	58690	5			*	4.6 h 0.2	3/2 <sup>-</sup>	14 18Ah01	J 1950	$\beta^+ \approx 100; \alpha \approx 0.15$	
<sup>243</sup> Bk <sup>p</sup>	58710	19	20	20	AD*	> 30# $\mu$ s	(7/2 <sup>+</sup> )			IT $\approx$ 100; $\beta^+ ?$	
<sup>243</sup> Cf	60990#	180#				10.8 m 0.3	(1/2 <sup>+</sup> )	14 18Ko05	T 1967	$\beta^+ \approx 86 3; \alpha \approx 14 3$	*
<sup>243</sup> Es	64750#	210#			*	22.1 s 1.4	(7/2 <sup>+</sup> )	14 10An08	JTD 1973	$\alpha=61 6; \beta^+ ?; SF < 1$	*
<sup>243</sup> Es <sup>m</sup>	64800#	220#	50#	50#	*	> 50# $\mu$ s	3/2 <sup>-</sup> #	10An08	I	IT ?; $\alpha ?; \beta^+ ?$	
<sup>243</sup> Fm	69320#	130#				231 ms 9	7/2 <sup>-</sup> #	14 20Kh10	D 1981	$\alpha=91 3; SF=9 3; \beta^+ ?$	*
* <sup>243</sup> Pu	T : average 19Le09=4.948(10) 69Ho10=4.958(5) 68Di09=4.955(3)										**
* <sup>243</sup> Am	T : average 20Ma.A=7342(14), 7345(14) y 07Ag02=7364(22) y, deduced from										**
* <sup>243</sup> Am	T : T1/2( <sup>243</sup> Am)=7357(23) y and T1/2( <sup>241</sup> Am)=432.6(0.6) y,										**
* <sup>243</sup> Am	T : 74Po17=7380(34) y 68Br22=7336.9(57.2), 7390(50) y										**
* <sup>243</sup> Cf	T : average 18Ko05=10.9(0.5) 67Fi04=12.5(1.0) 67Si08=10.3(0.5)										**
* <sup>243</sup> Cf	D : % $\alpha$ , % $\beta^+$ from I(7060 $\alpha$ )/I(7171 $\alpha$ ) $\approx$ 2.5 in										**
* <sup>243</sup> Cf	D : 67Fi04 and ( $\beta^+ + \alpha$ )/I(7060 $\alpha$ )=10(2) in 67Si08										**
* <sup>243</sup> Es	T : average 19Br06=24(3) 10An08=23(3) 89Ha27=21(5) 73Es02=21(2)										**
* <sup>243</sup> Es	J : from 10An08										**



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
$^{246}\text{Pu}$	65395	15				10.84 d 0.02	$0^+$	11	1955	$\beta^- = 100$
$^{246}\text{Am}$	64994#	18#				39 m 3	$7^-$	11	1955	$\beta^- = 100$ *
$^{246}\text{Am}^m$	65024	15	30#	10#		25.0 m 0.2	$2(-)$	11	1955	$\beta^- \approx 100; \text{IT} ?$ *
$^{246}\text{Am}^n$	66990#	800#	2000#	800#		73 $\mu\text{s}$ 10		11	1972	SF $\approx 100; \text{IT} ?$
$^{246}\text{Cm}$	62616.9	1.5				4.706 ky 0.040	$0^+$	11	1954	$\alpha = 99.97385 ?$ ; SF=0.02615 7
$^{246}\text{Cm}^m$	63796.6	1.5	1179.66	0.13		1.12 s 0.24	$8^-$	11 19Sh34	ETJ 2012	IT=100
$^{246}\text{Bk}$	63970	60				1.80 d 0.02	$2(-)$	11	1954	$\beta^+ \approx 100; \alpha ?$
$^{246}\text{Cf}$	64090.2	1.5				35.7 h 0.5	$0^+$	11	1951	$\alpha = 100; \text{SF} = 2.4e-4 4; \epsilon ?$
$^{246}\text{Es}$	67820	90				7.5 m 0.5	$4^- \#$	11	1954	$\beta^+ = 90.1 18; \alpha = 9.9 18;$ $\beta^+ \text{SF} \approx 0.003$
$^{246}\text{Es}^p$	68010	100	190	50			$2^- \#$	08An16	E	*
$^{246}\text{Es}^q$	68200	90	379.5	2.0				19Vo03	E	*
$^{246}\text{Fm}$	70191	14				1.54 s 0.04	$0^+$	11 11Ve03	TD 1966	$\alpha = 93.2 6; \text{SF} = 6.8 6; \epsilon < 1.3$ *
$^{246}\text{Md}$	76120#	260#			*	0.92 s 0.18	$1^- \#$	11 10An08	TD 1996	$\alpha = 100$ *
$^{246}\text{Md}^m$	76170#	260#	60	60	AD*	4.4 s 0.8	$4^- \#$	11 10An08	TD 2010	$\alpha = 57 3; \beta^+ > 77; \beta^+ \text{SF} > 10;$ $\alpha < 23$
$^{*246}\text{Am}$	J : direct $\beta^-$ feeding to $^{246}\text{Cm}^m$ (K=8-)									**
$^{*246}\text{Am}^m$	D : other %IT<0.02 in Ensdf2011, based on the observation of 6+ to 4+									**
$^{*246}\text{Am}^n$	D : gamma in $^{246}\text{Cm}$ , is not trusted by Nubase									**
$^{*246}\text{Es}^p$	E : other 19Vo03=151.9(2.0) keV									**
$^{*246}\text{Fm}$	D : %SF others 67Nu01=4.5(1.3) 96Ni09=15(5)									**
$^{*246}\text{Md}$	T : average 10An08=0.9(0.2) 96Ni09=1.0(0.4)									**
$^{247}\text{Pu}$	69210#	200#				2.27 d 0.23	$1/2^+ \#$	15	1983	$\beta^- = 100$
$^{247}\text{Am}$	67150#	100#				23.0 m 1.3	$5/2 \#$	15	1967	$\beta^- = 100$
$^{247}\text{Cm}$	65533	4				15.6 My0.5	$9/2^- *$	15	1954	$\alpha = 100$
$^{247}\text{Cm}^m$	65760	4	227.38	0.19		26.3 $\mu\text{s}$ 0.3	$5/2^+$	15	1968	IT=100
$^{247}\text{Cm}^n$	65938	4	404.90	0.03		100.6 ns 0.6	$1/2^+$	15	2003	IT=100
$^{247}\text{Bk}$	65490	5				1.38 ky 0.25	$3/2^-$	15	1965	$\alpha \approx 100; \text{SF} ?$
$^{247}\text{Cf}$	66109	14				3.11 h 0.03	$(7/2^+)$	15	1954	$\epsilon = 99.965 5; \alpha = 0.035 5$
$^{247}\text{Es}$	68578	19			*	4.55 m 0.26	$(7/2^+)$	15 89Ha27	J 1967	$\beta^+ \approx 93; \alpha \approx 7; \text{SF} ?$
$^{247}\text{Es}^m$	68630#	50#	50#	50#	*	> 20# $\mu\text{s}$	$(3/2^-)$			IT ?; $\beta^+ ?; \alpha ?$
$^{247}\text{Fm}$	71670#	180#				31 s 1	$(7/2^+)$	15	1967	$\alpha \approx 64; \beta^+ ?$
$^{247}\text{Fm}^m$	71720#	180#	49	8	AD	5.1 s 0.2	$(1/2^+)$	15	1967	$\alpha = 88 2; \beta^+ ?; \text{IT} ?$ *
$^{247}\text{Md}$	75940#	210#				1.19 s 0.09	$7/2^- \#$	15 10An08	TJD 1981	$\alpha \approx 100; \text{SF} < 0.1$ *
$^{247}\text{Md}^m$	76200#	210#	260	40	AD	250 ms 40	$1/2^- \#$	15 10An08	TJD 1993	$\alpha = 79 5; \text{SF} = 21 5$
$^{*247}\text{Fm}^m$	D : %IT from 06He27=12(2), but no direct gamma-ray decay was observed									**
$^{*247}\text{Md}$	T : average 10An08=1.2(0.1) 93Ho.A=1.12(0.22)									**
$^{248}\text{Am}$	70560#	200#				3# m	$3^+ \#$	14		$\beta^- ?$
$^{248}\text{Cm}$	67392.7	2.4				348 ky 6	$0^+$	14	1956	$\alpha = 91.61 16; \text{SF} = 8.39 16;$ $2\beta^- ?$
$^{248}\text{Cm}^m$	68850.8	2.6	1458.1	1.0		146 $\mu\text{s}$ 18	$8^- \#$	19Sh34	ETJ 2012	IT=100
$^{248}\text{Bk}$	68130	50			*	> 9 y	$6^+ \#$	14	1956	$\alpha ?; \epsilon ?$
$^{248}\text{Bk}^m$	68108	21	-20	50	*	23.7 h 0.2	$1(-)$	14	1956	$\beta^- = 70 5; \epsilon = 30 5; \alpha ?$
$^{248}\text{Bk}^p$	68180#	70#	50#	50#			$(5^-)$			
$^{248}\text{Cf}$	67238	5				333.5 d 2.8	$0^+$	14	1954	$\alpha \approx 100; \text{SF} = 0.0029 3$
$^{248}\text{Es}$	70300#	50#				24 m 3	$2^- \#$	14	1956	$\beta^+ \approx 100; \alpha \approx 0.25;$ *
										$\beta^+ \text{SF} = 3.5e-4 18$
$^{248}\text{Fm}$	71898	8				34.5 s 1.2	$0^+$	14	1958	$\alpha \approx 100; \beta^+ ?; \text{SF} = 0.10 5$
$^{248}\text{Fm}^m$	73100#	100#	1200#	100#		10.1 ms 0.6	$6^+ \#$	14	2010	IT ?; $\alpha ?; \beta^+ ?$
$^{248}\text{Md}$	76950#	180#				7 s 3		14	1973	$\beta^+ = 80 10; \alpha = 20 10;$ $\beta^+ \text{SF} < 0.05$
$^{248}\text{No}$	80690#	220#				< 2us	$0^+$	14 03Be18	I 2003	SF ?
$^{*248}\text{Es}$	D : % $\beta^+ \text{SF}$ from 01Sh09; other 80Ga07=3e-5%									**
$^{249}\text{Am}$	73100#	300#				3# m	$5/2 \#$			$\beta^- ?$



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>251</sup> Fm <sup>m</sup>	T : average 18Re07=23.7(1.1) 11As03=21.1(1.9) 06He20=21(3) 71Di03=15.2(2.3)								
* <sup>251</sup> Md	T : average 06Ch52=4.27(0.26) 73Es01=4.0(0.5)								
* <sup>251</sup> No	D : % $\alpha$ symmetrized from 01He35=91(+9-22)%								
* <sup>251</sup> No <sup>n</sup>	E : 142.4(0.5) + 203.1(0.5) + 782.5(0.6) keV gammas in a cascade to 7/2+ gs								
* <sup>251</sup> No <sup>n</sup>	J : expected conf=n <sup>3</sup> (1/2[631],7/2[624],9/2[734]),K=17/2-								
<sup>252</sup> Cm	79060#	300#			1# m	0 <sup>+</sup>	06		$\beta^-$ ?; $\alpha$ ?
<sup>252</sup> Bk	78540#	200#			1.8 m 0.5		06 92Kr.A	TD 1988	$\beta^-$ ?; $\alpha$ ?
<sup>252</sup> Cf	76034.6	2.4			2.645 y 0.008	0 <sup>+</sup>	06 18Be29	D 1954	$\alpha$ =96.8972 27; SF=3.1028 27
<sup>252</sup> Es	77290	50			471.7 d 1.9	(4 <sup>+</sup> )	06 FGK12a	J 1956	$\alpha$ =78 2; $\epsilon$ =22 2
<sup>252</sup> Fm	76817	5			25.39 h 0.04	0 <sup>+</sup>	06		$\alpha$ $\approx$ 100;SF=0.0023 2;2 $\beta^+$ ?
<sup>252</sup> Md	80470	90			2.3 m 0.8	1 <sup>+</sup> #	06		$\beta^+$ $\approx$ 100; $\alpha$ ?
<sup>252</sup> Md <sup>p</sup>	80550	80	80	120		am			
<sup>252</sup> No	82871	9			2.467 s 0.016	0 <sup>+</sup>	06 11Ga19	T 1967	$\alpha$ =67.6 5;SF=31.3 4; $\beta^+$ =1.1 3
<sup>252</sup> No <sup>m</sup>	84125	9	1254.1	1.6	109.1 ms 2.5	(8 <sup>-</sup> )	11Lo06	T 2007	IT=100
<sup>252</sup> Lr	88540#	190#			369 ms 75	7 <sup>-</sup> #	06 08Ne01	TD 2001	$\alpha$ $\approx$ 98;SF $\approx$ 2; $\beta^+$ ?
<sup>252</sup> Lr <sup>p</sup>	88710#	190#	170	30	AD				
* <sup>252</sup> Es	J : strong direct $\epsilon$ feeding to 3+ level at 969.8 keV in <sup>252</sup> Cf and the								
* <sup>252</sup> Es	J : expected p7/2[633]->n7/2[613] configuration change								
* <sup>252</sup> No	T : average 11Ga19=2.47(0.02) 06Le29=2.46(0.05) ( $\alpha$ (t)) 2.54(0.07)								
* <sup>252</sup> No	T : (SF(t)) 12Su22=2.43(0.13) 01Og08=2.44(0.04); others 12Sv02=2.3(0.1)								
* <sup>252</sup> No	T : 04He28=2.52(0.22) 03Be18=2.38(+0.26-0.22)								
* <sup>252</sup> No	D : %SF average 01Og08=32.2(0.5)% 11Ga19=29.3(0.9)% 03Be18=32(3)								
* <sup>252</sup> No	D : 77Be09=26.9(1.9); % $\beta^+$ and % $\alpha$ from $\beta^+/\alpha$ =0.016(0.005) in								
* <sup>252</sup> No	D : in 02He01; other $\alpha$ /SF=3.3(0.8) in 06Le29 in conflict								
* <sup>252</sup> No <sup>m</sup>	E : from a least-squares fit to the gamma rays in 07Su19								
* <sup>252</sup> No <sup>m</sup>	T : average 11Lo06=110(8) 08Ro21=109(6) 12Su22=109(3), supersedes								
* <sup>252</sup> No <sup>m</sup>	T : 07Su19=110(10)								
* <sup>252</sup> Lr	T : average 08Ne01=270(+180-80) 01He35=360(+110-70)								
* <sup>252</sup> Lr	D : %SF 76Og02 $\sim$ 2%								
<sup>253</sup> Bk	80930#	360#			60# m	3/2 <sup>-</sup> #	13 91Kr.A	I 1991	$\beta^-$ ?
<sup>253</sup> Cf	79302	4			17.81 d 0.08	(7/2 <sup>+</sup> )	13		$\beta^-$ =99.69 4; $\alpha$ =0.31 4
<sup>253</sup> Es	79010.5	1.2			20.47 d 0.03	7/2 <sup>+</sup> *	13 05Ah03	D 1954	$\alpha$ =100;SF=8.7e-6 3
<sup>253</sup> Es <sup>m</sup>	79117	4	106	4	> 10# $\mu$ s	3/2 <sup>-</sup> #	13 93Mo18	IJ	IT ?
<sup>253</sup> Fm	79345.5	1.5			3.00 d 0.12	1/2 <sup>+</sup>	13		$\epsilon$ =88 1; $\alpha$ =12 1
<sup>253</sup> Fm <sup>m</sup>	79486	6	140	6	> 100# ns	7/2 <sup>+</sup> #			IT ?
<sup>253</sup> Fm <sup>n</sup>	79697	6	351	6	560 ns 60	11/2 <sup>-</sup> #	13 11An13	ETJ 2011	IT=100
<sup>253</sup> Md	81170#	30#			12 m 8	(7/2 <sup>-</sup> )	13 05He27	D 1992	$\beta^+$ $\approx$ 100; $\alpha$ $\approx$ 0.7
<sup>253</sup> Md <sup>p</sup>	81230#	40#	60	30	1# m	1/2 <sup>-</sup> #	13		$\alpha$ ?;IT ?
<sup>253</sup> No	84359	7			1.57 m 0.02	9/2 <sup>-</sup> *	13 18Ra11	J 1967	$\alpha$ =55 3; $\beta^+$ ?;SF ?
<sup>253</sup> No <sup>m</sup>	84527	7	167.5	0.5	30.3 $\mu$ s 1.6	5/2 <sup>+</sup>	13 09He23	T 1973	$\alpha$ =100
<sup>253</sup> No <sup>n</sup>	85560	110	1196	107	706 $\mu$ s 24	19/2 <sup>+</sup> #	11Lo06	TJ 2011	IT=100
<sup>253</sup> No <sup>p</sup>	85620	110	1256	113	552 $\mu$ s 15	25/2 <sup>+</sup> #	13 11Lo06	JD 2011	IT=100
<sup>253</sup> Lr	88520	160		*	632 ms 46	(7/2 <sup>-</sup> )	13 01He35	TJD 1985	$\alpha$ =90 10;SF=1.0 6; $\beta^+$ ?
<sup>253</sup> Lr <sup>m</sup>	88560#	190#	30#	100#	1.32 s 0.14	(1/2 <sup>-</sup> )	13 09He20	TJD 1985	$\alpha$ =90 10;SF=12 3; $\beta^+$ ?; IT ?
<sup>253</sup> Rf	93640#	410#		*	13 ms 5	(7/2)( <sup>+</sup> #)	06 97He29	TJD 1997	SF $\approx$ 100; $\alpha$ ?
<sup>253</sup> Rf <sup>m</sup>	93840#	440#	200#	150#	52 $\mu$ s 14	(1/2)( <sup>+</sup> #)	06 97He29	TJD 1995	SF $\approx$ 100; $\alpha$ ?
* <sup>253</sup> Es	D : %SF from $\alpha$ /SF=1.15(0.03)e7 in 65Me02								
* <sup>253</sup> Fm	J : favored $\alpha$ decay to 416.6-keV level in <sup>249</sup> Cf (J=1/2+)								
* <sup>253</sup> Fm <sup>m</sup>	E : from 130-150 keV in 11An13								
* <sup>253</sup> Fm <sup>n</sup>	E : 211 keV above <sup>253</sup> Fm <sup>m</sup>								
* <sup>253</sup> Md	T : symmetrized from 92Ka08=6.4(+11.6-3.6)								
* <sup>253</sup> No	T : average 18Ac08=1.67(0.09) 17Mi01=1.7(0.2) 09He23=1.56(0.02)								
* <sup>253</sup> No	T : 09Qi04=1.57(+0.18-0.15) 67Mi03=95(10) 67Gh01=105(20)								
* <sup>253</sup> No	D : $\epsilon/e^+$ =0.45(0.03) in 11An13								
* <sup>253</sup> No <sup>m</sup>	T : average 09He23=28(3) 07Lo11=31.1(2.1) 73Be33=31.3(4.1); others								
* <sup>253</sup> No <sup>m</sup>	T : 11An13=22.7(0.5) 10St14=24(2) outliers								



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>253</sup> No <sup>n</sup>	E : greater than 1011 keV and less than 1380 keV									**
* <sup>253</sup> No <sup>n</sup>	T : 11Lo06=706(24); others: 07Lo11=970(210), 11An13=627(5) (r-ce-g(t))									**
* <sup>253</sup> No <sup>n</sup>	T : 650(15) (r-ce-802,713,209g(t))									**
* <sup>253</sup> No <sup>p</sup>	E : less than 120 keV above <sup>253</sup> No <sup>n</sup> in 11Lo06t									**
* <sup>253</sup> No <sup>p</sup>	T : from 11An13 using (r-ce-K x rays(t))									**
* <sup>253</sup> Lr	T : average 09He20=670(60) 11An13=552(15) 01He35=570(+70-60)									**
* <sup>253</sup> Lr	D : %SF from 17He08; other 01He35=1.3(+3.0-1.0)%									**
* <sup>253</sup> Lr <sup>m</sup>	T : 09He20 supersedes 01He35=1.49(+0.30-0.21); other 10He11=1.2(+0.7-0.4)									**
* <sup>253</sup> Lr <sup>m</sup>	D : %SF from 17He08; other 01He35=8(5)%									**
* <sup>253</sup> Rf	T : symmetrized from 97He29=11(+6-3)									**
* <sup>253</sup> Rf <sup>m</sup>	T : symmetrized from 97He29=48(+17-10)									**
<sup>254</sup> Bk	84390#	300#				1# m		19		$\beta^- ?$
<sup>254</sup> Cf	81341	11				60.5 d 0.2	0 <sup>+</sup>	19	1955	SF=99.69 2; $\alpha$ =0.31 2; $2\beta^- ?$
<sup>254</sup> Es	81994.2	2.9				275.7 d 0.5	7 <sup>+</sup>	19	1954	$\alpha \approx 100$ ; $\epsilon ?$ ; $\beta^- = 1.74e-4$ 8; SF<3e-6
<sup>254</sup> Es <sup>m</sup>	82074.5	2.7	80.4	1.1	AD	39.3 h 0.2	2 <sup>+</sup> *	19	FGK207 E	1954 $\beta^- = 98$ 2; IT<3; $\alpha = 0.32$ 1; $\epsilon = 0.076$ 7; SF<0.045
<sup>254</sup> Fm	80902.5	1.8				3.240 h 0.002	0 <sup>+</sup>	19	1954	$\alpha = 99.9408$ 3; SF=0.0592 3
<sup>254</sup> Md	83450#	100#	*			10 m 3	0 <sup>-</sup> #	19	1970	$\beta^+ \approx 100$ ; $\alpha ?$
<sup>254</sup> Md <sup>m</sup>	83500#	140#	50#	100#	*	28 m 8	3 <sup>-</sup> #	19	1970	$\beta^+ \approx 100$ ; $\alpha ?$
<sup>254</sup> No	84723	10				51.2 s 0.4	0 <sup>+</sup>	19	1966	$\alpha = 90$ 1; $\beta^+ = 10$ 1; SF=0.17 2
<sup>254</sup> No <sup>m</sup>	86019	10	1296.4	1.1		264.9 ms 1.4	(8 <sup>-</sup> )	19	1973	IT=100; SF=0.020 12; $\alpha < 0.01$
<sup>254</sup> No <sup>n</sup>	87940#	300#	3217#	300#		184 $\mu$ s 3	16 <sup>+</sup> #	19	10He10 EJT	2006 IT=100; SF $\leq$ 0.012
<sup>254</sup> Lr	89650	90				12.0 s 0.9	4 <sup>+</sup> #	19	19Vo03 T	1981 $\alpha = 71.7$ 19; $\beta^+ = 28.3$ 19; SF<0.1
<sup>254</sup> Lr <sup>m</sup>	89750	90	107	23	AD	20.3 s 4.1	1 <sup>+</sup> #	19	19Vo03 TE	2019 $\alpha ?$ ; $\beta^+ ?$ ; IT ?
<sup>254</sup> Rf	93200#	280#				22.9 $\mu$ s 1.0	0 <sup>+</sup>	19	20Kh01 T	1997 SF $\approx$ 100; $\alpha < 1.5$
<sup>254</sup> Rf <sup>m</sup>	94500#	340#	1300#	200#		4.3 $\mu$ s 0.7	8 <sup>-</sup> #	19	15Da12 JTD	2015 IT $\approx$ 100; SF<10
<sup>254</sup> Rf <sup>n</sup>	95200#	570#	2000#	500#		247 $\mu$ s 73	16 <sup>+</sup> #	19	15Da12 JT	2015 IT $\approx$ 100; SF<40
* <sup>254</sup> Es	J : favored $\alpha$ decay to <sup>250</sup> Bk <sup>n</sup> (J=7+)									**
* <sup>254</sup> Es <sup>m</sup>	T : other 19De11=51.8(16.1), probably a mixture between gs and isomer decays									**
* <sup>254</sup> No	T : other (recent) 18Mi11=54(4)									**
* <sup>254</sup> No <sup>m</sup>	T : average 11Lo06=259(17) 10Cl01=263(2) 10He10=275(7) 06He19=266(2)									**
* <sup>254</sup> No <sup>m</sup>	T : 06Ta19=266(10); other 73Gh03=280(40)									**
* <sup>254</sup> No <sup>n</sup>	E : 10He10=2917(3) + x keV; x=300#(300#); 10Cl01=2930(2), but their level									**
* <sup>254</sup> No <sup>n</sup>	E : scheme is disputed									**
* <sup>254</sup> Lr	T : average 19Vo03=11.9(0.9)(GSI) 01Ga20=13.4(4.2)(IMP); others (not used)									**
* <sup>254</sup> Lr	T : 08Ga25=17.8(+1.9-1.6)(LBNL) 08An16=18.4(1.8)(GSI)									**
* <sup>254</sup> Lr	T : 89Mu09=10.0(+4.5-2.4)(GSI) 85He22=13(+3-2)(GSI) 06Fo02=22(+9-6)(LBNL),									**
* <sup>254</sup> Lr	T : presumably affected by the longer-lived isomer									**
* <sup>254</sup> Lr	D : other (not used) % $\alpha$ =60(+11-15) % $\beta^+$ =40(+15-11) in 06Fo02									**
* <sup>254</sup> Rf	T : average 20Kh01=20(3) 15Da12=23.2(1.1) 97He29=23(3); other									**
* <sup>254</sup> Rf	T : 08Dr05=29.6(+0.7-0.6)									**
* <sup>254</sup> Rf <sup>m</sup>	T : average 15Da12=4.7(1.1) 20Kh01=4(1)									**
<sup>255</sup> Cf	84810#	200#				85 m 18	(7/2 <sup>+</sup> )	13	1981	$\beta^- = 100$ ; SF ?; $\alpha ?$
<sup>255</sup> Es	84089	11				39.8 d 1.2	(7/2 <sup>+</sup> )	13	1954	$\beta^- = 92.0$ 4; $\alpha = 8.0$ 4; SF=0.0041 2
<sup>255</sup> Fm	83800	4				20.07 h 0.07	7/2 <sup>+</sup>	13	1954	$\alpha = 100$ ; SF=2.4e-5 10
<sup>255</sup> Fm <sup>p</sup>	84031	4	231.1	0.2			9/2 <sup>+</sup>	13	2013	IT=100
<sup>255</sup> Md	84842	6	*			27 m 2	7/2 <sup>-</sup>	13	1958	$\beta^+ = 93$ 1; $\alpha = 7$ 1; SF ?
<sup>255</sup> Md <sup>p</sup>	84850#	70#	10#	70#	*	2# m	1/2 <sup>-</sup> #	13		$\alpha ?$ ; IT ?
<sup>255</sup> No	86812	14				3.52 m 0.18	(1/2 <sup>+</sup> )	13	11As03 TJ	1967 $\beta^+ = 70$ 5; $\alpha = 30$ 5
<sup>255</sup> No <sup>m</sup>	87020#	100#	210#	100#		1# s	11/2 <sup>-</sup> #			IT ?; $\alpha ?$
<sup>255</sup> No <sup>p</sup>	86910#	70#	100#	70#	Nm	> 100 ns	7/2 <sup>+</sup> #			IT ?
<sup>255</sup> Lr	89947	18				31.1 s 1.1	(1/2 <sup>-</sup> )	13	06Ch52 TJ	1971 $\alpha = 99.7$ 1; $\beta^+ = 0.3$ 1; SF ?
<sup>255</sup> Lr <sup>m</sup>	89988	19	41	8	AD	2.54 s 0.05	(7/2 <sup>-</sup> )	13	06Ch52 J	2006 IT ?; $\alpha \approx 40$
<sup>255</sup> Lr <sup>n</sup>	90743	22	796	12		< 1 $\mu$ s	(15/2 <sup>+</sup> )	13	2009	IT $\approx$ 100

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>255</sup> Lr <sup>p</sup>	91412	22	1465	12		1.78 ms 0.05	(25/2 <sup>+</sup> )	13	2008	IT≈100;α<0.15	*
<sup>255</sup> Rf	94330#	180#				1.63 s 0.05	(9/2 <sup>-</sup> )	13 15An05	D 1975	α=52.8 22;SF=47.2 22; β <sup>+</sup> <6	*
<sup>255</sup> Rf <sup>m</sup>	94480#	180#	150	22	AD	43 μs 9	(5/2 <sup>+</sup> )	15An05	ETJ 2015	IT=100	*
<sup>255</sup> Rf <sup>n</sup>	95380#	200#	1050	87		16 μs 5	19/2 <sup>+</sup> #	20Mo11	TED2020	IT=100	*
<sup>255</sup> Rf <sup>p</sup>	95630#	200#	1300	87		41 μs 10	25/2 <sup>+</sup> #	20Mo11	TED2020	IT=100	*
<sup>255</sup> Db	99600#	280#			*	54 ms	9/2 <sup>+</sup> #	13 17He08	TD 1976	SF≈67;α ?	*
<sup>255</sup> Db <sup>m</sup>	99700#	300#	100#	100#	*	2.8 ms	1/2 <sup>-</sup> #	17He08	TD 1976	SF≈100;α ?	*
* <sup>255</sup> Md	J : favored α decay to 7/2- level at 461.5 keV in <sup>251</sup> Es										**
* <sup>255</sup> Lr <sup>m</sup>	E : 740.0 keV above 9/2+ level, which is <30 keV above <sup>255</sup> Lr <sup>m</sup>										**
* <sup>255</sup> Lr <sup>p</sup>	E : 1408.6 keV above 9/2+, which is <30 keV above <sup>255</sup> Lr <sup>m</sup>										**
* <sup>255</sup> Lr <sup>p</sup>	T : average 09Je02=1.70(0.03) 08An16=1.81(0.02) (Birge ratio=3.05); other										**
* <sup>255</sup> Lr <sup>p</sup>	T : 08Ha31=1.4(0.1)										**
* <sup>255</sup> Rf	T : average 20Mo11=1.60(0.07) 06He27=1.68(0.09) 01He35=1.64(0.11);										**
* <sup>255</sup> Rf	T : other 20Kh01=1.9(0.2)										**
* <sup>255</sup> Rf	D : %SF average 20Mo11=51(7) 19Kh01=53(8) 15An05=45(3) 97He29=45(6)										**
* <sup>255</sup> Rf	D : 97He29=52(6); %β <sup>+</sup> from 15An05										**
* <sup>255</sup> Rf	J : favored α decay to the (9/2-) level at 203.6 keV in <sup>251</sup> No										**
* <sup>255</sup> Rf <sup>m</sup>	T : other 20Kh01>30 us										**
* <sup>255</sup> Rf <sup>m</sup>	T : symmetrized from 20Mo11=15(+6-4)										**
* <sup>255</sup> Rf <sup>n</sup>	E : 900-1200 keV above the gs in 20Mo11										**
* <sup>255</sup> Rf <sup>p</sup>	T : symmetrized from 20Mo11=38(+12-7)										**
* <sup>255</sup> Rf <sup>p</sup>	E : 1150-1450 keV above the gs in 20Mo11										**
* <sup>255</sup> Db	T : other 83OgZW=1.6(+0.6-0.4)										**
<sup>256</sup> Cf	87040#	310#				12.3 m 1.2	0 <sup>+</sup>	17	1980	SF=100;α ?;2β <sup>-</sup> ?	
<sup>256</sup> Es	87190#	100#			*&	7.6 h	7 <sup>+</sup> #	17	1976	β <sup>-</sup> ≈100;β <sup>-</sup> SF=0.002	
<sup>256</sup> Es <sup>m</sup>	87190#	140#	0#	100#	*&	25.4 m 1.2	0 <sup>+</sup> #	17 81Lo15	T 1981	β <sup>-</sup> =100	*
<sup>256</sup> Fm	85485	3				157.1 m 1.3	0 <sup>+</sup>	17	1955	SF=91.9 3;α=8.1 3	
<sup>256</sup> Md	87460#	120#			*	77.7 m 1.8	(1 <sup>-</sup> )	17	1955	β <sup>+</sup> =90.8 7;α=9.2 7;SF<3	
<sup>256</sup> Md <sup>m</sup>	87620	70	160#	100#	*	100# m	7 <sup>-</sup> #			β <sup>+</sup> ?;α ?;SF ?	
<sup>256</sup> Md <sup>p</sup>	87700#	120#	240#	140#			am				
<sup>256</sup> No	87823	8				2.91 s 0.05	0 <sup>+</sup>	17	1963	α=99.45 5;SF=0.55 5;ε ?	*
<sup>256</sup> Lr	91750	80				27.9 s 1.0	(0 <sup>-</sup> , 3 <sup>-</sup> )#	17	1965	α=85 10;β <sup>+</sup> =15 10;SF<0.03	
<sup>256</sup> Lr <sup>p</sup>	91980#	90#	230#	40#				17			
<sup>256</sup> Rf	94222	18				6.60 ms 0.05	0 <sup>+</sup>	17 20Mo11	T 1975	SF=99.69 10;α=0.31 10	*
<sup>256</sup> Rf <sup>m</sup>	95340#	100#	1120#	100#		25 μs 2	4 <sup>-</sup> #	17 15Ko14	J 2009	IT=100;SF ?	
<sup>256</sup> Rf <sup>n</sup>	95620#	100#	1400#	100#		17 μs 2	8 <sup>-</sup> #	17	2009	IT=100;SF ?	
<sup>256</sup> Rf <sup>p</sup>	96620#	200#	2400#	200#		27 μs 5		17	2009	IT=100;SF ?	*
<sup>256</sup> Db	100300#	190#				1.7 s 0.4	9 <sup>-</sup> #	17 01He35	TD 2001	α=70 11;β <sup>+</sup> =30 11;SF ?	*
* <sup>256</sup> Es <sup>m</sup>	T : 81Lo15=25.4(2.4), but the uncertainty is 2 standard deviations										**
* <sup>256</sup> No	D : %SF symmetrized from 90Ho03=0.53(+6-3)										**
* <sup>256</sup> Rf	T : average 20Mo11=6.75(0.49) 20Ku23=6.90(0.23) 18Sv02=5.75(0.17)										**
* <sup>256</sup> Rf	T : 13Ri07,12Gr12=6.9(0.2) 11Ro20=6.9(0.4) 10St14=5.1(1.0-0.7)										**
* <sup>256</sup> Rf	T : 09Je01=6.67(0.09) 08Dr05=6.70(0.09) 97He29=6.2(0.2) 84Og02=6.7(0.2)										**
* <sup>256</sup> Rf	D : %α average 20Ku23=0.29(+0.13-0.10) 97He29=0.32(0.017); other										**
* <sup>256</sup> Rf	D : %SF 10St14=97(+2-6)%										**
* <sup>256</sup> Rf <sup>p</sup>	T : other 20Mo10=18(7)										**
* <sup>256</sup> Db	T : symmetrized from 01He35=1.6(+0.5-0.3); other 83Og.A=2.6(+1.4-0.8)										**
<sup>257</sup> Es	89400#	410#				7.7 d 0.2	7/2 <sup>+</sup> #	13	1987	β <sup>-</sup> =100;α ?	
<sup>257</sup> Fm	88590	4				100.5 d 0.2	9/2 <sup>+</sup>	13 13As02	J 1964	α=99.790 4;SF=0.210 4	
<sup>257</sup> Md	88992.5	1.6				5.52 h 0.05	(7/2 <sup>-</sup> )	13	1965	ε=85 3;α=15 3;SF ?	
<sup>257</sup> No	90247	6				24.5 s 0.5	(3/2 <sup>+</sup> )	13 02Ho11	D 1967	α=85 8;β <sup>+</sup> =15 8;SF ?	
<sup>257</sup> No <sup>p</sup>	90550#	120#	300#	120#			9/2 <sup>+</sup> #				
<sup>257</sup> Lr	92670#	40#			*	6.0 s 0.4	7/2 <sup>-</sup> #	13 16He08	J 1971	α≈100;β <sup>+</sup> ?;SF ?	*
<sup>257</sup> Lr <sup>m</sup>	92770#	60#	100#	50#	*	0.27 s 0.12	1/2 <sup>-</sup> #	16He08	TI 2018	α ?; IT ?	*
<sup>257</sup> Lr <sup>p</sup>	92820#	110#	150#	100#			am	13			
<sup>257</sup> Rf	95866	11				5.0 s 0.2	(1/2 <sup>+</sup> )	13 13Ri07	T 1969	α=89.3 14;β <sup>+</sup> =9.4 14; SF=1.3 3	*
<sup>257</sup> Rf <sup>m</sup>	95940	10	73	11	AD	4.5 s 0.2	11/2 <sup>-</sup> #	13 10St14	TJ 1997	α=88.5 15;β <sup>+</sup> =11.5 15;	*

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens	Reference	Year of discovery	Decay modes and intensities (%)	
$^{257}\text{Rf}^n$	97022	10	1155	11	AD	106 $\mu\text{s}$ 6	$21/2^+\#$	13	13Ri07 TJ	2009	SF ?	
$^{257}\text{Db}$	100150	160			*	2.3 s 0.2	$9/2^+\#$	13	09He20 T	1985	IT=100	*
$^{257}\text{Db}^m$	100290#	200#	140#	110#	*	670 ms 60	$(1/2^-)$	13	09He20 T	1985	$\alpha>94$ ;SF<6; $\beta^+$ ?	*
$^{*257}\text{Lr}$	T : average 10St14=6.3(+0.9-0.7) and 5.8(0.5); others (not used)											**
$^{*257}\text{Lr}$	T : 97He29=3.3(+0.5-0.4), 4.3(+1.3-0.8) 76Be.A=0.646(0.025)											**
$^{*257}\text{Lr}$	T : 71Es01=0.6(0.1)											**
$^{*257}\text{Lr}^m$	J : direct $\beta^+$ decay feeding from $^{257}\text{Rf}$ ( $J=1/2+$ )											**
$^{*257}\text{Lr}^m$	T : symmetrized from 16He08=0.203(+0.164-0.063)											**
$^{*257}\text{Rf}$	J : favorite $\alpha$ decay to the $(1/2+)$ level at 670 keV in $^{253}\text{No}$											**
$^{*257}\text{Rf}$	T : average 13Ri07=6.1(0.5) 10St14=5.5(0.4) 10Be16=4.8(0.2) 09Qi04=4.7(0.3);											**
$^{*257}\text{Rf}$	T : others 85So03=3.8(0.8) 74Be.A=4.8(0.3) 71Gh03=4.8(0.5)											**
$^{*257}\text{Rf}$	D : $\beta^+$ from 16He08, stated that 10St14=19.4(1.4) is a misprint; other											**
$^{*257}\text{Rf}$	D : 09Qi04=2(1)%											**
$^{*257}\text{Rf}^m$	E : other 97He29=118(4) keV from direct comparison of two alpha lines											**
$^{*257}\text{Rf}^m$	T : average 13Ri07=4.7(0.4) 10St14=4.9(0.7) 10Be16=4.6(0.3) 97He29=3.9(0.4)											**
$^{*257}\text{Rf}^m$	T : 08Dr05=4.1(+0.7-0.6) 09Qi04=4.1(+2.4-1.3)											**
$^{*257}\text{Rf}^m$	D : $\beta^+$ from 16He08											**
$^{*257}\text{Rf}^n$	E : 1082(2) keV above $^{257}\text{Rf}^m$ in 10Be16											**
$^{*257}\text{Rf}^n$	T : others 10Be16=134.9 (7.7), reanalyzed in 13Ri07 to 10Be16=110(5)											**
$^{*257}\text{Rf}^n$	T : 20Mo10=105(19) 09Je01=109(13) (same group as 13Ri07)											**
$^{*257}\text{Rf}^n$	T : 09Qi04=160(+42-31)											**
$^{*257}\text{Db}$	T : from 09He20, supersedes 01He35=1.50(+0.19-0.15); 10He11=1.5(+0.9-0.4)											**
$^{*257}\text{Db}^m$	T : from 09He20, supersedes 01He35=760(+150-110); 10He11=360(+220-90)											**
$^{*257}\text{Db}^m$	J : favorite $\alpha$ decay to $^{253}\text{Lr}^m$ [ $J=(1/2^-)$ ]											**
$^{258}\text{Es}$	92700#	400#				4# m					$\beta^-$ ?; $\alpha$ ?	
$^{258}\text{Fm}$	90430#	200#				370 $\mu\text{s}$ 14	$0^+$	17	86Hu05 T	1971	SF $\approx$ 100; $\alpha$ ?	
$^{258}\text{Md}$	91690	3			*	51.59 d 0.29	$8^-$	17	93Mo18 D	1970	$\alpha\approx$ 100; $\beta^+$ <0.0015; $\beta^-$ <0.0015	*
$^{258}\text{Md}^m$	91690#	200#	0#	200#	*	57.0 m 0.9	$1^-$	17	93Mo18 D	1980	$\epsilon=85$ 15;SF<15; $\beta^-$ ?; $\alpha<$ 1.2	*
$^{258}\text{No}$	91480#	100#				1.23 ms 0.12	$0^+$	17	18Br13 T	1989	SF $\approx$ 100; $\alpha$ ?	*
$^{258}\text{Lr}$	94780#	100#				3.92 s 0.33		17	14Ha04 TD	1971	$\alpha=97.4$ 18; $\beta^+=2.6$ 18	
$^{258}\text{Lr}^p$	95020#	140#	240#	100#			<i>am</i>					
$^{258}\text{Rf}$	96344	16				12.5 ms 0.5	$0^+$	17	20Mo11 T	1969	SF=95.1 16; $\alpha=4.9$ 16	*
$^{258}\text{Rf}^m$	97540#	300#	1200#	300#		3.4 ms 1.7		17	16He15 ITD	2016	IT= ?; $\alpha$ ?; $\beta^+$ ?	*
$^{258}\text{Rf}^n$	97840#	500#	1500#	500#		15 $\mu\text{s}$ 10		17	16He15 ITD	2016	IT ?; SF ?	
$^{258}\text{Db}$	101510	90			&	2.17 s 0.36	$0^-$	17	19Vo03 TJ	1985	$\alpha=64$ 10; $\beta^+=36$ 10	*
$^{258}\text{Db}^m$	101560	90	53	14	AD &	4.41 s 0.21	$5^+\#$	17	19Vo03 TEJ	1981	$\alpha=77$ 8; $\beta^+=23$ 8;SF ?	*
$^{258}\text{Sg}$	105300#	410#				2.7 ms 0.5	$0^+$	17	17He08 TD	1997	SF $\approx$ 100; $\alpha$ ?	*
$^{*258}\text{Md}$	D : derived from: "the sum of SF, $\epsilon$ and $\beta^-$ decay branches < 0.003%" in											**
$^{*258}\text{Md}$	D : 93Mo18 and T(SF)>150000 y, from 86Lo16, thus %SF<1e-4%											**
$^{*258}\text{Md}^m$	D : %SF<15 derived from 93Mo18 "the sum of SF and $\beta^-$ decay branches < 30%"											**
$^{*258}\text{No}$	T : average 18Br13=1.24(+0.16-0.14) 89Hu09=1.2(0.2)											**
$^{*258}\text{Rf}$	T : average 20Mo11=8.79(1.12) 19He17=14.2(+1.2-0.4) 16He15=10.0(1.1)											**
$^{*258}\text{Rf}$	T : 08Ga08=14.7(+1.2-1.0) 85So03=13(3) 69Gh01=11(2); other											**
$^{*258}\text{Rf}$	T : 20Ku23=12(+16-10)											**
$^{*258}\text{Rf}^m$	T : symmetrized from 16He15=2.4(+2.4/-0.8)											**
$^{*258}\text{Db}$	T : others 16He15=3.6(0.3), 2.8(0.6) 09He20=1.9(0.5)											**
$^{*258}\text{Db}$	D : from 09He20											**
$^{*258}\text{Db}^m$	T : others 16He15=4.4(1.0) 09He20=4.3(0.5) 06Fo02=4.8(+1.0-0.8)											**
$^{*258}\text{Db}^m$	T : 01Ga20=4.3(1.1) 85He22=4.4(+0.9-0.6)											**
$^{*258}\text{Db}^m$	D : from 09He20; others $\beta^+$ 06Fo02=39(+11-9) 85He22=33(+9-5)											**
$^{*258}\text{Sg}$	T : symmetrized from 17He08=2.6(+0.6-0.4), determined by combining data											**
$^{*258}\text{Sg}$	T : from 09Fo02, 02Pa.A and 97He29											**
$^{259}\text{Fm}$	93700#	280#				1.5 s 0.2		13		1980	SF=100	
$^{259}\text{Md}$	93560#	100#				1.60 h 0.06	$7/2^-$	13		1982	SF $\approx$ 100; $\alpha$ ?	
$^{259}\text{No}$	94079	6				58 m 5	$9/2^+$	13	13As02 J	1973	$\alpha=75$ 4; $\epsilon=25$ 4;SF<10	
$^{259}\text{No}^p$	94310#	150#	230#	150#								

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{259}\text{Lr}$	95850#	70#				6.2 s 0.3	$1/2^-$	13	1971	$\alpha=78.2$ ; SF=22.2; $\beta^+$ ?	
$^{259}\text{Lr}^p$	96200#	170#	350#	150#							
$^{259}\text{Rf}$	98370#	70#				2.63 s 0.26	$3/2^+$	13 08Ga08	T 1969	$\alpha=85.4$ ; $\beta^+=15.4$ ; SF<3	
$^{259}\text{Rf}^p$	98430#	100#	60	70	Nm		$(7/2^+)$				*
$^{259}\text{Rf}^q$	98570#	110#	210	90	Nm		$(9/2^+)$				*
$^{259}\text{Db}$	101990	60				510 ms 160	$9/2^+$	13 01Ga20	D 2001	$\alpha=100$	
$^{259}\text{Sg}$	106520#	180#				402 ms 56	$(11/2^-)$	13 15An05	TJD 1985	$\alpha\approx 100$ ; SF ?; $\beta^+$ ?	
$^{259}\text{Sg}^m$	106610#	180#	87	22	AD	226 ms 27	$(1/2^+)$	15An05	TJD 2015	$\alpha\approx 97.1$ ; SF $\approx 3.1$ ; $\beta^+$ ?	
$^{*259}\text{Rf}$	T : average 08Ga08=2.5(+0.4-0.3) 94Gr08=1.7(+0.8-0.5);										**
$^{*259}\text{Rf}$	T : others 06Gr24=1.9(+1.3-0.5) 04Fo08=2.2(+1.7-0.8) 03Gi05=4.0(+7.3-1.6)										**
$^{*259}\text{Rf}$	T : 98Ho13=2.6(+1.4-0.7) 85So03=3.4(1.7) 81Be03=3.0(1.3)										**
$^{*259}\text{Rf}$	T : 73Dr10=3.2(0.8) 69Gh01=3.2(0.8); 10Ni14(1 event)=107 ms										**
$^{*259}\text{Rf}$	I : 08Ga08 suggest existence of an isomer formed only in direct production										**
$^{*259}\text{Rf}$	D : % $\beta^+$ 08Ga08=15(4)% to $^{259}\text{Lr}$ followed by SF; %SF 17He08<3%										**
$^{260}\text{Fm}$	95770#	440#			EU	1# m	$0^+$			SF ?	
$^{260}\text{Md}$	96550#	320#				27.8 d 0.8		99 92Lo.B	TD 1989	SF $\approx 100$ ; $\alpha < 5$ ; $\epsilon < 5$ ; $\beta^- < 3.5$	
$^{260}\text{No}$	95610#	200#				106 ms 8	$0^+$	99	1985	SF=100	
$^{260}\text{Lr}$	98280#	130#				3.0 m 0.5		99	1971	$\alpha=80.20$ ; $\beta^+=20.20$	
$^{260}\text{Rf}$	99150#	200#				21 ms 1	$0^+$	99 85So03	T 1985	SF $\approx 100$ ; $\alpha$ ?; $\beta^+$ ?	
$^{260}\text{Db}$	103670#	90#				1.52 s 0.13		99 77Be36	TD 1970	$\alpha=90.4.6$ ; SF=9.6.6; $\beta^+$ ?	
$^{260}\text{Db}^p$	103770#	180#	100#	150#							*
$^{260}\text{Sg}$	106547	21				4.95 ms 0.33	$0^+$	99 09He20	TD 1984	SF=71.3; $\alpha=29.3$	
$^{260}\text{Bh}$	113120#	200#				41 ms 14		16 08Ne01	TD 2008	$\alpha\approx 100$ ; $\beta^+$ ?; SF ?	
$^{*260}\text{Fm}$	I : T1/2~4 ms and %SF=100 mode were reported in the 92Lo.B, but the										**
$^{*260}\text{Fm}$	I : results were not confirmed in the subsequent experiment by same										**
$^{*260}\text{Fm}$	I : group (97Lo.A)										**
$^{*260}\text{Md}$	T : from 92Lo.B supersedes 86Hu01=31.8(0.5), same group										**
$^{*260}\text{No}$	T : other 19De11=155(+212-57)										**
$^{*260}\text{Rf}$	T : others 08Ga08=22.2(+3.0-2.4) 08Go.A=21(+7.3,-4.3) 13Mu08=12(11)										**
$^{*260}\text{Db}$	T : others 04Mo26=1.5(+0.8-0.4) 04Ga29=0.89(+0.79-0.35) 70Gh02=1.6(0.3)										**
$^{*260}\text{Db}$	T : 71Dr01=1.4(+0.6)0.3										**
$^{*260}\text{Sg}$	T : supersedes 85Mu11=3.6(+0.9-0.6)										**
$^{*260}\text{Sg}$	D : other 85Mu11 %SF=50(+30-20)% and % $\alpha=50(+20-30)$ %										**
$^{*260}\text{Bh}$	T : symmetrized from 08Ne01=35(+19-9)										**
$^{261}\text{Md}$	98580#	510#				40# m	$7/2^-$			$\alpha$ ?	
$^{261}\text{No}$	98460#	200#				3# h				$\alpha$ ?	
$^{261}\text{Lr}$	99560#	200#				39 m 12	$1/2^-$	99	1987	SF $\approx 100$ ; $\alpha$ ?	
$^{261}\text{Rf}$	101320	70			*&	2.1 s 0.2	$3/2^+$	15 11Ha13	TD 1970	SF=82.4; $\alpha=18.4$	
$^{261}\text{Rf}^m$	101390#	120#	70#	100#	*&	74 s 5	$11/2^-$	15 13Mu08	T 1970	$\alpha\approx 100$ ; $\beta^+$ ?; SF ?	
$^{261}\text{Rf}^p$	101550#	120#	230#	100#							*
$^{261}\text{Db}$	104310#	110#				4.7 s 1.0	$9/2^+$	99 13Su04	TD 1970	SF=73.11; $\alpha=27.11$	
$^{261}\text{Db}^p$	104590#	230#	280#	200#							*
$^{261}\text{Sg}$	108005	18				183 ms 5	$(3/2^+)$	99 10St14	TJD 1984	$\alpha=98.1.4$ ; $\beta^+=1.3.3$ ; SF=0.6.2	
$^{261}\text{Sg}^m$	108110#	50#	100#	50#		9.3 $\mu$ s 1.8	$7/2^+$	99 10Be16	T 2010	IT=100	
$^{261}\text{Bh}$	113080	180				12.8 ms 3.2	$(5/2^-)$	99 10He11	TJD 1989	$\alpha\approx 100$ ; SF ?	
$^{*261}\text{Rf}$	T : average 12Ha05=2.6(+0.7-0.5) 11Ha13=1.9(0.4) ( $\alpha(t)$ ) and										**
$^{*261}\text{Rf}$	T : 1.8(0.4) (SF(t)) 08Go.A=2.2(+0.9-0.5) 08Dv02=3(1) 96La11=2.1(0.2);										**
$^{*261}\text{Rf}$	T : others 02Ho11=4.2(+3.4-1.3), 13Mu08=3.9(3.0) 15Mo25=4.7(+3.6-1.4)										**
$^{*261}\text{Rf}$	T : 08Mo09=2 events at 2.97 and 8.3s 94La22=1.2(+1.0-0.5) originally										**
$^{*261}\text{Rf}$	T : attributed to $^{262}\text{Rf}$ , but re-assigned in 08Go.A and 11Ha13 to										**
$^{*261}\text{Rf}$	T : $^{261}\text{Rf}$										**
$^{*261}\text{Rf}$	D : %SF average 11Ha13=73(6) 12Ha05=82(9) 13Mu08=88(5); other 08Dv02=91										**
$^{*261}\text{Rf}^m$	T : average 02Ho11=78(+11-6) 00Sy01=74(+7-6) 71Gh01=65(10); others										**
$^{*261}\text{Rf}^m$	T : 13Mu08=19(+5-3) 12Ha05=59(42) 08Dv02=20(+110-10) 08Ga08=71(+342-33)										**
$^{*261}\text{Db}$	T : average 13Su04=4.7(+3.6-1.4) 10St14=4.1(+1.4-0.8); others										**
$^{*261}\text{Db}$	T : 04Ga29=1.70(+0.79-0.49) 71Fl02=1.8(0.6) 71Gh01=1.8(0.6)										**
$^{*261}\text{Db}$	D : from 13Su04 where 11 SF and 4 $\alpha$ events were observed;										**
$^{*261}\text{Db}$	D : uncertainty estimated by Nubase										**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>261</sup> Sg	T : average 10St14=184(5), supersedes 09He20=195(15), 10Be16=178(14)						**
* <sup>261</sup> Sg <sup>m</sup>	T : symmetrized from 10Be16=9.0(+2.0-1.5)						**
* <sup>261</sup> Bh	T : symmetrized from 10He11=11.8(+3.9-2.4); others 06Fo02=10(+14-5)						**
* <sup>261</sup> Bh	T : 08Ne08=6.7(+3.8-1.8) 89Mu09=11.8(+5.3-2.8), superseded by 10He11						**
* <sup>261</sup> Bh	D : no SF decays were observed in 10He11 (%SF<5)						**
<sup>262</sup> Md	101670#	450#	3# m				SF ?; $\alpha$ ?
<sup>262</sup> No	100100#	360#	~ 5 ms	0 <sup>+</sup>	01	1988	SF=100; $\alpha$ ?
<sup>262</sup> Lr	102110#	200#	~ 4 h		01	1987	$\beta^+$ = ?; SF<10; $\alpha$ ?
<sup>262</sup> Rf	102390#	220#	250 ms 100	0 <sup>+</sup>	01 08Go.A	TD 1985	SF≈100
<sup>262</sup> Rf <sup>m</sup>	103390#	460#	47 ms 5	(8 <sup>-</sup> , 9 <sup>-</sup> )#	85So03	TD 1978	SF=100
<sup>262</sup> Db	106250#	140#	34 s 4		01 14Ha04	TD 1971	SF=52 4; $\alpha$ =48 4; $\beta^+$ ?
<sup>262</sup> Db <sup>p</sup>	106300#	160#					$\alpha$ ?
<sup>262</sup> Sg	108369	22	10.3 ms 1.7	0 <sup>+</sup>	01 17He08	D 2001	SF=94 6; $\alpha$ ?
<sup>262</sup> Sg <sup>p</sup>	109230	110	860 110 AD	9 <sup>-</sup> #			
<sup>262</sup> Bh	114250	90	84 ms 11		01 09He20	T 1981	$\alpha$ ≈100; SF<20
<sup>262</sup> Bh <sup>m</sup>	114470	110	9.5 ms 1.6		01 06Fo02	T 1981	$\alpha$ ≈100; SF ?
* <sup>262</sup> Rf	T : symmetrized from 08Go.A=210(+128-58), 7 SF events; others 85So03=1.3(1)						**
* <sup>262</sup> Rf	T : 96La11=2.1(0.2) 94La22=1.2(+1.0-0.5) 98Tu01=2.5(+2.4-1.6). 11Ha13 and						**
* <sup>262</sup> Rf	T : 08Go.A suggested that the long-lived activities belong to <sup>261</sup> Rf						**
* <sup>262</sup> Rf <sup>m</sup>	I : assigned as a K isomer in 96La11						**
* <sup>262</sup> Db	T : symmetrized from 14Ha04=33.8(+4.4-3.5); other 15Mo25=39(+53-14)						**
* <sup>262</sup> Sg	T : average 06Gr24=15(+5-3) 17He08=8.5(+2.3-1.5), determined by merging						**
* <sup>262</sup> Sg	T : data from 01Ho06 and 12Ac04						**
* <sup>262</sup> Sg <sup>p</sup>	J : favored $\alpha$ decay from <sup>266</sup> Hs <sup>m</sup> (J=9-#)						**
* <sup>262</sup> Bh	T : average 09He20=83(14), supersedes 89Mu09=102(26) 06Fo02=84(+21-16);						**
* <sup>262</sup> Bh	T : other 08Ne08=120(+55-29)						**
* <sup>262</sup> Bh <sup>m</sup>	T : average 06Fo02=9.6(+3.6-2.4) 97Ho14(11 events)=12.2(+5.5-2.8)						**
* <sup>262</sup> Bh <sup>m</sup>	T : 89Mu09=8.0(2.1); others 09He20=22(4) 08Ne08(4 events)=16(+14-5)						**
<sup>263</sup> No	103130#	490#	20# m				$\alpha$ ?; SF ?
<sup>263</sup> Lr	103670#	220#	5# h	1/2 <sup>-</sup> #			$\alpha$ ?
<sup>263</sup> Rf	104760#	150#	11 m 3		99 03Kr20	TD 2003	SF≈100; $\alpha$ ?
<sup>263</sup> Rf <sup>p</sup>	105060#	250#	300# 200#				
<sup>263</sup> Db	107110#	170#	29 s 9	9/2 <sup>+</sup> #	99 92Kr01	DT 1992	SF=56 14; $\alpha$ =37 14; $\beta^+$ =6.9 16
<sup>263</sup> Db <sup>p</sup>	107370#	260#	260# 200#				
<sup>263</sup> Sg	110200#	100#	940 ms 140	3/2 <sup>+</sup> #	99 06Gr24	TD 1974	$\alpha$ =87 8; SF=13 8
<sup>263</sup> Sg <sup>m</sup>	110250#	100#	51 19 Nm*	7/2 <sup>+</sup> #	99 04Fo08	T 1995	$\alpha$ ≈100; SF ?; IT ?
<sup>263</sup> Sg <sup>p</sup>	110290#	100#	100 30 AD				
<sup>263</sup> Bh	114500#	310#	200# ms	5/2 <sup>-</sup> #	99		$\alpha$ ?
<sup>263</sup> Hs	119680#	200#	0.9 ms 0.4	3/2 <sup>+</sup> #	99 09Dr02	TD 2009	$\alpha$ ≈100; SF ?
<sup>263</sup> Hs <sup>m</sup>	120000#	200#	1# ms	11/2 <sup>-</sup> #			$\alpha$ ≈100; SF ?
* <sup>263</sup> Rf	T : average 03Kr20=24(+19-7)m 93Gr.C=500(+300-200)s 92Cz.A=600(+300-200)s;						**
* <sup>263</sup> Rf	T : other 08Dv02=8(+40-4) s using one SF event						**
* <sup>263</sup> Db	D : %SF symmetrized from 92Kr01=57(+13-15)%; % $\beta^+$ average 03Kr20=3(+4-1)%						**
* <sup>263</sup> Db	D : 93Gr.C=8(2)%						**
* <sup>263</sup> Db	T : symmetrized from 92Kr01=27(+10-7); other 98Ik02=54(+98-21) from SF(t)						**
* <sup>263</sup> Sg	T : average 06Gr24=820(+370-190) 94Gr08=553(+336-152) 74Gh04=900(200), all						**
* <sup>263</sup> Sg	T : produced in direct reaction population						**
* <sup>263</sup> Sg <sup>m</sup>	T : average 04Fo08=290(+170-90) 04Mo40=549(+300-143) 03Gi05=222(+404-87)						**
* <sup>263</sup> Sg <sup>m</sup>	T : 98Ho13=310(+160-80), all produced via $\alpha$ decay of <sup>267</sup> Hs;						**
* <sup>263</sup> Sg <sup>m</sup>	T : other 10Ni14= $\tau$ =702 ms via $\alpha$ decay of <sup>267</sup> Hs						**
* <sup>263</sup> Hs	T : symmetrized from 09Dr02=0.74(+0.48-0.21) 6 events observed						**
* <sup>263</sup> Hs	D : no SF observed in 09Dr02 (%SF<8.4)						**
<sup>264</sup> No	105010#	590#	1# m	0 <sup>+</sup>			$\alpha$ ?; SF ?
<sup>264</sup> Lr	106380#	440#	10# h				$\alpha$ ?; SF ?
<sup>264</sup> Rf	106080#	360#	1# h	0 <sup>+</sup>			$\alpha$ ?
<sup>264</sup> Db	109260#	240#	3# m				$\alpha$ ?

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>264</sup> Sg	110780#	280#				78 ms 25	0 <sup>+</sup>	06 17He08	TD 2006	SF>80; $\alpha$ ?	*
<sup>264</sup> Bh	115960#	180#				1.07 s 0.21		99 04Mo26	TD 1995	$\alpha \approx 86$ ; SF=14; $\beta^+$ ?	*
<sup>264</sup> Bh <sup>p</sup>	116290#	230#	330#	150#			<i>am</i>				
<sup>264</sup> Hs	119563	29				0.7 s 0.3	0 <sup>+</sup>	99 17He08	TD 1986	$\alpha=70$ 30; SF=30 30	*
* <sup>264</sup> Sg	T : symmetrized from 17He08=68(+32-16), determined by merging data from										**
* <sup>264</sup> Sg	T : 10Ni14, 06Ni10 and 06Gr24										**
* <sup>264</sup> Sg	D : no $\alpha$ observed in 17He08 ( $\% \alpha < 36$ )										**
* <sup>264</sup> Bh	T : average 04Mo26=0.9(+0.3-0.2) 04Ga29=1.17(+0.88-0.44)										**
* <sup>264</sup> Bh	T : 02Ho11=1.02(+0.69-0.29)										**
* <sup>264</sup> Hs	T : symmetrized from 17He08=0.63(+0.34-0.16), determined by merging data										**
* <sup>264</sup> Hs	T : from 87Mu15 and 11Sa41										**
* <sup>264</sup> Hs	D : %SF symmetrized from 17He08=20(+40-20), determined by merging data										**
* <sup>264</sup> Hs	D : from 87Mu15 and 11Sa41										**
<sup>265</sup> Lr	108230#	550#				10# h	1/2 <sup>-</sup> #			$\alpha$ ?; SF ?	
<sup>265</sup> Rf	108690#	360#				1.6 m 0.6	9/2 <sup>+</sup> #	15 16Ho09	TD 2010	SF $\approx$ 100; $\alpha$ ?	*
<sup>265</sup> Db	110380#	220#				15# m	9/2 <sup>+</sup> #			$\alpha$ ?	
<sup>265</sup> Sg	112790#	140#			*	9.2 s 1.6	11/2 <sup>-</sup> #	15 12Ha05	TD 1994	$\alpha > 50$ ; SF ?	*
<sup>265</sup> Sg <sup>m</sup>	112790#	130#	-10#	160#	*	16.4 s 2.4		15 12Ha05	TD 1994	$\alpha > 50$ ; SF ?	*
<sup>265</sup> Bh	116400#	240#				1.19 s 0.52	5/2 <sup>-</sup> #	99 04Ga29	TD 2004	$\alpha = ?$	*
<sup>265</sup> Hs	120900	24				1.96 ms 0.16	3/2 <sup>+</sup> #	99 09He20	T 1984	$\alpha \approx 100$ ; SF ?	*
<sup>265</sup> Hs <sup>m</sup>	121130	24	229	22	AD	360 $\mu$ s 150	11/2 <sup>-</sup> #	99 09He20	T 1995	$\alpha \approx 100$ ; IT ?	*
<sup>265</sup> Mt	126620#	440#				2# ms				$\alpha$ ?	
* <sup>265</sup> Rf	T : average 17Og01, 15Uf02=1.0(+1.2-0.3) 16Ho09=61(+84-22)s; other										**
* <sup>265</sup> Rf	T : 10El06=105(+503-48) s, one SF at 152 s										**
* <sup>265</sup> Sg	T : average 12Ha05=8.5(+2.6,-1.6) 08Du09=8.9(+2.7-1.9)										**
* <sup>265</sup> Sg <sup>m</sup>	T : average 13Su04=20(+15-6) 12Ha05=14.4(+3.7,-2.5) 08Du09=16.2(+4.7-3.5);										**
* <sup>265</sup> Sg <sup>m</sup>	T : others 08Dv02=15(+7-4) 06Dv01=14.9(+9.1-4.1) 98Tu01=7.4(+3.3-2.7)										**
* <sup>265</sup> Sg <sup>m</sup>	T : 08Mo09 2 events at 23 and 80 s										**
* <sup>265</sup> Bh	T : symmetrized from 04Ga29=0.94(+0.70-0.31)										**
* <sup>265</sup> Hs	T : average 09He20=1.9(0.2) 99He11=2.0(+0.3-0.2)										**
* <sup>265</sup> Hs <sup>m</sup>	T : symmetrized from 09He20=300(+200-100); other 99He11=750(+170-120)										**
<sup>266</sup> Lr	111660#	540#				22 h 14		19 19Kh04	TD 2014	SF=100	*
<sup>266</sup> Rf	110140#	410#				4# h	0 <sup>+</sup>			$\alpha$ ?; SF ?	
<sup>266</sup> Db	112740#	280#				80 m 70		19 17Og01	T 2007	$\alpha$ ?; SF=?; $\beta^+$ ?	*
<sup>266</sup> Sg	113620#	250#				390 ms 110	0 <sup>+</sup>	19 17He08	D 2006	SF>90	*
<sup>266</sup> Bh	118100#	160#				10.6 s 2.2		19 20Ha27	T 2000	$\alpha \approx 100$ ; $\beta^+$ ?; SF ?	*
<sup>266</sup> Hs	121140	27				3.0 ms 0.6	0 <sup>+</sup>	19 12Ac04	TD 2001	$\alpha = 76$ 9; SF=24 9	*
<sup>266</sup> Hs <sup>m</sup>	122240	90	1100	90	AD	280 ms 220	9 <sup>-</sup> #	12Ac04	T 2011	$\alpha \approx 100$	*
<sup>266</sup> Mt	127670	100				2.0 ms 0.5		19 09Ne02	T 1982	$\alpha \approx 100$ ; SF ?	*
<sup>266</sup> Mt <sup>m</sup>	128810	120	1140	90	AD	6 ms 3		97Ho14	TD 1984	$\alpha = 100$	*
* <sup>266</sup> Lr	T : symmetrized from 19Kh04, 14Kh04=11(+21-5)										**
* <sup>266</sup> Db	T : symmetrized from 17Og01, 07Og02=22(+105-10), one event at 31.74 m										**
* <sup>266</sup> Sg	T : average 13Og03=280(+190-80) 08Dv02=360(+250-100), supersedes										**
* <sup>266</sup> Sg	T : 06Dv01=444(+444-148); others 98Tu01=21(+20-12) s 94La22=10-30 s										**
* <sup>266</sup> Bh	T : symmetrized from 20Ha27=10.0(+2.6-1.7); others: 15Mo25=2.2(+2.9-0.8)										**
* <sup>266</sup> Bh	T : 06Qi03=0.66(+0.59-0.26)										**
* <sup>266</sup> Hs	T : average 11Ac.A=2.97(+0.78-0.51) 01Ho06=2.3(+1.3-0.6)										**
* <sup>266</sup> Hs <sup>m</sup>	T : symmetrized from 12Ac04=74(+354-34); the possibility in 01Ho06 that										**
* <sup>266</sup> Hs <sup>m</sup>	T : 01Ho06=6.3(+8.6-2.3) is ruled out by the 12Ac04 data										**
* <sup>266</sup> Hs <sup>m</sup>	J : from 15Ko14, expected conf=n <sup>2</sup> (7/2[613], 11/2[735])										**
* <sup>266</sup> Mt	T : average 09Ne02=3.3(+2.5-1.0) 97Ho14=1.7(+0.6-0.4)										**
* <sup>266</sup> Mt <sup>m</sup>	T : symmetrized from 97Ho14=3.4(+4.7-1.3), 3 events at 7.8, 2.0 and 5.0 ms										**
<sup>267</sup> Rf	113440#	580#				2.5 h 1.5		05 17Og01	TD 2004	SF=100	*
<sup>267</sup> Rf <sup>p</sup>	113520#	580#	80#	100#							
<sup>267</sup> Db	114010#	370#				2.0 h 1.1	9/2 <sup>+</sup> #	05 17Og01	TD 2004	SF=100	*
<sup>267</sup> Sg	115810#	260#				1.8 m 0.7		08Dv02	TD 2008	SF=83; $\alpha = 17$	*
<sup>267</sup> Sg <sup>p</sup>	115830#	270#	20#	50#							

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>267</sup> Bh	118770#	260#				22 s 10	5/2 <sup>-</sup> #	05	2000	$\alpha=100$	*
<sup>267</sup> Hs	122660#	100#			*	55 ms 11		05 04Mo40 T	1995	$\alpha>80$ ;SF ?	*
<sup>267</sup> Hs <sup>m</sup>	122700#	100#	39	24	AD*	990 $\mu$ s 90		05 04Fo08 TD	2004	$\alpha=?$ ;IT ?	*
<sup>267</sup> Mt	127790#	500#				10# ms				$\alpha ?$	*
<sup>267</sup> Ds	133880#	200#				10 $\mu$ s 8	3/2 <sup>+</sup> #	05 95Gh04 T	1995	$\alpha=100$	*
* <sup>267</sup> Rf	T : symmetrized from 17Og01,06Og05=1.3(+2.3-0.5), supersedes 04Og12 one										**
* <sup>267</sup> Rf	T : event at 2.3 h										**
* <sup>267</sup> Db	T : symmetrized from 17Og01=1.3(+1.6-0.5)										**
* <sup>267</sup> Sg	T : symmetrized from 08Dv02=80(+60-20) s; other 99Og.B=19 ms not trusted										**
* <sup>267</sup> Bh	T : symmetrized from 00Wi15=17(+14-6); other 00Ei05=14(+9-4)										**
* <sup>267</sup> Hs	T : symmetrized from 04Mo40=52(+13-8), combining the 04Mo40 ( $\tau=77(+31-7)$ )										**
* <sup>267</sup> Hs	T : and 98Ho13 ( $\tau=72(+28-16)$ ) data										**
* <sup>267</sup> Hs <sup>m</sup>	T : 04Fo08(2 events)=940(+120-45)us; other 04Mo40(1 event)=0.80(+3.8-0.37)s										**
* <sup>267</sup> Ds	T : 95Gh04=2.8(+13.0-1.3), one event with $\tau=4$ us										**
<sup>268</sup> Rf	115480#	660#				1# h	0 <sup>+</sup>			$\alpha ?$ ;SF ?	*
<sup>268</sup> Db	117060#	530#				29 h 3		19 17Og01 T	2004	SF $\approx$ 100; $\beta^+$ ?; $\alpha ?$	*
<sup>268</sup> Db <sup>p</sup>	117210#	540#	150	80							*
<sup>268</sup> Sg	116800#	470#				2# m	0 <sup>+</sup>			$\alpha ?$ ;SF ?	*
<sup>268</sup> Bh	120710#	380#				190# s				$\alpha ?$ ;SF ?	*
<sup>268</sup> Hs	122970#	300#				1.4 s 1.1	0 <sup>+</sup>	10Ni14 TD	2010	$\alpha\approx$ 100	*
<sup>268</sup> Mt	129150#	230#				23 ms 7		19 04Mo26 T	1995	$\alpha=100$	*
<sup>268</sup> Ds	133650#	300#				100# $\mu$ s	0 <sup>+</sup>			$\alpha ?$	*
* <sup>268</sup> Db	T : symmetrized from 17Og01=28.3(+3.3-2.6); others 16Fo10=28(3)										**
* <sup>268</sup> Db	T : 13Ru11=26(+7-5) 13Og01=25.9(+6.2-4.2),supersedes 12Og02=27.9(+7.8-5.0)										**
* <sup>268</sup> Db	T : 05Og02=29(+9-6) 04Og03=16(+19-6), 07St18=28(+11-4)										**
* <sup>268</sup> Hs	T : symmetrized from 10Ni14=0.38(+1.8-0.17)										**
* <sup>268</sup> Mt	T : symmetrized from 04Mo26=21(+8-5), 14 events; other 02Ho11=42(+29,-12),										**
* <sup>268</sup> Mt	T : 6 events										**
<sup>269</sup> Db	119150#	620#				3# h	9/2 <sup>+</sup> #			$\alpha ?$ ;SF ?	*
<sup>269</sup> Sg	119690#	370#				5 m 2		19 17Og01 T	2010	$\alpha\approx$ 100;SF ?	*
<sup>269</sup> Bh	121480#	370#				1# m	5/2 <sup>-</sup> #			$\alpha ?$	*
<sup>269</sup> Hs	124490#	130#				15 s 7	9/2 <sup>+</sup> #	05 15Mo25 T	1996	$\alpha=100$	*
<sup>269</sup> Mt	129300#	310#				100# ms				$\alpha ?$	*
<sup>269</sup> Ds	134830	30				230 $\mu$ s 110		05 95Ho03 T	1995	$\alpha=100$	*
* <sup>269</sup> Sg	T : average 17Og01,15Ut02=3.1(+3.7-1.1)m 16Ho09=185(+254-68)s; other										**
* <sup>269</sup> Sg	T : 10Ei06=128(+613-58) s, one alpha event at 185 s										**
* <sup>269</sup> Hs	T : symmetrized from 15Mo25,13Su04=12(+9-4)										**
* <sup>269</sup> Ds	T : symmetrized from 95Ho03=170(+160-60)										**
<sup>270</sup> Db	122400#	580#				1.7 h 1.0		19 14Kh04 TD	2010	SF $\approx$ 87; $\alpha\approx$ 13	*
<sup>270</sup> Sg	121430#	460#				3# m	0 <sup>+</sup>			$\alpha ?$ ;SF ?	*
<sup>270</sup> Bh	124230#	300#				3.8 m 3.0		19 17Og01 TD	2007	$\alpha=100$	*
<sup>270</sup> Bh <sup>p</sup>	124920#	360#	690#	200#							*
<sup>270</sup> Hs	125110#	250#				9 s 4	0 <sup>+</sup>	19 13Og03 TD	2003	$\alpha\approx$ 100;SF ?	*
<sup>270</sup> Mt	130710#	190#				800 ms 400		19 15Mo25 TD	2004	$\alpha\approx$ 100	*
<sup>270</sup> Ds	134680	40				205 $\mu$ s 48	0 <sup>+</sup>	19 12Ac04 TD	2001	$\alpha\approx$ 100;SF ?	*
<sup>270</sup> Ds <sup>m</sup>	136070	60	1390	60	AD	4.3 ms 1.2	10 <sup>-</sup> #	19 12Ac04 T	2001	$\alpha\approx$ 70;IT $\approx$ 30	*
* <sup>270</sup> Db	T : symmetrized from 14Kh04=1.0(+1.5-0.4), combines 14Kh04 and 13Og04 data;										**
* <sup>270</sup> Db	T : other 19Kh04,14Kh04=1.0(+1.9-0.4)										**
* <sup>270</sup> Bh	T : symmetrized from 17Og01=61(+292-28)										**
* <sup>270</sup> Hs	T : symmetrized from 13Og03=7.6(+4.9-2.2); other 03Tu05=3.6(+0.8-1.4)										**
* <sup>270</sup> Mt	T : symmetrized from 15Mo25=0.48(+0.66-0.18)s										**
* <sup>270</sup> Ds	T : average 12Ac04=200(+70-40) 01Ho06=100(+140-40)										**
* <sup>270</sup> Ds <sup>m</sup>	T : symmetrized from 12Ac04=3.9(+1.5-0.8); other 01Ho06=6.0(+8.2-2.2)										**
* <sup>270</sup> Ds <sup>m</sup>	J : from 15Ko14, expected conf= $m^2(9/2[615],11/2[725])$ ,K=10-										**
<sup>271</sup> Sg	124620#	590#				2.2 m 1.1		06 17Og01 TD	2004	$\alpha=42$ 23;SF=58 23	*

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)		Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)		
<sup>271</sup> Bh	125860#	380#			2.9 s 1.9		05 17Og01 TD	2013	$\alpha=100$	*	
<sup>271</sup> Hs	127690#	280#			10# s			2008	$\alpha ?;SF ?$		
<sup>271</sup> Mt	131100#	330#			400# ms				$\alpha ?$		
<sup>271</sup> Ds	135950#	100#			144 ms 53		05 16Ho09 TD	1998	SF=75; $\alpha=25$	*	
<sup>271</sup> Ds <sup>m</sup>	136020#	100#	68	27	AD*& 1.7 ms 0.4		05	1995	$\alpha=100$	*	
* <sup>271</sup> Sg	T : symmetrized from 17Og01=1.6(+1.5-0.5)										**
* <sup>271</sup> Bh	T : symmetrized from 17Og01=1.5(+2.8-0.6)										**
* <sup>271</sup> Ds	T : average 16Ho09=96(+96-32) 04Mo40=86(+117-22)										**
* <sup>271</sup> Ds <sup>m</sup>	T : symmetrized from 04Mo40=1.63(+0.44-0.29), combining the 04Mo40										**
* <sup>271</sup> Ds <sup>m</sup>	T : ( $\tau=2.9(+1.3-0.7)$ ) and 98Ho13 ( $\tau=1.8(+0.8-0.4)$ ) data										**
<sup>272</sup> Sg	126520#	690#			4# m	0 <sup>+</sup>			$\alpha ?;SF ?$		
<sup>272</sup> Bh	128790#	530#			11.3 s 1.8		19 16Fo10 T	2004	$\alpha \approx 100$	*	
<sup>272</sup> Hs	129000#	510#			10# s	0 <sup>+</sup>			$\alpha ?;SF ?$		
<sup>272</sup> Mt	133480#	490#			400# ms				$\alpha ?;SF ?$		
<sup>272</sup> Ds	136080#	420#			200# ms	0 <sup>+</sup>			SF ?		
<sup>272</sup> Rg	142770#	230#			4.2 ms 1.1		19 15Mo25 T	1995	$\alpha=100$	*	
* <sup>272</sup> Bh	T : symmetrized from 16Fo10=10.5(+1.5-1.1); other 17Og01=10.6(+1.6-1.1),										**
* <sup>272</sup> Bh	T : same raw data as 16Fo10										**
* <sup>272</sup> Rg	T : symmetrized from 15Mo25=3.8(+1.4-0.8); other: 02Ho11=1.6(+1.1-0.5)										**
<sup>273</sup> Sg	129920#	400#			5# m				SF ?		
<sup>273</sup> Bh	130680#	660#			1# m				$\alpha ?;SF ?$		
<sup>273</sup> Hs	131770#	370#			1060 ms 500		15 17Og01 T	2010	$\alpha \approx 100; SF ?$	*	
<sup>273</sup> Hs <sup>p</sup>	131970#	390#	200#	100#					$\alpha ?;SF ?$		
<sup>273</sup> Mt	134780#	420#			800# ms				$\alpha ?;SF ?$		
<sup>273</sup> Ds	138290#	140#			240 $\mu$ s 100		05 15Mo25 T	1996	$\alpha \approx 100$	*	
<sup>273</sup> Ds <sup>m</sup>	138490#	140#	198	20	EU 120 ms		05	1996	$\alpha=100$		
<sup>273</sup> Rg	142890#	400#			2# ms				$\alpha ?$		
* <sup>273</sup> Hs	T : symmetrized from 17Og01,15Ut02=760(+710-240); other 16Ho09=765(+765-255)										**
* <sup>273</sup> Ds	T : symmetrized 15Mo25,13Su04=190(+140-60)										**
<sup>274</sup> Bh	133760#	580#			57 s 27		19 17Og01 TD	2010	$\alpha=100$	*	
<sup>274</sup> Hs	133410#	470#			500# ms	0 <sup>+</sup>			$\alpha ?;SF ?$		
<sup>274</sup> Mt	137250#	380#			850 ms 540		19 17Og01 TD	2007	$\alpha=100$	*	
<sup>274</sup> Ds	139200#	390#			10# ms	0 <sup>+</sup>			$\alpha ?;SF ?$		
<sup>274</sup> Rg	144610#	210#			20 ms 11		05 15Mo25 TD	2004	$\alpha \approx 100$	*	
* <sup>274</sup> Bh	T : symmetrized from 17Og01=44(+34-13) (recommended), based on										**
* <sup>274</sup> Bh	T : 14Kh04=30(+54-12) 13Og04=54(+65-19)										**
* <sup>274</sup> Mt	T : symmetrized from 17Og01=440(+810-170)ms, based on data from 07Og02										**
* <sup>274</sup> Rg	T : symmetrized from $\tau=18(+24-7)$ in 15Mo25										**
<sup>275</sup> Bh	135780#	600#			1# m	5/2 <sup>-</sup> #			SF ?		
<sup>275</sup> Hs	136490#	590#			280 ms 130		05 17Og01 TD	2004	$\alpha=100$	*	
<sup>275</sup> Hs <sup>p</sup>	136750#	600#	260#	100#							
<sup>275</sup> Mt	138770#	390#			31 ms 17		05 17Og01 TD	2004	$\alpha=100$	*	
<sup>275</sup> Ds	141670#	340#			10# ms				$\alpha ?;SF ?$		
<sup>275</sup> Rg	145400#	450#			5# ms				$\alpha ?$		
* <sup>275</sup> Hs	T : symmetrized 17Og01=200(+180-60); other 16Ho09=201(+201-67)										**
* <sup>275</sup> Mt	T : symmetrized from 17Og01=20(+24-7)										**
<sup>276</sup> Bh	138950#	600#			60# s				$\alpha ?;SF ?$		
<sup>276</sup> Hs	138190#	720#			100# ms	0 <sup>+</sup>			$\alpha ?;SF ?$		
<sup>276</sup> Mt	141310#	530#			700 ms 80		19 16Fo10 T	2004	$\alpha=100$	*	
<sup>276</sup> Mt <sup>m</sup>	141560#	540#	250	80	AD* 7 s 3		19 17Og01 TD	2012	$\alpha=100$	*	
<sup>276</sup> Ds	142540#	550#			16# ms	0 <sup>+</sup>			$\alpha ?;SF ?$		
<sup>276</sup> Rg	147390#	630#			10# ms				$\alpha ?;SF ?$		
<sup>276</sup> Cn	150360#	500#			100# $\mu$ s	0 <sup>+</sup>			$\alpha ?;SF ?$		



**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)
* <sup>276</sup> Mt	T : symmetrized from 16Fo10=690(+90-70); other 17Og01=520(100)						**
* <sup>276</sup> Mt <sup>m</sup>	T : symmetrized from 17Og01=4(+5-1)						**
<sup>277</sup> Bh	141100#	600#	10# s				$\alpha$ ?;SF ?
<sup>277</sup> Hs	141380#	450#	12 ms 9	*	14 16Ho09	TD 2010	SF≈100; $\alpha$ ?
<sup>277</sup> Hs <sup>m</sup>	141480#	460#	100# 100#	*	14 12Ho12	TD 2012	SF=100
<sup>277</sup> Hs <sup>p</sup>	142000#	490#	620# 200#				
<sup>277</sup> Mt	143010#	660#	9 s 6		14 17Og01	TD 2013	SF=100; $\alpha$ ?
<sup>277</sup> Ds	145090#	390#	6 ms 3		15 17Og01	T 2010	$\alpha$ ≈100;SF ?
<sup>277</sup> Rg	148410#	470#	4# ms				$\alpha$ ?;SF ?
<sup>277</sup> Cn	152330#	150#	790 $\mu$ s 330		05 15Mo25	TD 1996	$\alpha$ =100
* <sup>277</sup> Hs	T : symmetrized from 16Ho09=3.1(+14.9-1.4) 10Du06=3.0(+14.4-1.4)						**
* <sup>277</sup> Hs	T : 17Og01=3(+15-1); other 99Og10 one SF event at 16.5m, not trusted						**
* <sup>277</sup> Hs <sup>m</sup>	T : symmetrized from 12Ho12=34(+166-16) (SF 1 event)						**
* <sup>277</sup> Mt	T : symmetrized from 17Og01,13Og04=5(+9-2)						**
* <sup>277</sup> Ds	T : symmetrized from 17Og01,15U02=4.1(+3.7-1.3); other 16Ho09=4.1(+4.1-1.4)						**
* <sup>277</sup> Cn	T : symmetrized from 15Mo25,13Su04=610(+460-180)						**
<sup>278</sup> Bh	144370#	400#	2# s		19 16Ho09	TD 2016	SF≈100; $\alpha$ ?
<sup>278</sup> Hs	143220#	300#	2# s	0 <sup>+</sup>	19 16Ho09	TD 2016	SF≈100; $\alpha$ ?
<sup>278</sup> Mt	145770#	580#	6 s 3		19 17Og01	TD 2010	$\alpha$ =100
<sup>278</sup> Mt <sup>p</sup>	146160#	590#	390# 100#				
<sup>278</sup> Ds	146250#	510#	270# ms	0 <sup>+</sup>			$\alpha$ ?;SF ?
<sup>278</sup> Rg	150520#	390#	8 ms 5		19 17Og01	TD 2007	$\alpha$ =100
<sup>278</sup> Cn	152840#	440#	2# ms	0 <sup>+</sup>			$\alpha$ ?;SF ?
<sup>278</sup> Nh	159030#	220#	2.3 ms 1.3		19 15Mo25	TD 2004	$\alpha$ ≈100
* <sup>278</sup> Bh	T : 16Ho09=690(+3300,-310)s not trusted by evaluator, based on TNN						**
* <sup>278</sup> Hs	T : 16Ho09=690(+3300,-310)s not trusted by evaluator, based on TNN						**
* <sup>278</sup> Mt	T : symmetrized from 17Og01=4.5(+3.5-1.3) (recommended), based on data from						**
* <sup>278</sup> Mt	T : 14Kh04=3.6(+6.5-1.4) 13Og04=5.2(+6.2-1.8)						**
* <sup>278</sup> Rg	T : symmetrized from 17Og01,07Og02=4.2(+7.5-1.7)						**
* <sup>278</sup> Nh	T : symmetrized from $\tau$ =2.0(+2.7-0.7) in 15Mo25,08Mo09						**
<sup>279</sup> Hs	146500#	600#	1# s				$\alpha$ ?;SF ?
<sup>279</sup> Mt	147590#	670#	20# s				$\alpha$ ?;SF ?
<sup>279</sup> Ds	149020#	610#	210 ms 40		05 17Og01	TD 2004	SF=88 5; $\alpha$ =12 5
<sup>279</sup> Ds <sup>p</sup>	149250#	610#	230# 100#				
<sup>279</sup> Rg	151720#	420#	170 ms 110		05 17Og01	TD 2004	$\alpha$ =100
<sup>279</sup> Rg <sup>p</sup>	151760#	430#	40# 100#				
<sup>279</sup> Cn	155020#	400#	60# $\mu$ s				$\alpha$ ?;SF ?
<sup>279</sup> Nh	159460#	600#	1# ms				$\alpha$ ?;SF ?
* <sup>279</sup> Ds	T : from 17Og01=210(40); other 16Ho09=290(+69-47)						**
* <sup>279</sup> Ds	D : %SF symmetrized from 17Og01=89(+4-6)						**
* <sup>279</sup> Rg	T : symmetrized from 17Og01=90(+170-40)						**
<sup>280</sup> Hs	148420#	600#	100# ms	0 <sup>+</sup>			$\alpha$ ?;SF ?
<sup>280</sup> Mt	150510#	600#	10# s				$\alpha$ ?;SF ?
<sup>280</sup> Ds	150320#	750#	25 ms 20	0 <sup>+</sup>	19 17Ka66	TDI 1999	SF=100
<sup>280</sup> Rg	153890#	530#	4.3 s 0.5		19 17Og01	T 2004	$\alpha$ =100
<sup>280</sup> Cn	155650#	580#	5# ms	0 <sup>+</sup>			$\alpha$ ?;SF ?
<sup>280</sup> Nh	161240#	400#	10# ms				$\alpha$ ?;SF ?
* <sup>280</sup> Ds	I : the identification in 17Ka66 is tentative and it needs to be confirmed						**
* <sup>280</sup> Ds	T : symmetrized from 17Ka66=6.7(+31.9-3); others 01Og01-3 events at 6.93 s,						**
* <sup>280</sup> Ds	T : 14.3 s and 7.4 s yield 6.6(+9.0-2.4) s, but data were later reassigned						**
* <sup>280</sup> Ds	T : to the <sup>293</sup> Lv chain						**
* <sup>280</sup> Rg	T : symmetrized from 17Og01=4.2(+0.6,-0.4); other 16Fo10=4.4(+0.5-0.4)						**
<sup>281</sup> Mt	152400#	600#	1# s				$\alpha$ ?;SF ?

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)		Excitation Energy (keV)			Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>281</sup> Ds	153270#	490#			*	14 s 3		05 17Ka66 T	2004	SF=90.7;α=10.7	*
<sup>281</sup> Ds <sup>m</sup>	153350#	460#	80#	240#	*	0.9 s 0.7		12Ho12 TD	2012	α=100	*
<sup>281</sup> Ds <sup>p</sup>	153340#	500#	70#	100#							
<sup>281</sup> Rg	155330#	770#				19 s 5		10 17Og01 TD	2010	SF=87.8;α=13.8	*
<sup>281</sup> Cn	157950#	400#				180 ms 80		15 17Og01 T	2010	α≈100;SF?	*
<sup>281</sup> Nh	161810#	300#				100# ms				α?;SF?	
* <sup>281</sup> Ds	T : average 17Ka66=9.9(+13.6-3.6) 17Og01=12.7(+4.0-2.5);										**
* <sup>281</sup> Ds	T : other 16Ho09=13.0(+4.5-2.7)										**
* <sup>281</sup> Ds	D : %SF symmetrized from 17Og01=93(+5-9)										**
* <sup>281</sup> Ds <sup>m</sup>	T : symmetrized from 12Ho12=0.25(+1.18-0.11) s										**
* <sup>281</sup> Rg	T : symmetrized from 17Og01=17(+6-3); other 16Fo16=21(+10-5)										**
* <sup>281</sup> Rg	D : %SF symmetrized from 17Og01=88(+7-9)										**
* <sup>281</sup> Cn	T : symmetrized from 17Og01,15Ut02=130(+120-40); other:										**
* <sup>281</sup> Cn	T : 16Ho09=128(+128-43) (analyzing same data as 17Og01)										**
<sup>282</sup> Mt	155460#	450#				100# ms		19 16Ho09 TD	2016	α≈100; SF?	*
<sup>282</sup> Ds	154790#	300#				4.2 m 3.3	0 <sup>+</sup>	19 16Ho09 TD	2016	α≈100; SF?	*
<sup>282</sup> Rg	157740#	590#				130 s 50		19 17Og01 TD	2010	α=100	*
<sup>282</sup> Cn	158830#	550#				1.1 ms 0.3	0 <sup>+</sup>	19 16Ho09 TD	2004	SF≈100;α?	*
<sup>282</sup> Nh	163730#	400#				140 ms 90		19 17Og01 TD	2007	α=100	*
* <sup>282</sup> Mt	T : 16Ho09=67(+320-30)s not trusted by evaluator, based on TNN										**
* <sup>282</sup> Ds	T : symmetrized from 16Ho09=67(+320-30)s										**
* <sup>282</sup> Rg	T : symmetrized from 17Og01=100(+70-30)										**
* <sup>282</sup> Cn	T : symmetrized from 16Ho09=0.96(0.35-0.20); other: 17Og01=0.91(0.33-0.19)										**
* <sup>282</sup> Nh	T : symmetrized from 17Og01,07Og02=73(+134-29)										**
<sup>283</sup> Ds	157830#	500#				1# m				α?;SF?	
<sup>283</sup> Rg	159380#	680#				2# m				α?;SF?	
<sup>283</sup> Cn	161340#	620#				4.7 s 0.8		06 16Ho09 TD	2004	α=81;SF=19	*
<sup>283</sup> Nh	164560#	440#				140 ms 90		05 17Og01 TD	2004	α=100	*
* <sup>283</sup> Cn	T : symmetrized from 16Ho09=4.48(+0.98-0.68); other 17Og01=4.2(+1.1-0.7)										**
* <sup>283</sup> Nh	T : symmetrized from 17Og01=75(+136-30)										**
<sup>284</sup> Ds	159460#	500#				1# m	0 <sup>+</sup>			α?;SF?	
<sup>284</sup> Rg	161970#	500#				1# m				α?;SF?	
<sup>284</sup> Cn	162420#	760#				102 ms 17	0 <sup>+</sup>	19 17Og01 TD	2004	SF=100	*
<sup>284</sup> Nh	166590#	530#				0.97 s 0.11		19 17Og01 TD	2004	α=100	*
<sup>284</sup> Fl	168780#	660#				3.1 ms 1.3	0 <sup>+</sup>	19 17Og01 TD	2015	SF≈100;α?	*
* <sup>284</sup> Cn	T : symmetrized from 17Og01=98(20-14); other 16Ho09=118(+24-17)										**
* <sup>284</sup> Nh	T : symmetrized from 17Og01=0.97(0.12-0.10); other 16Fo10=0.97(0.12-0.10)										**
* <sup>284</sup> Fl	T : symmetrized from 17Og01=2.5(+1.8-0.8); other 16Ho09=2.0(+2.7,-0.7)										**
<sup>285</sup> Rg	163730#	600#				30# s				α?;SF?	
<sup>285</sup> Cn	165090#	510#			*	30 s 8		05 17Og01 TD	2004	α=100	*
<sup>285</sup> Cn <sup>m</sup>	165620#	460#	530#	270#	*	15 s 12		12Ho12 TD	2012	α=100	*
<sup>285</sup> Nh	167770#	780#				4.6 s 1.1		10 17Og01 TD	2010	α=100	*
<sup>285</sup> Fl	170930#	400#				210 ms 100		15 17Og01 T	2010	α≈100;SF<20	*
* <sup>285</sup> Cn	T : symmetrized from 17Og01=28(+9-6); other 16Ho09=28.9(+10.1-5.9)										**
* <sup>285</sup> Cn <sup>m</sup>	T : symmetrized from 12Ho12=4.0(+19.1-1.8) s										**
* <sup>285</sup> Nh	T : symmetrized from 07Og01=4.2(+1.4-0.8); other 16Fo16=2.9(+1.4-0.7),										**
* <sup>285</sup> Nh	T : reanalyzed data of 13Og04=4.2(+1.4-0.8), 12Og06=4.9(+6.7-1.8),										**
* <sup>285</sup> Nh	T : 10Og01=5.5(+5.0-1.8)										**
* <sup>285</sup> Fl	T : symmetrized from 17Og01,15Ut02=150(+140-50); other: 16Ho09=152(+152-51),										**
* <sup>285</sup> Fl	T : analyzed same data as 17Og01										**
<sup>286</sup> Rg	166510#	460#				10# s		19 16Ho09 TD	2016	α≈100;SF?	*
<sup>286</sup> Cn	166450#	700#				30 s 30	0 <sup>+</sup>	19 17Ka66 T	2016	α≈100;SF?	*
<sup>286</sup> Nh	169960#	590#				12 s 5		19 17Og01 TD	2010	α=100	*

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
<sup>286</sup> Fl	171610#	550#	130 ms 30	0 <sup>+</sup>	19 17Og01	TD 2004	$\alpha=59$ 11;SF=41 11	*
* <sup>286</sup> Rg	T : 16Ho09=640(+3100-300)s not trusted by evaluator, based on TNN							**
* <sup>286</sup> Cn	T : symmetrized from 17Ka66=8.4(+40.5-3.9); other: 16Ho09=640(+3100-300)s							**
* <sup>286</sup> Nh	T : symmetrized from 17Og01=9.5(+6.3-2.7) (recommended), based on data from							**
* <sup>286</sup> Nh	T : 14Kh04=2.9(+5.5-1.1) 13Og04=13(+12-4)							**
* <sup>286</sup> Fl	T : symmetrized from 17Og01=120(+40-20); other: 16Ho09=166(+40-27)							**
* <sup>286</sup> Fl	D : % $\alpha$ symmetrized from 17Og01=60(+10-11); other: 16Ho09=52%							**
<sup>287</sup> Cn	169370#	700#	30# s				$\alpha$ ?;SF ?	
<sup>287</sup> Nh	171460#	710#	20# s				$\alpha$ ?;SF ?	
<sup>287</sup> Fl	173930#	620#	510 ms 120		05 17Og01	T 2004	$\alpha\approx 100$ ; SF ?	*
<sup>287</sup> Mc	177750#	440#	60 ms 30		05 17Og01	TD 2004	$\alpha=100$	*
* <sup>287</sup> Fl	T : symmetrized from 17Og01=480(+140-90); other: 16Ho09=540(+170-100)							**
* <sup>287</sup> Mc	T : symmetrized from 17Og01=37(+44-13)							**
<sup>288</sup> Cn	170930#	700#	10# s	0 <sup>+</sup>			$\alpha$ ?;SF ?	
<sup>288</sup> Nh	173970#	700#	20# s				$\alpha$ ?;SF ?	
<sup>288</sup> Fl	174920#	760#	653 ms 113	0 <sup>+</sup>	19 16Ho09	TD 2004	$\alpha\approx 100$ ; SF ?	*
<sup>288</sup> Mc	179670#	540#	177 ms 20		19 17Og01	TD 2004	$\alpha=100$	*
* <sup>288</sup> Fl	T : average 16Ho09=644(+136-97) 17Ka66=274(+500-108);							**
* <sup>288</sup> Fl	T : other: 17Og01=660(+140-100)							**
* <sup>288</sup> Mc	T : symmetrized from 17Og01=174(+22-18); other 16Fo10=170(20)							**
<sup>289</sup> Nh	175550#	500#	30# s				$\alpha$ ?;SF ?	
<sup>289</sup> Fl	177470#	510#	2.1 s 0.6		05 17Og01	TD 2004	$\alpha\approx 100$ ;SF ?	*
<sup>289</sup> Fl <sup>m</sup>	178220#	470#	750# 280#	1.1 s 0.8	12Ho12	TD 2012	$\alpha=100$	*
<sup>289</sup> Mc	180680#	780#	410 ms 150		10 17Og01	T 2010	$\alpha=100$	*
<sup>289</sup> Lv	184460#	500#	RN	16# ms	00 02Ni10	I	$\alpha$ ?	*
* <sup>289</sup> Fl	T : symmetrized from 17Og01=1.9(+0.7-0.4); others 16Ho09=1.87(+0.65-0.38)							**
* <sup>289</sup> Fl	T : 17Ka66=3.9(+5.3-1.4)							**
* <sup>289</sup> Fl <sup>m</sup>	T : symmetrized from 12Ho12=0.28(+1.35-0.13) s							**
* <sup>289</sup> Mc	T : symmetrized from 17Og01=330(+120-80); other 16Fo16=270(+120-60),							**
* <sup>289</sup> Mc	T : reanalyzed data of 13Og04=330(+120-80), 12Og06=430(+590-160),							**
* <sup>289</sup> Mc	T : 10Og01=220(+260-80)							**
* <sup>289</sup> Lv	T : 99Ni03=600(+860-300), $\alpha$ decay retracted by authors in 02Ni10							**
<sup>290</sup> Nh	178320#	470#	8 s 6		19 16Ho09	TD 2016	$\alpha\approx 100$ ; SF $\leq 50$	*
<sup>290</sup> Fl	178730#	700#	80 s 60	0 <sup>+</sup>	19 16Ho09	TD 2016	$\alpha\approx 100$ ; SF ?; $\beta^+ < 50$	*
<sup>290</sup> Mc	182790#	590#	840 ms 360		19 17Og01	T 2010	$\alpha=100$	*
<sup>290</sup> Lv	185030#	550#	9 ms 3	0 <sup>+</sup>	19 17Og01	T 2004	$\alpha\approx 100$ ; SF ?	*
* <sup>290</sup> Nh	T : symmetrized from 16Ho09=2.0(+9.6-0.9)							**
* <sup>290</sup> Fl	T : symmetrized from 16Ho09=21(+101-10); other 16Ho09=19(+91-9) $\beta^+$ branch							**
* <sup>290</sup> Mc	T : symmetrized from 17Og01=650(+490,-200) (recommended); based on data							**
* <sup>290</sup> Mc	T : from 13Og04=240(+280-90) 14Kh04=1300(+2300-500)							**
* <sup>290</sup> Lv	T : symmetrized from 17Og01=8.3(+3.5-1.9); other 16Ho09=8.3(+3.6-1.9)							**
<sup>291</sup> Fl	181500#	700#	10# s				$\alpha$ ?;SF ?	
<sup>291</sup> Mc	184180#	740#	1# s				$\alpha$ ?;SF ?	
<sup>291</sup> Lv	187240#	620#	26 ms 12		05 17Og01	T 2004	$\alpha\approx 100$ ;SF ?	*
<sup>291</sup> Ts	191650#	600#	2# ms				$\alpha$ ?;SF ?	
* <sup>291</sup> Lv	T : symmetrized from 17Og01=19(+17-6); other 16Ho09=18(+25-7)							**
<sup>292</sup> Mc	186600#	700#	5# s				$\alpha$ ?;SF ?	
<sup>292</sup> Lv	188130#	760#	16 ms 6	0 <sup>+</sup>	19 17Og01	T 2004	$\alpha\approx 100$ ; SF ?	*
<sup>292</sup> Ts	193620#	670#	10# ms				$\alpha$ ?;SF ?	
* <sup>292</sup> Lv	T : average 17Og01=13(+7-4) 17Ka66=11.9(+21.7-2.6);							**
* <sup>292</sup> Lv	T : other 16Ho09=12.8(+7.0-3.3)							**

**Table I. The NUBASE2020 table (Explanation of Table on page 030001-16)**

Nuclide	Mass excess (keV)	Excitation Energy (keV)	Half-life	$J^\pi$	Ens Reference	Year of discovery	Decay modes and intensities (%)	
$^{293}\text{Lv}$	190570#	520#	70 ms 30		05 17Og01	T 2004	$\alpha \approx 100$ ; SF ?	*
$^{293}\text{Lv}^m$	191290#	470#	80 ms 60		12Ho12	TD 2012	$\alpha = 100$	*
$^{293}\text{Ts}$	194430#	780#	25 ms 6		10 17Og01	TD 2010	$\alpha = 100$	*
$^{293}\text{Og}$	198800#	710#	1# ms		00 02Ni10	I 2010	$\alpha ?$	*
* $^{293}\text{Lv}$	T : symmetrized from 17Og01,15Og05=57(+43-17); others 16Ho09=57(+46-18)							**
* $^{293}\text{Lv}$	T : 17Ka66t=188(+342-74)							**
* $^{293}\text{Lv}^m$	T : symmetrized from 12Ho12=20(+96-9)							**
* $^{293}\text{Ts}$	T : symmetrized from 17Og01,15Og05=22(+8-4); other: 16Fo16=18(+8-4),							**
* $^{293}\text{Ts}$	T : reanalyzed data of 13Og04=22(+8-4), 12Og06=27(+12-6),							**
* $^{293}\text{Ts}$	T : 10Og01=14(+11-4)							**
* $^{293}\text{Og}$	T : 99Ni03=120(+180-60) $\alpha$ decay retracted by authors in 02Ni10							**
$^{294}\text{Ts}$	196400#	590#	70 ms 30		19 17Og01	TD 2010	$\alpha = 100$	*
$^{294}\text{Og}$	199320#	550#	0.7 ms 0.3	$0^+$	05 18Br13	T 2004	$\alpha \approx 100$ ; SF ?	*
* $^{294}\text{Ts}$	T : symmetrized from 17Og01,15Og05=51(+38-16) (recommended); based on data							**
* $^{294}\text{Ts}$	T : from 14Kh04=51(+94-20) 13Og04=50(+60-18)							**
* $^{294}\text{Og}$	T : symmetrized from 18Br13=0.58(+0.44-0.18), supersedes							**
* $^{294}\text{Og}$	T : 17Og01=0.69(+0.64-0.22) 16Ho09=0.69(+69-23)							**
$^{295}\text{Og}$	201370#	660#	680 ms 540		16Ho09	TD 2006	$\alpha \approx 100$	*
* $^{295}\text{Og}$	T : symmetrized from 16Ho09=181(+866-83)							**